

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 94 (2019)
Heft: 3

Artikel: Fourier optimization and prime gaps
Autor: Carneiro, Emanuel / Milinovich, Micah B. / Soundararajan, Kannan
DOI: <https://doi.org/10.5169/seals-846789>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Fourier optimization and prime gaps

Emanuel Carneiro, Micah B. Milinovich and Kannan Soundararajan

Abstract. We investigate some extremal problems in Fourier analysis and their connection to a problem in prime number theory. In particular, we improve the current bounds for the largest possible gap between consecutive primes assuming the Riemann hypothesis.

Mathematics Subject Classification (2010). 41A30, 11M06, 11M26, 11N05.

Keywords. Bandlimited functions, Fourier uncertainty, prime gaps, Riemann hypothesis.

1. Introduction

In this paper we study a new set of extremal problems in Fourier analysis, motivated by a problem in prime number theory. These problems (which will be described shortly) are of the kind where one prescribes some constraints for a function and its Fourier transform, and then wants to optimize a certain quantity. When available, a solution to such a problem usually requires two main ingredients: a tool to prove optimality and a tool to construct an extremal function. A classical example in approximation theory is the problem of finding the best $L^1(\mathbb{R})$ -approximation of real-valued functions by bandlimited functions (i.e. functions with compactly supported Fourier transforms). For the two-sided problem (i.e. unrestricted approximation), one usually works with the so called *extremal signatures* to establish optimality, whereas for the one-sided problem (in which one is interested in majorizing or minorizing a given function) the Poisson summation formula is useful as a tool to prove optimality. For an account of such methods see, for instance, [10, 37, 42] and the references therein. Optimal bandlimited majorants and minorants have several applications to inequalities in analysis and number theory, for instance in connection to the theory of the Riemann zeta-function, e.g. [6–8, 11]. Slightly different extremal problems appear in the work [33], in connection with the question of bounding the least quadratic nonresidue modulo a prime. Another example of a Fourier optimization problem was proposed by Cohn and Elkies [12], in connection to the sphere packing problem. This recently attracted considerable attention with its resolution in dimensions 8 and 24 (see [13, 43]).

As we see below, the Fourier optimization problems considered here are simple enough to be stated in very accessible terms but rather delicate in the sense that the usual tools in the literature to prove optimality and construct extremal functions are not particularly helpful. While we have been unable to determine explicitly the solutions to our optimization problems, we are able to make progress on the existence and uniqueness of extremizers, and to establish good upper and lower bounds for the values of the sharp constants. In addition, we establish a connection between these extremal problems in Fourier analysis and the problem of bounding the largest possible gap between consecutive primes (assuming the Riemann hypothesis).

1.1. Fourier optimization problems. For $F \in L^1(\mathbb{R})$, we let

$$\widehat{F}(t) = \int_{-\infty}^{\infty} e^{-2\pi i x t} F(x) dx$$

denote the Fourier transform of F . We also let $x_+ := \max\{x, 0\}$ and $1 \leq A \leq \infty$ be a given parameter (note that we include the possibility that $A = \infty$), and we consider the following problems.

Extremal problem 1. Given $1 \leq A < \infty$, find

$$\mathcal{C}(A) := \sup_{\substack{F \in \mathcal{A} \\ F \neq 0}} \frac{1}{\|F\|_1} \left(|F(0)| - A \int_{[-1,1]^c} |\widehat{F}(t)| dt \right), \quad (1.1)$$

where the supremum is taken over the class \mathcal{A} of continuous functions $F: \mathbb{R} \rightarrow \mathbb{C}$, with $F \in L^1(\mathbb{R})$. In the case $A = \infty$, determine

$$\mathcal{C}(\infty) = \sup_{\substack{F \in \mathcal{E} \\ F \neq 0}} \frac{|F(0)|}{\|F\|_1}, \quad (1.2)$$

where the supremum is over the subclass $\mathcal{E} \subset \mathcal{A}$ of continuous functions $F: \mathbb{R} \rightarrow \mathbb{C}$, with $F \in L^1(\mathbb{R})$ and $\text{supp}(\widehat{F}) \subset [-1, 1]$.

Extremal problem 2. Given $1 \leq A < \infty$, find

$$\mathcal{C}^+(A) := \sup_{\substack{F \in \mathcal{A}^+ \\ F \neq 0}} \frac{1}{\|F\|_1} \left(F(0) - A \int_{[-1,1]^c} (\widehat{F}(t))_+ dt \right), \quad (1.3)$$

where the supremum is taken over the class \mathcal{A}^+ of even and continuous functions $F: \mathbb{R} \rightarrow \mathbb{R}$, with $F \in L^1(\mathbb{R})$. In the case $A = \infty$, determine

$$\mathcal{C}^+(\infty) = \sup_{\substack{F \in \mathcal{E}^+ \\ F \neq 0}} \frac{F(0)}{\|F\|_1}, \quad (1.4)$$

where the supremum is over the subclass $\mathcal{E}^+ \subset \mathcal{A}^+$ of even and continuous functions $F: \mathbb{R} \rightarrow \mathbb{R}$, with $F \in L^1(\mathbb{R})$ and $\widehat{F}(t) \leq 0$ for $|t| \geq 1$.

There has been some previous works in connection to problem (1.2) and its analogue for trigonometric polynomials, see for instance [2, 24, 29, 40]. The current best numerical upper and lower bounds for $\mathcal{C}(\infty)$, reviewed in (1.5) below, are due to Hörmander and Bernhardsson [29]. We were not able to find any mention to the other problems in the literature. *If one further imposes the condition that F is nonnegative on \mathbb{R}* , then (1.2) reduces to a folkloric problem for bandlimited functions while (1.4) reduces to the Cohn–Elkies problem [12, Theorem 3.1] in dimension 1. In both cases Poisson summation shows that the required maximum is 1, being attained by any constant multiple of the Fejér kernel $F(x) = (\sin(\pi x)/(\pi x))^2$. Classical interpolation formulas of Vaaler [42, Theorem 9] show that these are indeed the unique extremizers for this simplified version of (1.2), whereas this simplified version of (1.4) admits other extremizers (see [12, Section 5]).

We restricted the parameter A to the range $1 \leq A \leq \infty$ because in the range $0 < A < 1$ the corresponding problems (1.1) and (1.3) are trivial in the sense that $\mathcal{C}(A) = \mathcal{C}^+(A) = \infty$. This can be seen by taking $F_\varepsilon(x) = \frac{1}{\sqrt{\varepsilon}} e^{-\pi x^2/\varepsilon}$ with $\varepsilon \rightarrow 0^+$. It is also clear that the mappings $A \mapsto \mathcal{C}(A)$ and $A \mapsto \mathcal{C}^+(A)$ are non-increasing for $1 \leq A \leq \infty$.

The extremal problems presented here are certainly related to the phenomenon of Fourier uncertainty, and works like [17, 18], that discuss L^1 -uncertainty principles, provide interesting insights. The recent works [3, 23] on the “root-uncertainty principle” for the Fourier transform also consider interesting extremal problems related to the theory of zeta-functions in number fields. Toward the problems of determining the exact values of the sharp constants $\mathcal{C}(A)$ and $\mathcal{C}^+(A)$ we establish the following results.

Theorem 1.1. *Let $1 \leq A \leq \infty$. With respect to problems (1.1) and (1.2), the following propositions hold:*

(a) *If $A = \infty$, then:*

- (a.1) *There exists an even and real-valued function $G \in \mathcal{E}$, with $G(0) = 1$, that extremizes (1.2).*
- (a.2) *All the extremizers of (1.2) are of the form $F(x) = c G(x)$, where $c \in \mathbb{C}$ with $c \neq 0$.*
- (a.3) *The extremal function G verifies the identity*

$$\mathcal{C}(\infty) \int_{-\infty}^{\infty} \operatorname{sgn}(G(x)) F(x) dx = F(0)$$

for any $F \in \mathcal{E}$.

- (a.4) *(cf. [29]) The sharp constant $\mathcal{C}(\infty)$ verifies the inequality*

$$1.0818576438 < \mathcal{C}(\infty) < 1.0818576441. \quad (1.5)$$

(b) If $A = 1$ then $\mathcal{C}(1) = 2$, but there are no extremizers for (1.1).

(c) If $1 < A < \infty$, then:

(c.1) There exists an even and real-valued function $G \in \mathcal{A}$ that extremizes (1.1).

(c.2) Let $c_0 = \frac{4}{\pi} \left(\int_{-1}^1 \frac{\sin \pi t}{\pi t} dt \right)^{-1} = 1.07995 \dots$ and let $d_0 = 1.09769 \dots$ be defined by (4.5). Let $\lambda = \lambda(A)$ be the unique solution of

$$1 - \frac{1}{A} = \sin\left(\frac{\pi\lambda}{2}\right) - \frac{\pi\lambda}{2} \cos\left(\frac{\pi\lambda}{2}\right)$$

with $0 < \lambda < 1$. Then

$$\begin{aligned} \max \left\{ 2A - 2\sqrt{A(A-1)}, \frac{\pi A c_0}{2} \cos\left(\frac{\pi\lambda(A)}{2}\right) \right\} \\ \leq \mathcal{C}(A) \leq \min \left\{ \left(\frac{d_0}{1 - 0.3/(A-2)} \right), 2 \right\}, \end{aligned} \quad (1.6)$$

where the first upper bound on the right-hand side of (1.6) is only available in the range $2.6 \leq A < \infty$.

Remark. The function

$$H(x) = \frac{\cos 2\pi x}{1 - 16x^2} \quad (1.7)$$

belongs to the class \mathcal{E} and verifies $\|H\|_1 = 1/c_0$. We then have $H(0)/\|H\|_1 = c_0 = 1.07995 \dots$, and this yields a slightly inferior lower bound for $\mathcal{C}(\infty)$ when compared to the one in (1.5) (which is obtained in [29] by means of more complicated numerical examples). Due to its simplicity, this particular function $H(x)$ plays an important role in our work, being used in the proof of the lower bound in (1.6) and in the proof of Theorem 1.5.

Theorem 1.2. Let $1 \leq A \leq \infty$. With respect to problems (1.3) and (1.4), the following propositions hold:

(a) If $A = \infty$, then:

(a.1) There exists a function $G \in \mathcal{E}^+$ that extremizes (1.4).

(a.2) The sharp constant $\mathcal{C}^+(\infty)$ verifies the inequality

$$\mathcal{C}(\infty) \leq \mathcal{C}^+(\infty) < 1.2.$$

(b) If $A = 1$ then $\mathcal{C}^+(1) = 2$, but there are no extremizers for (1.3).

(c) If $1 < A < \infty$, then:

(c.1) There exists an even and real-valued function $G \in \mathcal{A}^+$ that extremizes (1.3).

(c.2) The sharp constant $\mathcal{C}^+(A)$ verifies the inequality

$$\mathcal{C}(A) \leq \mathcal{C}^+(A) \leq \min \left\{ \left(\frac{1.2}{1 - 0.222/(A-1)} \right), 2 \right\}, \quad (1.8)$$

where the first upper bound on the right-hand side of (1.8) is only available in the range $1.222 < A < \infty$.

(c.3) In particular, if $A = 36/11$ a numerical example yields the lower bound

$$\frac{25}{21} < \mathcal{C}^+\left(\frac{36}{11}\right). \quad (1.9)$$

Remark. Note that for small values of A , the right-hand side of (1.8) gives a better bound than the right-hand side of (1.6), and can be used instead. The reason, as we shall see, is that such bounds come from modifying the test functions in the dual problem for the case $A = \infty$. In our construction, these modifications do not necessarily maintain the hierarchy as A approaches 1.

1.2. Bounds for prime gaps on RH. Let p_n denote the n th prime. Assuming the Riemann hypothesis (RH), a classical result of Cramér [14] yields the bound

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\sqrt{p_n} \log p_n} \leq c, \quad (1.10)$$

where c is a universal constant. Building upon the works of Goldston [21] and of Ramaré and Saouter [39], the current best form of this bound is due to Dudek [19, Theorem 1.3], who obtained (1.10) with constant $c = 1$. Here we improve this and other bounds in this theory by establishing an interesting connection with the extremal problems presented in the previous section.

Our strategy consists of three main ingredients: (i) the explicit formula, (ii) the Brun–Titchmarsh inequality, and (iii) the derived extremal problems in Fourier analysis. Letting $\pi(x)$ denote the number of primes less than or equal to x , we define the Brun–Titchmarsh constant \mathbf{B} in our desired scale by

$$\mathbf{B} := \limsup_{x \rightarrow \infty} \frac{\pi(x + \sqrt{x}) - \pi(x)}{\sqrt{x} / \log x} \quad (1.11)$$

and we observe that

$$1 \leq \mathbf{B} \leq \frac{36}{11}. \quad (1.12)$$

The lower bound in (1.12) follows from the prime number theorem $\pi(x) \sim x / \log x$ as $x \rightarrow \infty$ and the upper bound on \mathbf{B} follows from the work of Iwaniec [30, Theorem 14].

We prove the following general result.

Theorem 1.3. *Assume the Riemann hypothesis. Let $\mathcal{C}^+(\cdot)$ be defined in (1.3) and \mathbf{B} be defined in (1.11). Then, for any $\alpha \geq 0$, we have*

$$\inf \left\{ c > 0; \liminf_{x \rightarrow \infty} \frac{\pi(x + c\sqrt{x} \log x) - \pi(x)}{\sqrt{x}} > \alpha \right\} \leq \frac{(1 + 2\alpha)}{\mathcal{C}^+(\mathbf{B})} < \frac{21}{25}(1 + 2\alpha). \quad (1.13)$$

The last inequality comes from (1.9) and (1.12).

The case $\alpha = 0$ in Theorem 1.3 yields an affirmative answer for a question posed in [19], on whether one could establish (1.10) with a constant $c < 1$.

Corollary 1.4. *Assume the Riemann hypothesis. Let $\mathcal{C}^+(\cdot)$ be defined in (1.3) and \mathbf{B} be defined in (1.11). Then*

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\sqrt{p_n} \log p_n} \leq \frac{1}{\mathcal{C}^+(\mathbf{B})} < \frac{21}{25}. \quad (1.14)$$

We note from (1.12) and Theorem 1.2 (b) that the limit of this method would yield a constant $\frac{1}{2}$ on the right-hand side of (1.14). On the other hand, under stronger assumptions, namely the Riemann hypothesis and Montgomery's pair correlation conjecture, it is known that the limit supremum in (1.14) is actually zero (see, for instance, [26, 27, 36]).

The case $\alpha = 1$ in Theorem 1.3 yields the constant

$$c = \frac{3}{\mathcal{C}^+(\mathbf{B})} < \frac{63}{25}$$

on the right-hand side of (1.13). This also sharpens the previous best result, due to Dudek [19], who had obtained this inequality with constant $c = 3$.

By working with a particular dilation of the bandlimited function (1.7) and an explicit version of the Brun–Titchmarsh inequality due to Montgomery and Vaughan [35], we are able to make all of our error terms effective and, assuming the Riemann hypothesis, prove that

$$p_{n+1} - p_n \leq \frac{22}{25} \sqrt{p_n} \log p_n$$

for all primes $p_n > 3$.

Theorem 1.5. *Assume the Riemann hypothesis. Then, for $x \geq 4$, there is always a prime number in the interval $[x, x + \frac{22}{25} \sqrt{x} \log x]$.*

This theorem improves a result of Dudek, Grenié, and Molteni [20, Theorem 1.1], who had previously reached a similar conclusion with $c = \frac{22}{25}$ replaced by $c = c(x) = 1 + \frac{4}{\log x}$. Cramér [15] has conjectured that

$$p_{n+1} - p_n = O(\log^2 p_n),$$

and this problem remains open to this date. It has been verified by Oliveira e Silva, Herzog, and Pardi [38, Section 2.2] that

$$p_{n+1} - p_n < \log^2 p_n \quad (1.15)$$

for all primes $11 \leq p_n \leq 4 \cdot 10^{18}$. Estimate (1.15) plainly implies the conclusion of Theorem 1.5 for all $4 \leq x \leq 4 \cdot 10^{18}$. Therefore, in our proof, we assume that $x \geq 4 \cdot 10^{18}$.

We now proceed to the proofs of the main results stated in this introduction. This is carried out in Sections 2–6. In Section 7 we have a general discussion on some related extremal problems in Fourier analysis, which includes for example the existence of extremizers for the Fourier optimization problem of Cohn and Elkies [12] related to sphere packing. Some of this material may be of independent interest.

2. Existence of extremizers

In this section we discuss the existence of extremizers for the extremal problems (1.1)–(1.4). We prove here parts (a.1), (b), and (c.1) of Theorems 1.1 and 1.2. We begin by making some simplifying observations, that will be helpful for the rest of the paper. Note that we may restrict ourselves to the situation when $\hat{F} \in L^1(\mathbb{R})$ (otherwise the quotients on right-hand sides of (1.1), (1.3), and (1.4) yield $-\infty$), and we assume this throughout the rest of the paper. In particular, F decays at infinity and $\|F\|_\infty$ is attained at some point.

The class \mathcal{A} in Theorem 1.1 includes complex-valued functions, but for our extremal problems we can restrict attention to even, real-valued functions. Indeed, given a non-identically zero $F \in \mathcal{A}$, the following steps either increase the quotients on the right-hand sides of (1.1)–(1.2) or leave them unaltered:

- by translating F over \mathbb{R} , we may assume that $|F(0)| = \|F\|_\infty$;
- by dilating F , we may assume that $\|F\|_1 = 1$;
- by multiplying F by a unimodular complex number, we may assume that $F(0) > 0$;
- by replacing $F(x)$ by $(F(x) + \overline{F(x)})/2$ we may assume that F is real-valued;
- by replacing $F(x)$ by $(F(x) + F(-x))/2$ we may assume that F is even.

From the definitions it is clear that $\mathcal{C}(A)$ and $\mathcal{C}^+(A)$ are non-increasing functions of A . The observations above show that in (1.1)–(1.2) we can restrict attention to even, real-valued functions, so that $\mathcal{C}(A) \leq \mathcal{C}^+(A)$. The Fejér kernel $F(x) = (\sin(\pi x)/(\pi x))^2$ reveals that $\mathcal{C}(\infty) \geq 1$. For every $F \in \mathcal{A}$ we have

$$|F(0)| - \int_{[-1,1]^c} |\hat{F}(t)| dt \leq \left| \int_{-1}^1 \hat{F}(t) dt \right| \leq 2\|F\|_1, \quad (2.1)$$

so that $\mathcal{C}(1) \leq 2$. A similar argument gives $\mathcal{C}^+(1) \leq 2$. Putting together all of these observations, for $1 \leq A \leq \infty$, we obtain the chain of inequalities

$$1 \leq \mathcal{C}(\infty) \leq \mathcal{C}(A) \leq \mathcal{C}^+(A) \leq \mathcal{C}^+(1) \leq 2.$$

2.1. Proof of Theorem 1.1 (a.1). This is the case $A = \infty$ and we are restricted to the class $\mathcal{E} \subset \mathcal{A}$ of continuous functions $F: \mathbb{R} \rightarrow \mathbb{C}$, with $F \in L^1(\mathbb{R})$ and $\text{supp}(\hat{F}) \subset [-1, 1]$. Let $\{F_n\}_{n \geq 1}$ be an extremizing sequence verifying the conditions above, i.e. a sequence $\{F_n\}_{n \geq 1} \subset \mathcal{E}$ of even and real-valued functions, with $\|F_n\|_1 = 1$, $\|F_n\|_\infty = F_n(0) > 0$, and

$$\lim_{n \rightarrow \infty} F_n(0) = \mathcal{C}(\infty).$$

Since $\mathcal{C}(\infty) \leq 2$, it follows that $\{F_n\}_{n \geq 1}$ is a bounded sequence in $L^2(\mathbb{R})$. Hence, there exists $G \in L^2(\mathbb{R})$ such that (after passing to a subsequence, if necessary) $F_n \rightharpoonup G$ weakly in $L^2(\mathbb{R})$. In this case, $\text{supp}(\hat{G}) \subset [-1, 1]$ and by Fourier inversion G can be taken continuous. For any $y \in \mathbb{R}$, we have

$$\begin{aligned} F_n(y) &= \int_{-1}^1 e^{2\pi i y t} \hat{F}_n(t) dt = \int_{-\infty}^{\infty} \frac{\sin 2\pi(x-y)}{\pi(x-y)} F_n(x) dx \\ &\rightarrow \int_{-\infty}^{\infty} \frac{\sin 2\pi(x-y)}{\pi(x-y)} G(x) dx = \int_{-1}^1 e^{2\pi i y t} \hat{G}(t) dt = G(y), \end{aligned}$$

as $n \rightarrow \infty$. It follows that G is even, real-valued and $G(0) = \mathcal{C}(\infty)$. Moreover, by Fatou's lemma, we have $\|G\|_1 \leq 1$. Hence $G \in \mathcal{E}$, and from the definition of $\mathcal{C}(\infty)$ we must have $\|G\|_1 = 1$ which makes G an extremizer. Multiplying this G by the constant factor $\mathcal{C}(\infty)^{-1}$ we arrive at the extremizer stated in the theorem (that assumes the value 1 at $x = 0$).

2.2. Proof of Theorem 1.1 (b). We already observed in (2.1) that $\mathcal{C}(1) \leq 2$. By taking $F_\varepsilon(x) = \frac{1}{\sqrt{\varepsilon}} e^{-\pi x^2/\varepsilon}$ with $\varepsilon \rightarrow 0^+$ we see that $\mathcal{C}(1) = 2$. In order to obtain equality in (2.1) we must have $\hat{F}(t) = c\|F\|_1$ for all $t \in [-1, 1]$, for some constant $c \in \mathbb{C}$ with $|c| = 1$. This is not possible, and hence there are no extremizers in this case.

2.3. Proof of Theorem 1.1 (c.1). Here $1 < A < \infty$. Suppose $F \in \mathcal{A}$ is non-identically zero, with

$$\frac{1}{2} \leq \frac{|F(0)| - A \int_{[-1,1]^c} |\hat{F}(t)| dt}{\|F\|_1}. \quad (2.2)$$

Since

$$\int_{[-1,1]^c} |\hat{F}(t)| dt \geq \left| F(0) - \int_{-1}^1 \hat{F}(t) dt \right| \geq |F(0)| - 2\|F\|_1,$$

we may use (2.2) to see that

$$|F(0)| \leq \frac{2A - \frac{1}{2}}{A - 1} \|F\|_1. \quad (2.3)$$

Inserting this estimate into (2.2), we also have

$$\int_{[-1,1]^c} |\hat{F}(t)| dt \leq \frac{3}{2(A-1)} \|F\|_1. \quad (2.4)$$

Let $\{F_n\}_{n \geq 1} \subset \mathcal{A}$ be an extremizing sequence of even and real-valued functions, with $\|F_n\|_1 = 1$, $\|F_n\|_\infty = F_n(0) > 0$, and $\hat{F}_n \in L^1(\mathbb{R})$. Thus

$$\lim_{n \rightarrow \infty} \left(F_n(0) - A \int_{[-1,1]^c} |\hat{F}_n(t)| dt \right) = \mathcal{C}(A).$$

Since $\mathcal{C}(A) \geq 1$, from our observation in (2.3) we see that $\{F_n(0)\}_{n \geq 1}$ is a bounded sequence, and from (2.4) that $\{\|\hat{F}_n\|_1\}_{n \geq 1}$ is also bounded.

2.3.1. Step 1. Since $\|F_n\|_\infty = F_n(0)$, the sequence $\{F_n\}_{n \geq 1}$ is bounded in $L^2(\mathbb{R})$. Passing to a subsequence, if necessary, we may assume that $F_n(0) \rightarrow c$, for some constant $c \geq \mathcal{C}(A)$, and that $F_n \rightharpoonup G$ weakly in $L^2(\mathbb{R})$ for some $G \in L^2(\mathbb{R})$. By Mazur's lemma [4, Corollary 3.8 and Exercise 3.4], there exists a sequence $H_k \rightarrow G$ strongly in $L^2(\mathbb{R})$, with $H_k \in \text{Conv}(\{F_n\}_{n \geq k})$ (i.e. each H_k is a *finite* linear convex combination of functions F_n with $n \geq k$). Note that H_k is even and real-valued, $\|H_k\|_\infty = H_k(0) \rightarrow c$, $\|H_k\|_1 \leq 1$, and $\{\|\hat{H}_k\|_1\}_{k \geq 1}$ remains bounded. By passing to a further subsequence, we may also assume that $H_k \rightarrow G$ and $\hat{H}_k \rightarrow \hat{G}$, pointwise almost everywhere. Hence G is also even and real-valued. Note that $\{H_k\}_{k \geq 1}$ is also an extremizing sequence.

2.3.2. Step 2. By Fatou's lemma $\|G\|_1 \leq \liminf_{k \rightarrow \infty} \|H_k\|_1 \leq 1$ and $\|\hat{G}\|_1 \leq \liminf_{k \rightarrow \infty} \|\hat{H}_k\|_1 < \infty$. By Fourier inversion, we may assume that G is continuous (after eventually modifying it on a set of measure zero), hence $G \in \mathcal{A}$. First we claim that G is nonzero. In fact, since $\{H_k\}_{k \geq 1}$ is an extremizing sequence and $H_k(0) \rightarrow c \geq \mathcal{C}(A)$, from (2.3) we find that $\liminf_{k \rightarrow \infty} \|H_k\|_1 \geq c_1 > 0$. From the L^2 -convergence (applied below just in the interval $[-1, 1]$) and Fatou's lemma, we have

$$\begin{aligned} G(0) - A \int_{[-1,1]^c} |\hat{G}(t)| dt &= \int_{-1}^1 \hat{G}(t) dt - \int_{[-1,1]^c} (|\hat{G}(t)| - \hat{G}(t)) dt - (A-1) \int_{[-1,1]^c} |\hat{G}(t)| dt \\ &\geq \limsup_{k \rightarrow \infty} \left(\int_{-1}^1 \hat{H}_k(t) dt - \int_{[-1,1]^c} (|\hat{H}_k(t)| - \hat{H}_k(t)) dt \right. \\ &\quad \left. - (A-1) \int_{[-1,1]^c} |\hat{H}_k(t)| dt \right) \end{aligned}$$

$$\begin{aligned}
&= \limsup_{k \rightarrow \infty} \left(H_k(0) - A \int_{[-1,1]^c} |\hat{H}_k(t)| dt \right) \\
&\geq c_1 \mathcal{C}(A).
\end{aligned}$$

This shows that G is nonzero. The same computation above (up to its third line) shows that G is indeed an extremizer, since $\|G\|_1 \leq \liminf_{k \rightarrow \infty} \|H_k\|_1$.

2.4. Proof of Theorem 1.2 (a.1), (b), and (c.1). The proof of part (b) follows along the same lines as the argument in §2.2 (with the same extremizing family). The proofs of parts (a.1) and (c.1) follow the outline of §2.3 and we simply indicate the minor modifications needed.

In seeking extremizers when $1 < A < \infty$, we may assume that $F(0) > 0$ and that $\hat{F} \in L^1(\mathbb{R})$ (recall that here we are already working within the class of even and real-valued functions). Suppose that $F \in \mathcal{A}^+$ is non-identically zero, with

$$\frac{1}{2} \leq \frac{F(0) - A \int_{[-1,1]^c} (\hat{F}(t))_+ dt}{\|F\|_1}. \quad (2.5)$$

Since

$$\int_{[-1,1]^c} (\hat{F}(t))_+ dt \geq F(0) - \int_{-1}^1 \hat{F}(t) dt \geq F(0) - 2\|F\|_1,$$

we may use (2.5) to see that

$$F(0) \leq \frac{2A - \frac{1}{2}}{A - 1} \|F\|_1. \quad (2.6)$$

As before, inserting this estimate into (2.5) we obtain

$$\int_{[-1,1]^c} (\hat{F}(t))_+ dt \leq \frac{3}{2(A-1)} \|F\|_1. \quad (2.7)$$

Let $\{F_n\}_{n \geq 1} \subset \mathcal{A}^+$ be an extremizing sequence with $\|F_n\|_1 = 1$, $F_n(0) > 0$, and $\hat{F}_n \in L^1(\mathbb{R})$. Note that, in principle, we do not necessarily have $\|F_n\|_\infty = F_n(0)$. Since $\mathcal{C}^+(A) \geq 1$, from (2.6) we see that $\{F_n(0)\}_{n \geq 1}$ is a bounded sequence, and from (2.7) we see that $\{\|\hat{F}_n\|_1\}_{n \geq 1}$ is also bounded.

The rest of the proof follows as in Steps 1 and 2 of §2.3. Note that the corresponding sequence $\{H_k\}_{k \geq 1}$ will be extremizing, due to the general property that $(f+g)_+ \leq f_+ + g_+$, and inequality (2.6) shows that $\liminf_{k \rightarrow \infty} \|H_k\|_1 \geq c_1 > 0$. For the final computation, one uses the identity

$$\begin{aligned}
G(0) - A \int_{[-1,1]^c} (\hat{G}(t))_+ dt &= \int_{-1}^1 \hat{G}(t) dt - \int_{[-1,1]^c} -(\hat{G}(t))_- dt \\
&\quad - (A-1) \int_{[-1,1]^c} (\hat{G}(t))_+ dt.
\end{aligned}$$

For the case $A = \infty$ (part (a.1)), the required modifications are similar and we omit the details.

3. Uniqueness of extremizers

In this section we continue the study of the extremal problem (1.2). We prove the uniqueness of a bandlimited extremizer (up to multiplication by a complex scalar) and provide its variational characterization as described in parts (a.2) and (a.3) of Theorem 1.1.

3.1. Proof of Theorem 1.1 (a.2). Let $G \in \mathcal{E} \subset \mathcal{A}$ be an even and real-valued extremizer of (1.2) with $G(0) = 1$. Let $G_1 \in \mathcal{E}$ be another extremizer of (1.2), with $G_1(0) = 1$. It suffices to show that $G_1 = G$.

Let $F = (G + G_1)/2$. Then, by the triangle inequality, we have

$$\int_{-\infty}^{\infty} |F(x)| dx \leq \frac{1}{2} \int_{-\infty}^{\infty} (|G(x)| + |G_1(x)|) dx = \frac{1}{\mathcal{C}(\infty)}, \quad (3.1)$$

and $F(0) = 1$. To avoid strict inequality in (3.1) we must have

$$|G(x) + G_1(x)| = |G(x)| + |G_1(x)|$$

for all $x \in \mathbb{R}$. In particular, this shows that $G_1: \mathbb{R} \rightarrow \mathbb{C}$ is real-valued and that

$$G(x) G_1(x) \geq 0$$

for all $x \in \mathbb{R}$. Let $R = G \cdot G_1$. Then R is a nonnegative and integrable function with $\text{supp}(\hat{R}) \subset [-2, 2]$. By a classical result of Krein [1, p. 154], we have $R(x) = |S(x)|^2$, for some $S \in L^2(\mathbb{R})$ with $\text{supp}(\hat{S}) \subset [-1, 1]$. Observe that $|S(0)| = 1$ and that

$$\begin{aligned} \int_{-\infty}^{\infty} |S(x)| dx &= \int_{-\infty}^{\infty} \sqrt{G(x) G_1(x)} dx \\ &\leq \frac{1}{2} \int_{-\infty}^{\infty} (|G(x)| + |G_1(x)|) dx = \frac{1}{\mathcal{C}(\infty)}. \end{aligned} \quad (3.2)$$

In particular $S \in L^1(\mathbb{R})$. To avoid strict inequality in (3.2) we must have $G(x) = G_1(x)$ for all $x \in \mathbb{R}$, completing the proof.

3.2. Proof of Theorem 1.1 (a.3). Let G be the unique extremal function of (1.2) with $G(0) = 1$. Let $F \in \mathcal{E}$ be a real-valued function with $F(0) = 0$ and define, for $\varepsilon \in \mathbb{R}$,

$$\Phi(\varepsilon) := \int_{-\infty}^{\infty} |G(x) + \varepsilon F(x)| dx = \int_{-\infty}^{\infty} ((G(x) + \varepsilon F(x))^2)^{1/2} dx.$$

This is a differentiable function of the variable ε and, since G is an extremizer, we must have

$$0 = \frac{\partial \Phi}{\partial \varepsilon}(0) = \int_{-\infty}^{\infty} \text{sgn}(G(x)) F(x) dx. \quad (3.3)$$

If $F_1 \in \mathcal{E}$ is a generic real-valued function (not necessarily with $F_1(0) = 0$), by (3.3) we obtain that

$$\begin{aligned} \mathcal{C}(\infty) \int_{-\infty}^{\infty} \operatorname{sgn}(G(x)) F_1(x) dx &= \mathcal{C}(\infty) \int_{-\infty}^{\infty} \operatorname{sgn}(G(x)) (F_1(x) - F_1(0)G(x)) dx \\ &\quad + \mathcal{C}(\infty) \int_{-\infty}^{\infty} \operatorname{sgn}(G(x)) F_1(0) G(x) dx \end{aligned} \quad (3.4)$$

$$= F_1(0). \quad (3.5)$$

Finally, if $F_2 \in \mathcal{E}$ is a generic complex-valued function, we may write $F_2(x) = A(x) - iB(x)$, where $A(x) = (F_2(x) + \overline{F_2(x)})/2$ and $B(x) = i(F_2(x) - \overline{F_2(x)})/2$ are real-valued functions in \mathcal{E} , and use (3.4) to arrive at

$$\mathcal{C}(\infty) \int_{-\infty}^{\infty} \operatorname{sgn}(G(x)) F_2(x) dx = F_2(0).$$

4. Upper and lower bounds

In this section we conclude the proofs of Theorems 1.1 and 1.2 by establishing the proposed upper and lower bounds for the sharp constants $\mathcal{C}(A)$ and $\mathcal{C}^+(A)$.

4.1. Approximations. For the purpose of finding the values of the sharp constants $\mathcal{C}(A)$ and $\mathcal{C}^+(A)$ in problems (1.1)–(1.4), without loss of generality we may work with smooth functions. For instance, let us show that we can simply consider $\hat{F} \in C_c^\infty(\mathbb{R})$. This observation is useful in some passages later in the paper.

Starting with $0 \neq F \in \mathcal{A}$ (or $0 \neq F \in \mathcal{A}^+$ in the case of (1.3)), we write

$$J(F) := \frac{|F(0)| - A \int_{[-1,1]^c} |\hat{F}(t)| dt}{\|F\|_1}$$

and
$$J^+(F) := \frac{F(0) - A \int_{[-1,1]^c} (\hat{F}(t))_+ dt}{\|F\|_1}.$$

In either situation we may also assume that $\hat{F} \in L^1(\mathbb{R})$ and that $J(F)$ and $J^+(F)$ are positive. Let $K(x) = (\sin(\pi x)/(\pi x))^2$ be the Fejér kernel and, for $\lambda > 0$, define $K_\lambda(x) = \lambda^{-1}K(x/\lambda)$. By Young's inequality we have $\|F * K_\lambda\|_1 \leq \|F\|_1$, and using dominated convergence it follows that $\limsup_{\lambda \rightarrow 0} J(F * K_\lambda) \geq J(F)$ and $\limsup_{\lambda \rightarrow 0} J^+(F * K_\lambda) \geq J^+(F)$. Hence we may assume that our test function F is bandlimited.

Let $\eta \in C_c^\infty(\mathbb{R})$ be an even, nonnegative, and radially non-increasing function such that $\eta(0) = 1$, $\text{supp}(\eta) \subset [-1, 1]$, and $\int_{-1}^1 \eta(x) dx = 1$. Again, let $\eta_\lambda(x) = \lambda^{-1} \eta(x/\lambda)$. If $\text{supp}(\hat{F}) \subset [-\Lambda, \Lambda]$, then $\hat{F} * \eta_\lambda \in C_c^\infty(\mathbb{R})$ and $\text{supp}(\hat{F} * \eta_\lambda) \subset [-\Lambda - \lambda, \Lambda + \lambda]$. By dominated convergence, we have $\lim_{\lambda \rightarrow 0} J(F \cdot \hat{\eta}_\lambda) = J(F)$ and $\lim_{\lambda \rightarrow 0} J^+(F \cdot \hat{\eta}_\lambda) = J^+(F)$. This verifies our claim in the cases $1 \leq A < \infty$. In the cases $A = \infty$ one has to slightly dilate F beforehand in order to apply the procedure above and arrive at a function in the class $\mathcal{E} \subset \mathcal{A}$ for (1.2) and $\mathcal{E}^+ \subset \mathcal{A}^+$ for (1.4).

4.2. Proof of Theorem 1.1 (a.4). The bounds

$$1.08185 \dots \leq \mathcal{C}(\infty) \leq 1.09769 \dots \quad (4.1)$$

were proved in the very interesting work of Gorbachev [25, Theorem 3], to which we refer the reader for details. These are slightly inferior to the ones presented in (1.5) but will be more convenient for our purposes, especially the upper bound, since the test functions used are described more explicitly. These bounds improved upon the work of Andreev, Konyagin, and Popov [2], who had previously obtained

$$c_0 = 1.07995 \dots \leq \mathcal{C}(\infty) \leq 1.17898. \quad (4.2)$$

As already pointed out in the introduction, the lower bound in (4.2) comes from the simple example

$$H(x) = \frac{\cos(2\pi x)}{1 - 16x^2}.$$

The Fourier transform of H is $\hat{H}(t) = \frac{\pi}{4} \cos(\pi t/2) \chi_{[-1,1]}(t)$, which may be verified by starting with our expression for $\hat{H}(t)$ and computing its Fourier transform to recover H . Thus H belongs to the class \mathcal{E} , and $H(0)$ is clearly 1. To compute the L^1 -norm of H we observe that $\text{sgn}(H(x)) = 2\chi_{[-\frac{1}{4}, \frac{1}{4}]}(x) - \text{sgn}(\cos 2\pi x)$, and use Plancherel's theorem and the fact that $\text{sgn}(\cos 2\pi x)$ has distributional Fourier transform supported outside $(-1, 1)$ to get¹

$$\begin{aligned} \|H\|_1 &= \int_{-\infty}^{\infty} |H(x)| dx = \int_{-\infty}^{\infty} (2\chi_{[-\frac{1}{4}, \frac{1}{4}]}(x) - \text{sgn}(\cos 2\pi x)) H(x) dx \\ &= \int_{-\infty}^{\infty} 2\chi_{[-\frac{1}{4}, \frac{1}{4}]}(x) H(x) dx = \int_{-1}^1 \left(\frac{2 \sin(\pi t/2)}{\pi t} \right) \left(\frac{\pi}{4} \cos(\pi t/2) \right) dt \\ &= \frac{\pi}{4} \int_{-1}^1 \frac{\sin \pi t}{\pi t} dt = 1/c_0. \end{aligned}$$

This example will be useful later on to generate lower bounds for $\mathcal{C}(A)$ in the general case $1 < A < \infty$. The lower bound of Gorbachev [25] in (4.1) comes from more complicated numerical examples.

¹The function $x \mapsto \text{sgn}(\cos 2\pi x)$ is an example of a *high pass function*, as studied in [34].

The upper bound in (4.1) comes from a dual formulation of the problem. Suppose that $\psi \in L^\infty(\mathbb{R})$ is such that its distributional Fourier transform is identically equal to 1 on the interval $(-1, 1)$. Let $\mathcal{S}(\mathbb{R})$ denote the Schwartz class. Then, for $F \in \mathcal{E} \cap \mathcal{S}(\mathbb{R})$ (as discussed in §4.1), we have

$$\|\psi\|_\infty \int_{-\infty}^{\infty} |F(x)| dx \geq \left| \int_{-\infty}^{\infty} F(x) \psi(x) dx \right| = \left| \int_{-1}^1 \hat{F}(t) \hat{\psi}(t) dt \right| = |F(0)|,$$

which implies that

$$\mathcal{C}(\infty) \leq \|\psi\|_\infty.$$

With this dual formulation, it suffices to exhibit a nice test function ψ .

We now briefly describe the construction of Gorbachev [25, Lemma 9]. To simplify the notation (and align with the terminology of [25] to facilitate the references) we let

$$j(x) = \frac{\sin(2\pi x)}{2\pi x}$$

in what follows. For $\tau = 29289/100000 = 0.29289$ we define a continuous and piecewise linear function $\alpha: [0, 1/2] \rightarrow \mathbb{R}$ by

$$\alpha(x) = \begin{cases} 2x - 1, & 0 \leq x \leq \tau; \\ 2\tau - 1 + 2(1 - \tau)(x - \tau)/\varepsilon, & \tau \leq x \leq \tau + \varepsilon; \\ 1, & \tau + \varepsilon \leq x \leq 1/2 - 2\varepsilon; \\ 1 - y(x - 1/2 + 2\varepsilon)/\varepsilon, & 1/2 - 2\varepsilon \leq x \leq 1/2 - \varepsilon; \\ 1 - y + y(x - 1/2 + \varepsilon)/\varepsilon, & 1/2 - \varepsilon \leq x \leq 1/2, \end{cases} \quad (4.3)$$

where

$$\varepsilon = \frac{\tau^2 - 2\tau + 1/2}{1 + y - 2\tau} > 0, \quad (4.4)$$

and, having defined (4.3) and (4.4), y is finally chosen so that

$$\int_0^{1/2} \frac{(1 - \alpha(x))}{j(x)} \cos(2\pi x) dx = 0.$$

One arrives at the values $y = 0.43056\dots$ and $\varepsilon = 0.0000053884\dots$. Let

$$d_0 = \left(\int_0^{1/2} \frac{1 - \alpha(x)}{j(x)} dx \right)^{-1} = 1.09769\dots \quad (4.5)$$

and define 1-periodic even functions $a(x)$ and $b(x)$ by

$$a(x) = d_0 \alpha(x) \quad \text{and} \quad b(x) = \frac{d_0 - a(x)}{2j(x)} - 1, \quad \text{for } x \in [0, 1/2].$$

As observed in [25], with this construction the functions a and b have Fourier series expansions

$$a(x) = \sum_{n=1}^{\infty} 2a_n \cos(2\pi nx), \quad b(x) = \sum_{n=2}^{\infty} 2b_n \cos(2\pi nx),$$

$$\sum_{n=1}^{\infty} |a_n| < \infty, \quad \text{and} \quad \sum_{n=2}^{\infty} |b_n| < \infty$$

(notice that the first Fourier coefficients verify $a_0 = b_0 = b_1 = 0$). A numerical evaluation leads to

$$a_1 = -0.5622\dots, \quad a_2 = 0.0684\dots, \quad a_3 = 0.1005\dots,$$

and since $\|a\|_{L^2[-\frac{1}{2}, \frac{1}{2}]}^2 = 0.7238\dots$ and $2a_1^2 = 0.6321\dots$, an application of Plancherel's theorem gives us that $|a_n| \leq |a_1|$ for all n . For the function b we adopt a slightly different approach to bounding the Fourier coefficients b_n (since $\|b\|_{L^2[-\frac{1}{2}, \frac{1}{2}]}$ is very large). A numerical integration yields

$$|b_n| \leq \int_{-1/2}^{1/2} |b(x)| dx = 0.8283\dots$$

for all $n \geq 2$.

Finally, let $\phi(x) = 2j(x)(1 + b(x))$, and define

$$\psi(x) = \phi(x) + a(x). \quad (4.6)$$

This is the test function constructed by Gorbachev [25], which verifies $\|\psi\|_{\infty} = d_0$ and has distributional Fourier transform identically equal to 1 on the interval $(-1, 1)$. In fact, we have

$$\begin{aligned} \hat{\psi}(t) &= \hat{\phi}(t) + \sum_{n=1}^{\infty} a_n (\delta(t-n) + \delta(t+n)) \\ &= \chi_{[-1,1]}(t) + \sum_{n=2}^{\infty} b_n (\chi_{[-1,1]}(t-n) + \chi_{[-1,1]}(t+n)) \\ &\quad + \sum_{n=1}^{\infty} a_n (\delta(t-n) + \delta(t+n)), \end{aligned} \quad (4.7)$$

where δ is the Dirac delta distribution. We shall use this construction to generate upper bounds for $\mathcal{C}(A)$ in the general case $1 < A < \infty$. The observation that $\|\hat{\phi}\|_{\infty} = 1$ will be relevant later on.

Remark. In an earlier version of this manuscript, without being aware of the references [2, 25], and [29], we had initially arrived at the test function

$$\begin{aligned}\tilde{\psi}(x) = 2\tilde{a}_0\chi_{[-\frac{1}{4}, \frac{1}{4}]}(x) + \sum_{n=1}^{\infty} 2\tilde{a}_n(\chi_{[-\frac{1}{4}, \frac{1}{4}]}(x - \frac{n}{2}) + \chi_{[-\frac{1}{4}, \frac{1}{4}]}(x + \frac{n}{2})) \\ - \tilde{a}_0 \operatorname{sgn}(\cos(2\pi x)),\end{aligned}$$

where $\tilde{a}_n = \frac{4}{\pi} \sum_{j=n}^{\infty} \frac{(-1)^j}{(2j+1)^2}$ are the Fourier coefficients in the expansion

$$\frac{(\pi t/2)}{\sin(\pi t/2)} = \tilde{a}_0 + 2 \sum_{n=1}^{\infty} \tilde{a}_n \cos(n\pi t)$$

for $-1 \leq t < 1$. This leads to the bound $\mathcal{C}(\infty) \leq \|\tilde{\psi}\|_{\infty} = \tilde{a}_0 = 1.16624\dots$, which is intermediate between (4.1) and (4.2).

4.3. Proof of Theorem 1.1 (c.2).

4.3.1. Lower bounds. As before, let $H(x) = (\cos 2\pi x)/(1 - 16x^2)$. Take $F(x) = H(x/\lambda)$ for a suitable parameter $\lambda \in (0, 1]$ to be optimized. Then $F(0) = 1$ and $\|F\|_1 = \lambda \|H\|_1 = \lambda/c_0$ with $c_0 = 1.079950\dots$. The ratio to be maximized is

$$\frac{c_0}{\lambda} \left(1 - A \lambda \frac{\pi}{4} \int_{1 \leq |t| \leq \frac{1}{\lambda}} \cos\left(\frac{\pi \lambda t}{2}\right) dt \right) = \frac{c_0}{\lambda} \left(1 - A \left(1 - \sin\left(\frac{\pi \lambda}{2}\right) \right) \right).$$

Calculus shows that this is maximized by choosing λ such that

$$1 - \frac{1}{A} = \sin\left(\frac{\pi \lambda}{2}\right) - \frac{\pi \lambda}{2} \cos\left(\frac{\pi \lambda}{2}\right). \quad (4.8)$$

For $\lambda = \lambda(A)$ verifying (4.8), this examples demonstrates that

$$\mathcal{C}(A) \geq \frac{\pi A c_0}{2} \cos\left(\frac{\pi \lambda(A)}{2}\right).$$

Note that as $A \rightarrow 1^+$, this lower bound goes to $\pi c_0/2$ and is not very effective. Alternatively, we can then use a dilation of the Fejér kernel $K(x) = (\sin(\pi x)/(\pi x))^2$ (note that $\hat{K}(t) = (1 - |t|)_+$). Again we consider $F(x) = K(x/\lambda)$ and optimize the dilation parameter $\lambda \in (0, 1]$. The ratio we seek to maximize is

$$\frac{1}{\lambda} \left(1 - A \lambda \int_{1 \leq |t| \leq \frac{1}{\lambda}} (1 - |\lambda t|)_+ dt \right) = \frac{1}{\lambda} - A \left(\frac{1}{\lambda} + \lambda - 2 \right).$$

The optimal choice is $\lambda = \sqrt{(A-1)/A}$, which leads to the bound

$$\mathcal{C}(A) \geq 2A - 2\sqrt{A(A-1)}.$$

4.3.2. Upper bounds. We already know that $\mathcal{C}(A) \leq \mathcal{C}(1) = 2$. The other upper bound comes from duality considerations. Suppose that $\varphi \in L^\infty(\mathbb{R})$ is such that its distributional Fourier transform is identically equal to 1 on the interval $(-1, 1)$ and $|\hat{\varphi}(t) - 1| \leq A$ for all $t \in \mathbb{R}$. Then, for $F \in \mathcal{A} \cap \mathcal{S}(\mathbb{R})$ (as discussed in §4.1), we have

$$\begin{aligned} \|\varphi\|_\infty \int_{-\infty}^{\infty} |F(x)| dx &\geq \left| \int_{-\infty}^{\infty} F(x) \varphi(x) dx \right| \\ &= \left| \int_{-\infty}^{\infty} \hat{F}(t) \hat{\varphi}(t) dt \right| \geq |F(0)| - A \int_{[-1,1]^c} |\hat{F}(t)| dt. \end{aligned}$$

This leads to $\mathcal{C}(A) \leq \|\varphi\|_\infty$.

Let ψ be defined by (4.6). The idea is to mollify this function (used in the case $A = \infty$) in order to “bring down the delta functions” in its Fourier transform into the acceptable range $|\hat{\varphi}(t) - 1| \leq A$ for all $t \in \mathbb{R}$. First we dilate $\hat{\psi}$ defined by (4.7) to push the delta functions away from the interval $[-1, 1]$, in other words, for $\gamma > 1$, we observe that

$$\hat{\psi}(t/\gamma) = \hat{\phi}(t/\gamma) + \sum_{n=1}^{\infty} \gamma a_n (\delta(t - \gamma n) + \delta(t + \gamma n)).$$

Let $R(t) = \chi_{[-1/2, 1/2]}(t)$. For $\lambda > 0$, we write $R_\lambda(t) = \lambda^{-1} R(t/\lambda)$ and define

$$\begin{aligned} \hat{\varphi}(t) &:= (\hat{\psi}(\cdot/\gamma) * R_\lambda)(t) \\ &= (\hat{\phi}(\cdot/\gamma) * R_\lambda)(t) + \sum_{n=1}^{\infty} \frac{\gamma a_n}{\lambda} (\chi_{[-\frac{\lambda}{2}, \frac{\lambda}{2}]}(t - \gamma n) + \chi_{[-\frac{\lambda}{2}, \frac{\lambda}{2}]}(t + \gamma n)). \end{aligned} \tag{4.9}$$

Recall that $|a_n| \leq |a_1| < 0.6$ for all $n \geq 1$. Let $c = 0.6$, so that all the delta functions in (4.7) have coefficients at most c . Let us assume that $A \geq 2 + c$ (so that our particular choices of λ and γ below verify $0 < \lambda \leq \gamma$). We choose $\gamma - 1 = \frac{\lambda}{2}$ (so that the support of the mollified delta functions in (4.9) stay away from the interval $(-1, 1)$) and $\frac{c\gamma}{\lambda} = A - 2$ (so that the height of the mollified delta functions in (4.9) is at most $A - 2$). This leads to the explicit forms

$$\lambda = \frac{2}{\frac{2}{c}(A-2) - 1} \quad \text{and} \quad \gamma = \frac{1}{1 - \frac{c}{2(A-2)}}.$$

From (4.9) we conclude that $\hat{\varphi}(t) = 1$ for $t \in (-1, 1)$ and, since $\|\hat{\phi}\|_\infty = 1$, we also have $|\hat{\varphi}(t)| \leq A - 1$ for all $t \in \mathbb{R}$, which in particular implies that $|\hat{\varphi}(t) - 1| \leq A$ for all $t \in \mathbb{R}$ (note that the mollified delta functions on the right-hand side of (4.9) have disjoint supports due to the fact that $\lambda \leq \gamma$). Since $\varphi(x) = \gamma \psi(\gamma x) \hat{R}(x/\lambda)$, our upper bound is then $\|\varphi\|_\infty = \gamma \|\psi\|_\infty = \gamma d_0$.

4.4. Proof of Theorem 1.2 (a.2). We proceed again via duality considerations. Suppose that $\Psi \in L^\infty(\mathbb{R})$ is a real-valued function such that its distributional Fourier transform is identically equal to 1 on the interval $(-1, 1)$ and $\hat{\Psi}(t) - 1 \leq 0$ for all $t \in \mathbb{R}$. Then, for $F \in \mathcal{E}^+ \cap \mathcal{S}(\mathbb{R})$ (as discussed in §4.1), we have

$$\|\Psi\|_\infty \int_{-\infty}^{\infty} |F(x)| dx \geq \int_{-\infty}^{\infty} F(x) \Psi(x) dx = \int_{-\infty}^{\infty} \hat{F}(t) \hat{\Psi}(t) dt \geq F(0),$$

which implies that

$$\mathcal{C}^+(\infty) \leq \|\Psi\|_\infty.$$

Experimentation gave the following numerical example. Let $a = 0.018$, $b = 0.027$, and $c = 0.002$, and consider

$$\begin{aligned} \Psi(x) = & \frac{\sin(2\pi x)}{\pi x} + \frac{2 \sin(a\pi x)}{\pi x} \cos(3\pi x) + \frac{2 \sin(b\pi x)}{\pi x} \cos(4\pi x) \\ & + \frac{2 \sin(c\pi x)}{\pi x} \cos(10\pi x) - 0.888 \cos(2\pi x) - 0.01 \cos(6\pi x), \end{aligned} \quad (4.10)$$

which has Fourier transform

$$\begin{aligned} \hat{\Psi}(t) = & \chi_{[-1,1]}(t) + \chi_{[-a/2,a/2]}(t - \tfrac{3}{2}) + \chi_{[-a/2,a/2]}(t + \tfrac{3}{2}) \\ & + \chi_{[-b/2,b/2]}(t - 2) + \chi_{[-b/2,b/2]}(t + 2) \\ & + \chi_{[-c/2,c/2]}(t - 5) + \chi_{[-c/2,c/2]}(t + 5) \\ & - 0.444(\delta(t + 1) + \delta(t - 1)) - 0.005(\delta(t + 3) + \delta(t - 3)). \end{aligned} \quad (4.11)$$

For this test function we have $\|\Psi\|_\infty < 1.2$.




Figure 1. Graph of the function Ψ in (4.10) in two different scales.

4.5. Proof of Theorem 1.2 (c.2). We have already seen that $\mathcal{C}^+(A) \leq \mathcal{C}^+(1) = 2$. The other upper bound comes from the following dual formulation. Suppose that $\Phi \in L^\infty(\mathbb{R})$ is a real-valued function such that its distributional Fourier transform is

identically equal to 1 on the interval $(-1, 1)$ and $-A \leq \hat{\Phi}(t) - 1 \leq 0$ for all $t \in \mathbb{R}$. Then, for $F \in \mathcal{A}^+ \cap \mathcal{S}(\mathbb{R})$ (as discussed in §4.1), we have

$$\begin{aligned} \|\Phi\|_\infty \int_{-\infty}^{\infty} |F(x)| \, dx &\geq \int_{-\infty}^{\infty} F(x) \Phi(x) \, dx \\ &= \int_{-\infty}^{\infty} \hat{F}(t) \hat{\Phi}(t) \, dt \geq F(0) - A \int_{[-1,1]^c} (\hat{F}(t))_+ \, dt, \end{aligned}$$

which leads to

$$\mathcal{C}^+(A) \leq \|\Phi\|_\infty.$$

The idea is to mollify the test function in (4.10) to bring down the delta functions to the required range, as done in §4.3.2. Let $c = 0.444$ be the largest coefficient of a delta function in (4.11) and assume a priori that $A > 1 + \frac{c}{2}$ (so that our choice of λ below is in fact positive). With the same notation as in (4.9) we choose $\gamma - 1 = \frac{\lambda}{2}$ and $\frac{c\gamma}{\lambda} = A - 1$. Note that the four delta functions in (4.11) have negative coefficients, while the rest of the Fourier transform lies between 0 and 1, so we may take $A - 1$ here instead of $A - 2$. Moreover, since these delta functions are supported in non-consecutive integers, the condition $\gamma - 1 = \frac{\lambda}{2}$ already guarantees that the mollified delta functions will not overlap (hence we do not need to assume here that $\lambda \leq \gamma$). This yields

$$\lambda = \frac{2}{\frac{2}{c}(A-1) - 1} \quad \text{and} \quad \gamma = \frac{1}{1 - \frac{c}{2(A-1)}}.$$

Since $\Phi(x) = \gamma \Psi(\gamma x) \hat{R}(x/\lambda)$, our upper bound is $\|\Phi\|_\infty \leq \gamma \|\Psi\|_\infty < \gamma \times 1.2$.

4.6. Proof of Theorem 1.2 (c.3). For the specific value of $A = \frac{36}{11}$, the lower bound described in (1.8) and (1.6) yields $\mathcal{C}^+(\frac{36}{11}) \geq 1.1569\dots$. We found a better example through experimentation. The function

$$\begin{aligned} F(x) = & -4.8 x^2 e^{-3.3x^2} + 1.5 x^2 e^{-7.4x^2} \\ & + 520 x^{24} e^{-9.7x^2} + 1.3 e^{-2.8x^2} + 0.18 e^{-2x^2} \end{aligned} \quad (4.12)$$

gives

$$\frac{F(0) - A \int_{[-1,1]^c} (\hat{F}(t))_+ \, dt}{\|F\|_1} = 1.1943\dots > \frac{25}{21}.$$

We have found more complicated examples that do slightly better.




Figure 2. Graph of the function F in (4.12) on the left, and graph of \hat{F} on the right.

5. Prime gaps: asymptotic version

In this section we prove Theorem 1.3. The proof uses two main tools: the explicit formula connecting the prime numbers and the zeros of the Riemann zeta-function, and the Brun–Titchmarsh inequality as expressed in (1.11) and (1.12).

Lemma 5.1 (Guinand–Weil explicit formula). *Let $h(s)$ be analytic in the strip $|\operatorname{Im} s| \leq \frac{1}{2} + \varepsilon$ for some $\varepsilon > 0$, and assume that $|h(s)| \ll (1 + |s|)^{-(1+\delta)}$ for some $\delta > 0$ when $|\operatorname{Re} s| \rightarrow \infty$. Then*

$$\sum_{\rho} h\left(\frac{\rho - \frac{1}{2}}{i}\right) = h\left(\frac{1}{2i}\right) + h\left(-\frac{1}{2i}\right) - \frac{1}{2\pi} \hat{h}(0) \log \pi \\ + \frac{1}{2\pi} \int_{-\infty}^{\infty} h(u) \operatorname{Re} \frac{\Gamma'}{\Gamma}\left(\frac{1}{4} + \frac{iu}{2}\right) du - \frac{1}{2\pi} \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \left(\hat{h}\left(\frac{\log n}{2\pi}\right) + \hat{h}\left(-\frac{\log n}{2\pi}\right) \right),$$

where $\rho = \beta + i\gamma$ are the non-trivial zeros of $\zeta(s)$, Γ'/Γ is the logarithmic derivative of the Gamma function, and $\Lambda(n)$ is the Von Mangoldt function defined to be $\log p$ if $n = p^m$ with p a prime number and $m \geq 1$ an integer, and zero otherwise.

Proof. The proof follows from [31, Theorem 5.12]. □

5.1. Set-up. Motivated by the discussion in §4.1, throughout this section we fix $F: \mathbb{R} \rightarrow \mathbb{R}$ to be an even and bandlimited Schwartz function, with $F(0) > 0$. Let us assume that $\operatorname{supp}(\hat{F}) \subset [-N, N]$ for some parameter $N \geq 1$. It then follows that F extends to an entire function (which we continue calling F) and the fact that $x^2 F(x) \in L^\infty(\mathbb{R})$ implies, via the Phragmén–Lindelöf principle, that $|F(s)| \ll (1 + |s|)^{-2}$ when $|\operatorname{Re} s| \rightarrow \infty$. We may therefore apply the explicit formula (Lemma 5.1).

Our idea to approach this problem can be summarized as follows. We use the explicit formula above to measure the size of an interval that does not contain too

many primes. Note that the information about the primes is on the right-hand side of the formula, while on the left-hand side we have information on the zeros of $\zeta(s)$. We modify our test function F in such a way that \hat{F} emphasizes the information on said interval, translating and rescaling \hat{F} to concentrate the mass of \hat{F} on this interval. We then try to understand the effect of this localization in all the terms of the formula through an asymptotic analysis. Since the function \hat{F} must be small near its endpoints, it is advantageous to use the Brun–Titchmarsh inequality to (over) estimate the contribution from the primes on the edges of the interval.

Let $0 < \Delta \leq 1$ and $1 < a$ be free parameters to be chosen later. We anticipate that we will be choosing $\Delta \rightarrow 0^+$ and $a \rightarrow \infty$, so it is not harmful to further assume that

$$2\pi\Delta N \leq \log a. \quad (5.1)$$

Define $f(z) := \Delta F(\Delta z)$ and note that $\text{supp}(\hat{f}) \subset [-\Delta N, \Delta N]$. Assuming RH, an application of the explicit formula (Lemma 5.1) to the entire function $h(z) = f(z)a^{iz}$ yields the following inequality:

$$\begin{aligned} f\left(\frac{1}{2i}\right)a^{1/2} + f\left(-\frac{1}{2i}\right)a^{-1/2} &\leq \sum_{\gamma} |f(\gamma)| + \frac{1}{2\pi} \hat{f}\left(-\frac{\log a}{2\pi}\right) \log \pi \\ &\quad + \left| \frac{1}{2\pi} \int_{-\infty}^{\infty} f(u) a^{iu} \operatorname{Re} \frac{\Gamma'}{\Gamma}\left(\frac{1}{4} + \frac{iu}{2}\right) du \right| \\ &\quad + \frac{1}{2\pi} \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \left(\hat{f}\left(\frac{\log(n/a)}{2\pi}\right)_+ + \hat{f}\left(-\frac{\log na}{2\pi}\right)_+ \right). \end{aligned} \quad (5.2)$$

5.2. Proof of Theorem 1.3. The idea is to proceed with an asymptotic evaluation of both sides of (5.2). We start with its left-hand side. Note that

$$\begin{aligned} f\left(\frac{1}{2i}\right) &= \Delta F\left(\frac{\Delta}{2i}\right) = \Delta \int_{-N}^N e^{\pi t \Delta} \hat{F}(t) dt \\ &= \Delta \int_{-N}^N \hat{F}(t) dt + \Delta \int_{-N}^N (e^{\pi t \Delta} - 1) \hat{F}(t) dt \\ &= \Delta F(0) + O(\Delta^2). \end{aligned}$$

We may similarly estimate $f(-\frac{1}{2i})$. Therefore, the left-hand side of (5.2) equals

$$f\left(\frac{1}{2i}\right)a^{1/2} + f\left(-\frac{1}{2i}\right)a^{-1/2} = \Delta F(0)(a^{1/2} + a^{-1/2}) + O(\Delta^2 a^{1/2}).$$

For the right-hand side of (5.2), we first consider the error terms. From (5.1) we have

$$\frac{1}{2\pi} \hat{f}\left(-\frac{\log a}{2\pi}\right) \log \pi = \frac{1}{2\pi} \hat{F}\left(-\frac{\log a}{2\pi\Delta}\right) \log \pi = 0.$$

Also, using Stirling's formula $\frac{\Gamma'}{\Gamma}(s) = \log s + O(|s|^{-1})$ and (5.1), we get

$$\begin{aligned} \int_{-\infty}^{\infty} f(u) a^{iu} \operatorname{Re} \frac{\Gamma'}{\Gamma}\left(\frac{1}{4} + \frac{iu}{2}\right) du \\ &= \int_{-\infty}^{\infty} F(y) e^{2\pi i y (\frac{\log a}{2\pi\Delta})} \log \left| \frac{1}{4} + \frac{iy}{2\Delta} \right| dy + O(1) \\ &= \int_{-\infty}^{\infty} F(y) e^{2\pi i y (\frac{\log a}{2\pi\Delta})} \left(\frac{1}{2} \log(\Delta^2 + 4y^2) + \log\left(\frac{1}{4\Delta}\right) \right) dy + O(1) \\ &= \log\left(\frac{1}{4\Delta}\right) \hat{F}\left(-\frac{\log a}{2\pi\Delta}\right) + O(1) = O(1). \end{aligned}$$

Thus, we have deduced that

$$\begin{aligned} \Delta F(0)(a^{1/2} + a^{-1/2}) &\leq \sum_{\gamma} |f(\gamma)| \\ &+ \frac{1}{2\pi} \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \left(\hat{f}\left(\frac{\log(n/a)}{2\pi}\right)_+ + \hat{f}\left(-\frac{\log na}{2\pi}\right)_+ \right) + O(\Delta^2 a^{1/2}) + O(1). \end{aligned} \quad (5.3)$$

It remains to estimate the two remaining sums on right-hand side of this inequality.

5.2.1. The sum over zeros. Let $N(x)$ denote the number of zeros $\rho = \beta + i\gamma$ of $\zeta(s)$ with $0 < \gamma \leq x$. Using the fact that $N(x) = \frac{x}{2\pi} \log \frac{x}{2\pi} - \frac{x}{2\pi} + O(\log x)$, we evaluate the sum $\sum_{\gamma} |f(\gamma)|$ using summation by parts to get

$$\sum_{\gamma} |f(\gamma)| = \frac{1}{2\pi} \int_{-\infty}^{\infty} |f(x)| \log^+ \frac{|x|}{2\pi} dx + O(\|f\|_{\infty} + \|f'(x) \log^+ |x|\|_1),$$

where $\log^+ x = \max\{\log x, 0\}$ for $x > 0$. Recalling that $f(x) = \Delta F(\Delta x)$, this yields

$$\begin{aligned} \sum_{\gamma} |f(\gamma)| &= \frac{1}{2\pi} \int_{-\infty}^{\infty} |F(y)| \log^+ |y/2\pi\Delta| dy + O(1) \\ &= \frac{\log(1/2\pi\Delta)}{2\pi} \|F\|_1 + O(1). \end{aligned} \quad (5.4)$$

5.2.2. The sum over primes and the choice of parameters. Fix $\alpha \geq 0$ and assume that c is a fixed positive constant such that

$$\liminf_{x \rightarrow \infty} \frac{\pi(x + c\sqrt{x} \log x) - \pi(x)}{\sqrt{x}} \leq \alpha.$$

Then, given $\varepsilon > 0$, there exists a sequence of $x \rightarrow \infty$ such that

$$\frac{\pi(x + c\sqrt{x} \log x) - \pi(x)}{\sqrt{x}} \leq \alpha + \varepsilon \quad (5.5)$$

along this sequence. For each such x , we choose a and Δ such that

$$[x, x + c\sqrt{x} \log x] = [ae^{-2\pi\Delta}, ae^{2\pi\Delta}]. \quad (5.6)$$

Then (allowing the implicit constants in the big- O notation here to depend on c) we have

$$4\pi\Delta = \log\left(1 + c\frac{\log x}{\sqrt{x}}\right) = c\frac{\log x}{\sqrt{x}} + O\left(\frac{\log^2 x}{x}\right) \quad (5.7)$$

and

$$a = x\left(1 + c\frac{\log x}{\sqrt{x}}\right)^{1/2} = x + O(\sqrt{x} \log x). \quad (5.8)$$

By (5.1), the sum we want to evaluate is

$$\sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \left(\hat{f}\left(\frac{\log(n/a)}{2\pi}\right)_+ + \hat{f}\left(-\frac{\log na}{2\pi}\right)_+ \right) = \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \hat{F}\left(\frac{\log(n/a)}{2\pi\Delta}\right)_+. \quad (5.9)$$

Note that the last sum is supported on n with $ae^{-2\pi\Delta N} \leq n \leq ae^{2\pi\Delta N}$. The contribution of the (at most) $(\alpha + \varepsilon)\sqrt{x}$ primes in the interval $(x, x + c\sqrt{x} \log x] = (ae^{-2\pi\Delta}, ae^{2\pi\Delta}]$ to the sum (5.9) is bounded above by (using the trivial bound $(\hat{F}(t))_+ \leq \|F\|_1$)

$$\leq \|F\|_1 \sum_{p \in (ae^{-2\pi\Delta}, ae^{2\pi\Delta}]} \frac{\log p}{\sqrt{p}} \leq \|F\|_1 (\alpha + \varepsilon)\sqrt{x} \frac{\log x}{\sqrt{x}} = \|F\|_1 (\alpha + \varepsilon) \log x.$$

It is not hard to show that the contribution of the prime powers $n = p^k$ with $k \geq 2$ in the full interval $[ae^{-2\pi\Delta N}, ae^{2\pi\Delta N}]$ to the sum (5.9) is $O(1)$. It remains to estimate the contribution of the primes in the intervals $[ae^{-2\pi\Delta N}, ae^{-2\pi\Delta}]$ and $[ae^{2\pi\Delta}, ae^{2\pi\Delta N}]$, and for this we use the Brun–Titchmarsh inequality. Let \mathbf{B} be defined by (1.11) and let $\mathbf{B}' > \mathbf{B}$. For x sufficiently large we have

$$\begin{aligned} \sum_{1 \leq \left| \frac{\log(p/a)}{2\pi\Delta} \right| \leq N} \frac{\log p}{\sqrt{p}} \hat{F}\left(\frac{\log(p/a)}{2\pi\Delta}\right)_+ &\leq \mathbf{B}' \int_{1 \leq \left| \frac{\log(t/a)}{2\pi\Delta} \right| \leq N} \hat{F}\left(\frac{\log(t/a)}{2\pi\Delta}\right)_+ \frac{dt}{\sqrt{t}} + O(1) \\ &= \mathbf{B}' \sqrt{a} (2\pi\Delta) \int_{[-1,1]^c} (\hat{F}(t))_+ dt + O(1). \end{aligned}$$

The inequality above can be seen by covering the intervals $[ae^{-2\pi\Delta N}, ae^{-2\pi\Delta}]$ and $[ae^{2\pi\Delta}, ae^{2\pi\Delta N}]$ by subintervals of size \sqrt{a} , and applying the Brun–Titchmarsh

inequality in each summand of the corresponding Riemann–Stieltjes sum associated to this partition (the details of this argument are carried out in §6.2 for a specific function and can be modified to handle the general case). Combining estimates, we see that

$$\begin{aligned} \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \left(\widehat{f} \left(\frac{\log(n/a)}{2\pi} \right)_+ + \widehat{f} \left(-\frac{\log na}{2\pi} \right)_+ \right) \\ \leq \|F\|_1 (\alpha + \varepsilon) \log x + \mathbf{B}' \sqrt{a} (2\pi \Delta) \int_{[-1,1]^c} (\widehat{F}(t))_+ dt + O(1). \end{aligned} \quad (5.10)$$

5.2.3. Conclusion. Inserting the estimates in (5.4) and (5.10) into (5.3) and then rearranging terms, it follows that

$$\begin{aligned} \Delta \sqrt{a} \left(F(0) - \mathbf{B}' \int_{[-1,1]^c} (\widehat{F}(t))_+ dt \right) &\leq \frac{\log(1/2\pi \Delta)}{2\pi} \|F\|_1 \\ &\quad + \frac{1}{2\pi} \|F\|_1 (\alpha + \varepsilon) \log x + O(1), \end{aligned}$$

where we have used (5.7) and (5.8) to combine the error terms. Sending $x \rightarrow \infty$ along the sequence (5.5), we conclude that

$$c \leq (1 + 2\alpha + 2\varepsilon) \frac{\|F\|_1}{(F(0) - \mathbf{B}' \int_{[-1,1]^c} (\widehat{F}(t))_+ dt)},$$

where we naturally assume that the denominator above is positive. Since this holds for all $\varepsilon > 0$ and $\mathbf{B}' > \mathbf{B}$ we finally arrive at

$$c \leq (1 + 2\alpha) \frac{\|F\|_1}{(F(0) - \mathbf{B} \int_{[-1,1]^c} (\widehat{F}(t))_+ dt)}. \quad (5.11)$$

This is the connection to our extremal problem (1.3) and the discussion in §4.1 leads to the desired conclusion, since we may now optimize (5.11) over such bandlimited F .

6. Prime gaps: explicit version

We now move on to the proof of Theorem 1.5. Instead of initially following the proof outlined in Section 5 with a particular choice of test function F in the Guinand–Weil explicit formula (and carefully estimating the error terms), we start off slightly differently using a Mellin transform approach to the problem. For our fixed choice of test function, this approach simplifies some of our calculations. Moreover, it may be the case that the kernel we are using will be helpful in other applications. For a generic choice of test function, however, the Fourier transform approach to the problem used in the previous section is perhaps more illuminating.

Lemma 6.1. *Let ϑ and δ be positive numbers satisfying $\vartheta\delta = \pi/2$. Then, for $a > e^\delta$ and ϑ not an ordinate of a zero of $\zeta(s)$, we have*

$$\sum_{ae^{-\delta} \leq n \leq ae^\delta} \frac{\Lambda(n)}{\sqrt{n}} \cos\left(\vartheta \log \frac{a}{n}\right) = \frac{\vartheta \sqrt{a}}{\frac{1}{4} + \vartheta^2} (e^{\delta/2} + e^{-\delta/2}) - 2\vartheta \sum_{\gamma} \frac{a^{i\gamma} \cos(\delta\gamma)}{\vartheta^2 - \gamma^2} - \sum_{n=1}^{\infty} \frac{\vartheta a^{-2n-1/2}}{(2n+\frac{1}{2})^2 + \vartheta^2} (e^{(2n+1/2)\delta} + e^{-(2n+1/2)\delta}). \quad (6.1)$$

Here the first sum on the right-hand side runs over the nontrivial zeros $\rho = 1/2 + i\gamma$ of $\zeta(s)$ where $\gamma \in \mathbb{C}$ with $|\operatorname{Re}(\gamma)| < 1/2$.

Proof. For any $c > 0$, $\delta > 0$ and $\xi > 0$ we have

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \xi^s \left(\frac{e^{\delta s} - e^{-\delta s}}{s} \right) ds = \begin{cases} 1, & \text{if } e^{-\delta} < \xi < e^\delta, \\ 1/2, & \text{if } \xi = e^{\pm\delta}, \\ 0, & \text{otherwise.} \end{cases}$$

It then follows, for any $c > 1/2$, $a > 0$, $\delta > 0$ (assuming $ae^{\pm\delta} \notin \mathbb{N}$), and any real number ϑ , that

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} -\frac{\zeta'}{\zeta} \left(s + \frac{1}{2} + i\vartheta\right) a^{s+i\vartheta} \left(\frac{e^{\delta s} - e^{-\delta s}}{s} \right) ds = \sum_{ae^{-\delta} \leq n \leq ae^\delta} \frac{\Lambda(n)}{\sqrt{n}} \left(\frac{a}{n}\right)^{i\vartheta}.$$

For details on this calculation we refer to [16, Chapter 17]. Applying this formula at ϑ and $-\vartheta$ and then adding, we deduce that

$$\begin{aligned} & 2 \sum_{ae^{-\delta} \leq n \leq ae^\delta} \frac{\Lambda(n)}{\sqrt{n}} \cos\left(\vartheta \log \frac{a}{n}\right) \\ &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} -\frac{\zeta'}{\zeta} \left(w + \frac{1}{2}\right) a^w \left(\frac{e^{\delta(w-i\vartheta)} - e^{-\delta(w-i\vartheta)}}{w-i\vartheta} + \frac{e^{\delta(w+i\vartheta)} - e^{-\delta(w+i\vartheta)}}{w+i\vartheta} \right) dw. \end{aligned} \quad (6.2)$$

In the case $\vartheta\delta = \pi/2$, after dividing by 2, this formula simplifies to

$$\begin{aligned} & \sum_{ae^{-\delta} \leq n \leq ae^\delta} \frac{\Lambda(n)}{\sqrt{n}} \cos\left(\vartheta \log \frac{a}{n}\right) \\ &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} -\frac{\zeta'}{\zeta} \left(w + \frac{1}{2}\right) a^w \frac{\vartheta}{w^2 + \vartheta^2} (e^{\delta w} + e^{-\delta w}) dw; \end{aligned} \quad (6.3)$$

note the removable singularities of the integrand at $w = \pm i\vartheta$ and that the formula now holds when $ae^{\pm\delta} \in \mathbb{N}$, as well. Since $a > e^\delta$, we can shift the line of integration

left from $\operatorname{Re}(w) = c$ to $\operatorname{Re}(w) = -\infty$ and, using the calculus of residues, the integral in (6.3) equals

$$\begin{aligned} & \frac{\vartheta \sqrt{a}}{\frac{1}{4} + \vartheta^2} (e^{\delta/2} + e^{-\delta/2}) - 2\vartheta \sum_{\gamma} \frac{a^{i\gamma} \cos(\delta\gamma)}{\vartheta^2 - \gamma^2} \\ & - \sum_{n=1}^{\infty} \frac{\vartheta a^{-2n-1/2}}{(2n + \frac{1}{2})^2 + \vartheta^2} (e^{(2n+1/2)\delta} + e^{-(2n+1/2)\delta}). \end{aligned}$$

Combining estimates, the lemma follows. \square

Remark 6.2. Slightly more generally, if $\vartheta\delta \equiv \frac{\pi}{2} \pmod{\pi}$, then we can also evaluate the integral in (6.2) in terms of an absolutely convergent sum over the nontrivial zeros of $\zeta(s)$ (but not otherwise).

Since $\vartheta\delta = \pi/2$, the first term on the right-hand side of (6.1) is

$$\frac{\vartheta \sqrt{a}}{\frac{1}{4} + \vartheta^2} (e^{\delta/2} + e^{-\delta/2}) = 2\pi \sqrt{a} \left(\frac{\delta (e^{\delta/2} + e^{-\delta/2})}{\pi^2 + \delta^2} \right) \geq 2\pi \sqrt{a} \left(\frac{2\delta}{\pi^2} \right) = \frac{4\delta \sqrt{a}}{\pi}.$$

Our assumptions below imply that $e^{\delta}/a \leq 1/\sqrt{3}$, so the third term on the right-hand side of (6.1) is bounded in absolute value by

$$\frac{2}{\vartheta} \left(\frac{e^{\delta}}{a} \right)^{5/2} \sum_{n=0}^{\infty} \left(\frac{e^{\delta}}{a} \right)^{2n} \leq \frac{2}{\vartheta} \left(\frac{e^{\delta}}{a} \right)^{5/2} \sum_{n=0}^{\infty} \left(\frac{1}{3} \right)^n = \frac{3}{\vartheta} \left(\frac{e^{\delta}}{a} \right)^{5/2}.$$

Hence, taking absolute values in (6.1) and using the previous two estimates, it follows that

$$\frac{4\delta \sqrt{a}}{\pi} \leq \sum_{ae^{-\delta} \leq n \leq ae^{\delta}} \frac{\Lambda(n)}{\sqrt{n}} \cos\left(\vartheta \log \frac{a}{n}\right) + 2\vartheta \sum_{\gamma} \left| \frac{\cos(\delta\gamma)}{\vartheta^2 - \gamma^2} \right| + \frac{3}{\vartheta} \left(\frac{e^{\delta}}{a} \right)^{5/2}. \quad (6.4)$$

At this point, it is convenient to make a change of variables so that we can retrace our steps from the proof of Theorem 1.3 in Section 5 using a dilation of the Fourier transform pair

$$H(x) = \frac{\cos(2\pi x)}{1 - 16x^2} \quad \text{and} \quad \widehat{H}(t) = \frac{\pi}{4} \cos\left(\frac{\pi t}{2}\right) \chi_{[-1,1]}(t).$$

We set $f(x) = \Delta F(\Delta x)$ where $F(x) = H(x/\lambda)$ so that $\widehat{F}(t) = \lambda \widehat{H}(\lambda t)$. Then, letting

$$\delta = \frac{2\pi\Delta}{\lambda} \quad \text{and} \quad \vartheta = \frac{\lambda}{4\Delta}$$

in (6.4) (note that $\vartheta\delta = \pi/2$), after a little rearranging it follows that

$$\Delta\sqrt{a} \leq \sum_{\gamma} |f(\gamma)| + \frac{1}{2\pi} \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \widehat{F}\left(\frac{\log(n/a)}{2\pi\Delta}\right) + \frac{3}{2}\Delta\left(\frac{e^{2\pi\Delta/\lambda}}{a}\right)^{5/2}. \quad (6.5)$$

Note that the sum over n is supported on the interval $(ae^{-2\pi\Delta/\lambda}, ae^{2\pi\Delta/\lambda})$.

We assume that there are no primes in the interval $[x, x + c\sqrt{x}\log x]$ for $\frac{1}{2} \leq c \leq 1$, and we choose a and Δ to satisfy (5.6). In particular, the equalities in (5.7) and (5.8) still hold. As mentioned at the end of the introduction, we may assume that $x \geq 4 \cdot 10^{18}$. Using the fact that $\log(1+y) \leq y$ for $y \geq 0$ in (5.7), we note that $\Delta \leq \frac{1}{4\pi} \left(\frac{\log x}{\sqrt{x}}\right) < 10^{-8}$.

6.1. Sum over zeros. We now explicitly estimate the sum over the zeros of the zeta function on the right-hand side of (6.5).

Lemma 6.3. *Let $N(x)$ denote the number of zeros $\rho = \beta + i\gamma$ of $\zeta(s)$ with $0 < \gamma \leq x$. Then*

$$\left| N(x) - \frac{x}{2\pi} \log \frac{x}{2\pi e} - \frac{7}{8} \right| \leq 0.15 \log x + 3$$

for $x \geq e$.

Proof. The result holds for $e \leq x \leq 10$, since $N(10) = 0$. From [41, Corollary 1] we have

$$\left| N(x) - \frac{x}{2\pi} \log \frac{x}{2\pi e} - \frac{7}{8} \right| \leq 0.112 \log x + 0.278 \log \log x + 2.51 + \frac{0.2}{x}, \quad (6.6)$$

which holds for all $x \geq e$. The estimate

$$0.278 \log \log x \leq 0.038 \log x + 0.28 \quad (6.7)$$

holds for all $x \geq e$, while

$$\frac{0.2}{x} \leq 0.02 \quad (6.8)$$

holds for $x \geq 10$. Combining (6.6), (6.7), and (6.8), we arrive at our desired bound for $x \geq 10$. \square

Write

$$N(x) = \frac{x}{2\pi} \log \frac{x}{2\pi e} + \frac{7}{8} + R(x)$$

and let $x_0 = 9.676 \dots$ be such that

$$\frac{x_0}{2\pi} \log \frac{x_0}{2\pi e} + \frac{7}{8} = 0.$$

Then, assuming the Riemann hypothesis and using summation by parts and Lemma 6.3, we have

$$\begin{aligned} \sum_{\gamma > 0} |f(\gamma)| &= \int_{x_0}^{\infty} \left(\frac{1}{2\pi} \log \frac{x}{2\pi} \right) |f(x)| dx - \int_{x_0}^{\infty} R(x) |f'(x)| dx \\ &\leq \int_{x_0}^{\infty} \left(\frac{1}{2\pi} \log \frac{x}{2\pi} \right) |f(x)| dx + \int_{x_0}^{\infty} (0.15 \log x + 3) |f'(x)| dx \\ &= \int_{\Delta x_0}^{\infty} \left(\frac{1}{2\pi} \log \frac{y}{2\pi \Delta} \right) |F(y)| dy + \Delta \int_{\Delta x_0}^{\infty} (0.15 \log \frac{y}{\Delta} + 3) |F'(y)| dy. \end{aligned}$$

Therefore

$$\begin{aligned} \sum_{\gamma > 0} |f(\gamma)| &\leq \frac{1}{2\pi} \int_0^{\infty} \log^+ y |F(y)| dy + \frac{1}{2\pi} \log(1/2\pi \Delta) \int_0^{\infty} |F(y)| dy \\ &\quad + (0.15)\Delta \int_0^{\infty} \log^+ y |F'(y)| dy + (0.15)\Delta \log(1/\Delta) \int_0^{\infty} |F'(y)| dy \\ &\quad + 3\Delta \int_0^{\infty} |F'(y)| dy. \end{aligned}$$

The same bound holds for the zeros with $\gamma < 0$. Since $\Delta < 10^{-8}$ implies that $\Delta \log(1/\Delta) \leq 2 \times 10^{-7}$, we conclude that

$$\begin{aligned} \sum_{\gamma} |f(\gamma)| &\leq \frac{\log(1/2\pi \Delta)}{2\pi} \|F\|_1 + \frac{1}{2\pi} \|\log^+ |y| F(y)\|_1 \\ &\quad + (0.15) \times 10^{-8} \times \|\log^+ |y| F'(y)\|_1 \\ &\quad + (3 \times 10^{-8} + (0.15) \times 2 \times 10^{-7}) \|F'\|_1 \end{aligned} \quad (6.9)$$

$$< \frac{\log(1/2\pi \Delta)}{2\pi} \|F\|_1 + \frac{0.070}{2\pi}. \quad (6.10)$$

6.2. Sum over prime powers. We use a version of the Brun–Titchmarsh inequality due to Montgomery and Vaughan [35, Theorem 2] which states that

$$\pi(x+y) - \pi(x) < \frac{2y}{\log y}, \quad (6.11)$$

for all $x, y > 1$. For us, the relevant range is $y \geq \sqrt{x}$, so that (6.11) corresponds to an application of the Brun–Titchmarsh inequality with the bound $\mathbf{B} \leq 4$. This is slightly worse than Iwaniec’s bound (1.12) but is completely explicit. With $A = 4$, the lower bound in (1.6) was established in §4.3.1 with a dilation of the function $H(x)$ with dilation parameter $\lambda = \lambda(4) = 0.892422\dots$, leading to the

bound $\mathfrak{C}(4) \geq 1.141186\dots = (0.8762\dots)^{-1}$. For the sake of simplicity, we work instead with the dilation parameter $\lambda = 0.9$ and note that for $F(x) = H(x/\lambda)$ we have

$$J(F) = \frac{F(0) - 4 \int_{[-1,1]^c} |\hat{F}(t)| dt}{\|F\|_1} = 1.1405\dots > \frac{25}{22}. \quad (6.12)$$

With a and Δ chosen as in (5.6), we need to estimate the contribution of the primes p such that $1 < |\frac{\log p/a}{2\pi\Delta}| \leq \lambda^{-1}$ to the sum over n in (6.5). We cover the interval $(a e^{2\pi\Delta}, a e^{2\pi\Delta\lambda^{-1}}] \subset \cup_{j=0}^{J-1} (x_j, x_{j+1}]$, with $x_0 = a e^{2\pi\Delta}$ and $x_{j+1} = x_j + \sqrt{x_j}$. Using (6.11) in each subinterval $(x_j, x_{j+1}]$ and the fact that \hat{F} is decreasing on $[0, \lambda^{-1}]$ we obtain

$$\begin{aligned} \sum_{1 < \frac{\log p/a}{2\pi\Delta} \leq \lambda^{-1}} \frac{\log p}{\sqrt{p}} \hat{F}\left(\frac{\log(p/a)}{2\pi\Delta}\right) &\leq \sum_{j=0}^{J-1} \left(\frac{\log x_j}{\sqrt{x_j}} \hat{F}\left(\frac{\log(x_j/a)}{2\pi\Delta}\right)\right) \frac{4\sqrt{x_j}}{\log x_j} \\ &\leq 4\hat{F}(1) + \frac{4}{\sqrt{a}} \sum_{j=1}^{J-1} \hat{F}\left(\frac{\log(x_j/a)}{2\pi\Delta}\right) \sqrt{x_{j-1}} \\ &\leq 4\hat{F}(1) + \frac{4}{\sqrt{a}} \int_{x_0}^{x_J} \hat{F}\left(\frac{\log(t/a)}{2\pi\Delta}\right) dt \\ &= 4\hat{F}(1) + 4\sqrt{a} (2\pi\Delta) \int_1^{\lambda^{-1}} \hat{F}(y) e^{2\pi\Delta y} dy \\ &= 4\hat{F}(1) + 4\sqrt{a} (2\pi\Delta) \int_1^{\lambda^{-1}} \hat{F}(y) dy + 4\sqrt{a} (2\pi\Delta) \int_1^{\lambda^{-1}} \hat{F}(y) (e^{2\pi\Delta y} - 1) dy \\ &\leq 4\hat{F}(1) + 4\sqrt{a} (2\pi\Delta) \left(\int_1^{\lambda^{-1}} \hat{F}(y) dy\right) + 4\sqrt{a} (4\pi\Delta)^2 \hat{F}(1) (\lambda^{-1} - 1), \end{aligned} \quad (6.13)$$

where we have used the basic estimate $e^x - 1 \leq 2x$, for $x \leq 1$, in the last passage. We treat the other interval in a similar way, covering $[a e^{-2\pi\Delta\lambda^{-1}}, a e^{-2\pi\Delta}) \subset \cup_{j=0}^{L-1} [x_{j+1}, x_j)$, with $x_0 = a e^{-2\pi\Delta}$ and $x_j = x_{j+1} + \sqrt{x_{j+1}}$. Using (6.11) in each subinterval $[x_{j+1}, x_j)$ and the fact that \hat{F} is increasing on $[-\lambda^{-1}, 0]$ we obtain

$$\begin{aligned} \sum_{-\lambda^{-1} < \frac{\log p/a}{2\pi\Delta} \leq -1} \frac{\log p}{\sqrt{p}} \hat{F}\left(\frac{\log(p/a)}{2\pi\Delta}\right) &\leq \sum_{j=0}^{L-1} \left(\frac{\log x_{j+1}}{\sqrt{x_{j+1}}} \hat{F}\left(\frac{\log(x_j/a)}{2\pi\Delta}\right)\right) \frac{4\sqrt{x_{j+1}}}{\log x_{j+1}} \\ &\leq 4\hat{F}(-1) + \frac{4}{\sqrt{a/(e^{4\pi\Delta})}} \sum_{j=1}^{L-1} \hat{F}\left(\frac{\log(x_j/a)}{2\pi\Delta}\right) \sqrt{x_j} \\ &\leq 4\hat{F}(-1) + \frac{4e^{2\pi\Delta}}{\sqrt{a}} \int_{x_L}^{x_0} \hat{F}\left(\frac{\log(t/a)}{2\pi\Delta}\right) dt \end{aligned} \quad (6.14)$$

$$\begin{aligned}
&\leq 4\hat{F}(-1) + 4\sqrt{a}e^{2\pi\Delta}(2\pi\Delta) \int_{-\lambda^{-1}}^{-1} \hat{F}(y) dy \\
&\leq 4\hat{F}(-1) + 4\sqrt{a}(2\pi\Delta) \int_{-\lambda^{-1}}^{-1} \hat{F}(y) dy + 8\sqrt{a}(2\pi\Delta)^2 \hat{F}(-1)(\lambda^{-1} - 1).
\end{aligned}$$

Combining (6.13) and (6.14) we conclude that

$$\begin{aligned}
&\sum_{1 < |\frac{\log p/a}{2\pi\Delta}| \leq \lambda^{-1}} \frac{\log p}{\sqrt{p}} \hat{F}\left(\frac{\log(p/a)}{2\pi\Delta}\right) \\
&\leq 8\hat{F}(1) + 4\sqrt{a}(2\pi\Delta) \int_{[-1,1]^c} \hat{F}(y) dy + 24\sqrt{a}(2\pi\Delta)^2 \hat{F}(1)(\lambda^{-1} - 1) \\
&\leq 0.886 + 4\sqrt{a}(2\pi\Delta) \int_{[-1,1]^c} \hat{F}(y) dy.
\end{aligned} \tag{6.15}$$

Here we have used the estimate $8\hat{F}(1) \leq 0.885$ along with the inequalities $a \leq 4x$ and $\log(1+y) \leq y$ for $y \geq 0$ in (5.7) and (5.8) to see that $\sqrt{a}(2\pi\Delta)^2 \leq \frac{c^2 \log^2 x}{\sqrt{x}} \leq 10^{-7}$ for $c \leq 1$ and $x \geq 4 \cdot 10^{18}$ and thus that $24\sqrt{a}(2\pi\Delta)^2 \hat{F}(1)(\lambda^{-1} - 1) \leq 0.001$.

Since $\text{supp}(\hat{F}) \subset [-\lambda^{-1}, \lambda^{-1}] \subset [-2, 2]$ and $|\hat{F}(y)| \leq \pi\lambda/4 < 1$, the contribution from the prime powers $n = p^k$ with $k \geq 2$ to the sum over n in (6.5) is

$$\begin{aligned}
&\leq \sum_{k \geq 2} \sum_{\substack{ae^{-4\pi\Delta} \leq n \leq ae^{4\pi\Delta} \\ n = p^k}} \frac{\log p}{\sqrt{n}} \\
&\leq \frac{\log(ae^{-4\pi\Delta})}{2\sqrt{ae^{-4\pi\Delta}}} \sum_{\substack{k \geq 2 \\ k \leq \log(ae^{4\pi\Delta})/\log 2}} (1 + a^{\frac{1}{k}}(e^{4\pi\Delta/k} - e^{-4\pi\Delta/k})),
\end{aligned}$$

where we used a trivial estimate for the total number of k th powers that can lie in the interval $[ae^{-4\pi\Delta}, ae^{4\pi\Delta}]$. This is readily bounded by

$$\begin{aligned}
&\leq \frac{\log(ae^{-4\pi\Delta})}{2\sqrt{ae^{-4\pi\Delta}}} (2\sqrt{a}(e^{2\pi\Delta} - e^{-2\pi\Delta})) \frac{\log(ae^{4\pi\Delta})}{\log 2} \\
&= \frac{\log(ae^{-4\pi\Delta}) \log(ae^{4\pi\Delta})}{a e^{-2\pi\Delta} \log 2} (ae^{2\pi\Delta} - ae^{-2\pi\Delta}) \\
&= \frac{\log(ae^{-4\pi\Delta}) \log(ae^{4\pi\Delta})}{a e^{-2\pi\Delta} \log 2} c\sqrt{x} \log x < \frac{2(\log a + 1)^3}{\log 2 \sqrt{a}} < 0.001.
\end{aligned} \tag{6.16}$$

6.3. Finishing the proof. Note that $ae^{-2\pi\Delta/\lambda} \geq x/2 \geq 2 \cdot 10^{18}$ and thus

$$\frac{3}{2}\Delta \left(\frac{e^{2\pi\Delta/\lambda}}{a} \right)^{5/2} \leq \frac{0.001}{2\pi}.$$

Combining this estimate with (6.5), (6.9), (6.15), and (6.16), (after multiplying both sides by 2π) we derive that

$$\sqrt{a} (2\pi\Delta) \leq \log \left(\frac{1}{2\pi\Delta} \right) \|F\|_1 + 4\sqrt{a} (2\pi\Delta) \int_{[-1,1]^c} \hat{F}(y) dy + 0.958.$$

Rearranging and dividing by $\|F\|_1 = \lambda \|H\|_1 = 0.83337\dots$ we obtain (with $J(F)$ defined in (6.12))

$$J(F) \sqrt{a} (2\pi\Delta) \leq \log \left(\frac{1}{2\pi\Delta} \right) + 1.16. \quad (6.17)$$

Since $a \geq x$, $1 \geq c \geq \frac{1}{2}$, and $\log(1+y) \geq y - \frac{y^2}{2}$ for $y \geq 0$, we derive from (5.7) and (5.8) that

$$\sqrt{a} (2\pi\Delta) \geq \frac{c}{2} \log x - \frac{c^2 \log^2 x}{4\sqrt{x}} \geq \frac{c}{2} \log x - 0.001.$$

Using the inequalities $c \geq \frac{1}{2}$ and $\log(1+y) \geq y \log 2$, which holds for $0 \leq y \leq 1$, it follows that $2\pi\Delta = \frac{1}{2} \log \left(1 + c \frac{\log x}{\sqrt{x}} \right) \geq \frac{c \log 2}{2} \frac{\log x}{\sqrt{x}}$ and therefore

$$\begin{aligned} \log \left(\frac{1}{2\pi\Delta} \right) &\leq \log \left(\frac{2}{c \log 2} \frac{\sqrt{x}}{\log x} \right) \\ &\leq \frac{1}{2} \log x - \log \left(\frac{\log 2}{4} \log(4 \cdot 10^{18}) \right) \leq \frac{1}{2} \log x - 2. \end{aligned}$$

Inserting these estimates into (6.17), we derive that

$$\frac{c J(F)}{2} \log x \leq \frac{1}{2} \log x - \frac{1}{2}.$$

This is not possible if $c = \frac{1}{J(F)} < \frac{22}{25}$. Hence there must be a prime in the interval $[x, x + \frac{22}{25} \sqrt{x} \log x]$.

7. Concluding remarks

There are several related extremal problems in Fourier analysis that could be the sources of further investigation. We briefly discuss a few of these here.

7.1. Multidimensional analogues. The corresponding versions of the extremal problems (1.1)–(1.4) in \mathbb{R}^d arise as natural generalizations. The compact interval $[-1, 1] \subset \mathbb{R}$ could be replaced by any convex, compact, and symmetric set $K \subset \mathbb{R}^d$, for instance. Of those, the most basic ones are certainly the cube $Q = [-1, 1]^d$ and the unit Euclidean ball $B = \{x \in \mathbb{R}^d; |x| \leq 1\}$. The same ideas used here could be

applied to show the existence of extremizers in this general situation. By averaging over the group of symmetries of K , one can show that extremizers admit, without loss of generality, these symmetries. Note that a crucial step in our proof of the uniqueness of extremizers in Section 3 (for the bandlimited problem (1.2)) was the ability to write a nonnegative function with Fourier transform supported in $2K$ as the square of a function whose Fourier transform is supported in K . In general, this decomposition is not available for any given K , but in the case of the unit ball B , with respect to radial functions, this statement holds. This was proved, for instance, in [9, 28], exploring the connection with the theory of Hilbert spaces of entire functions of L. de Branges. Hence, in dimension $d \geq 1$ and for $K = B$, one has indeed the uniqueness of radial extremizers (up to multiplication by a complex scalar) for the multidimensional version of (1.2). Letting $\mathcal{C}_{d,K}(A)$ denote the sharp constant in the multidimensional version of (1.1)–(1.2), one can show that $\mathcal{C}_{d,Q}(\infty) = \mathcal{C}(\infty)^d$, and a tensor product of one-dimensional extremizers is an extremizer for the multivariable problem. In the general case, one has $\mathcal{C}_{d,K}(\infty) \leq \text{vol}(K)$. A lower bound for $\mathcal{C}_{d,K}(\infty)$ may come, for instance, from the solution of the “one-delta problem for K ”, which is the same problem as (1.2) with the additional constraint that $F \geq 0$. Such problem is also vastly open, having been solved only in a few particular cases such as the cube Q and the ball B (see the discussion in [5, 22, 32]). It would be interesting to have refined upper and lower bounds for all of these extremal problems, as we have here in our Theorems 1.1 and 1.2.

7.2. Sphere packing. The following extremal problem in Fourier analysis was proposed by Cohn and Elkies [12] in connection to the sphere packing problem. Find

$$C = \sup_{\substack{F \in \mathcal{E}_d^+ \\ F \neq 0}} \frac{F(0)}{\widehat{F}(0)}, \quad (7.1)$$

where the supremum is taken over the class \mathcal{E}_d^+ of real-valued, continuous, and integrable functions $F: \mathbb{R}^d \rightarrow \mathbb{R}$ with $F \geq 0$ and $\widehat{F}(y) \leq 0$ for $|y| \geq 1$. This is the multidimensional analogue of our extremal problem (1.4) with the additional constraint that $F \geq 0$. By averaging over the group of rotations $SO(d)$ we may restrict the search to radial functions and by following the outline of §2.3 and §2.4 we obtain the next result.²

Proposition 7.1. *There exists a radial extremizer for (7.1).*

As a matter of fact, Cohn and Elkies [12] proposed this optimization problem over the more restrictive class of admissible functions $F: \mathbb{R}^d \rightarrow \mathbb{R}$ such that $|F|$ and $|\widehat{F}|$ are bounded above by constant times $(1 + |x|)^{-d-\delta}$ for some $\delta > 0$. Standard

²This result has been previously communicated by E. Carneiro and Alvaro A. Gomez (with a slightly different proof than the one presented here), as part of the M.Sc. thesis of the latter under the supervision of the former.

approximation arguments show that the sharp constant over this restricted class is the same C in (7.1), although extremizers of (7.1), in principle, need not have this particular decay. In addition to dimension $d = 1$, the value of the sharp constant in (7.1) is known only in dimensions $d = 8$ and 24 (see [43] and [13], respectively). The extremizers found by Viazovska in [43] and by Cohn, Kumar, Miller, Radchenko, and Viazovska in [13] are indeed radial Schwartz functions.

Acknowledgements. The authors are thankful to Andrés Chirre for very helpful discussions during the preparation of this work, to Dan Goldston for encouragement, to Dimitar Dimitrov for bringing references [2, 24, 25, 40] to our attention, and to Dmitry Gorbachev for bringing reference [29] to our attention (the latter happened after the manuscript had been accepted for publication). E.C. acknowledges support from CNPq-Brazil, FAPERJ-Brazil, and the Fulbright Junior Faculty Award, and is also thankful to Stanford University for the support and warm hospitality. M.B.M. was supported in part by NSA Young Investigator Grants H98230-15-1-0231 and H98230-16-1-0311, and thanks Stanford University for hosting him on two research visits. K.S. was partially supported by NSF grant DMS 1500237, and a Simons Investigator grant from the Simons Foundation.

References

- [1] N. I. Achieser, *Theory of approximation*. Translated by Charles J. Hyman, Frederick Ungar Publishing Co., New York, 1956. Zbl 0072.28403 MR 95369
- [2] N. N. Andreev, S. V. Konyagin, and A. Yu. Popov, Extremum problems for functions with small support, *Math. Notes*, **60** (1996), no. 3-4, 241–247 (1997); translated from *Mat. Zametki*, **60** (1996), no. 3, 323–332, 478. Zbl 0911.42001 MR 1428847
- [3] J. Bourgain, L. Clozel, and J.P. Kahane, Principe d’Heisenberg et fonctions positives. (French) [The Heisenberg principle and positive functions], *Ann. Inst. Fourier (Grenoble)*, **60** (2010), no. 4, 1215–1232. Zbl 1298.11105 MR 2722239
- [4] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Universitext, Springer, New York, 2011. Zbl 1220.46002 MR 2759829
- [5] G. Bianchi and M. Kelly, A Fourier analytic proof of the Blaschke-Santaló inequality, *Proc. Amer. Math. Soc.*, **143** (2015), no. 11, 4901–4912. Zbl 1322.52007 MR 3391048
- [6] E. Carneiro, V. Chandee, F. Littmann, and M. B. Milinovich, Hilbert spaces and the pair correlation of zeros of the Riemann zeta-function, *J. Reine Angew. Math.*, **725** (2017), 143–182. Zbl 1362.11078 MR 3630120
- [7] E. Carneiro, V. Chandee, and M. B. Milinovich, Bounding $S(t)$ and $S_1(t)$ on the Riemann hypothesis, *Math. Ann.*, **356** (2013), no. 3, 939–968. Zbl 1325.11084 MR 3063902
- [8] E. Carneiro and A. Chirre, Bounding $S_n(t)$ on the Riemann hypothesis, *Math. Proc. Cambridge Philos. Soc.*, **164** (2018), no. 2, 259–283. Zbl 06903441 MR 3778242
- [9] E. Carneiro and F. Littmann, Extremal functions in de Branges and Euclidean spaces, *Adv. Math.*, **260** (2014), 281–349. Zbl 1295.41015 MR 3209354

- [10] E. Carneiro, F. Littmann, and J. D. Vaaler, Gaussian subordination for the Beurling-Selberg extremal problem, *Trans. Amer. Math. Soc.*, **365** (2013), no. 7, 3493–3534. Zbl 1276.41018 MR 3042593
- [11] V. Chandee and K. Soundararajan, Bounding $|\zeta(\frac{1}{2} + it)|$ on the Riemann hypothesis, *Bull. London Math. Soc.*, **43** (2011), no. 2, 243–250. Zbl 1238.11078 MR 2781205
- [12] H. Cohn and N. Elkies, New upper bounds on sphere packings. I, *Ann. of Math. (2)*, **157** (2003), no. 2, 689–714. Zbl 1041.52011 MR 1973059
- [13] H. Cohn, A. Kumar, S. D. Miller, D. Radchenko, and M. Viazovska, The sphere packing problem in dimension 24, *Ann. of Math. (2)*, **185** (2017), no. 3, 1017–1033. Zbl 1370.52037 MR 3664817
- [14] H. Cramér, Some theorems concerning prime numbers, *Ark. Mat. Astron. Fysik.*, **15** (1921), no. 5, 33pp. Zbl 47.0156.01
- [15] H. Cramér, On the order of magnitude of the difference between consecutive prime numbers, *Acta Arith.*, **2** (1936), no. 1, 23–46. Zbl 0015.19702
- [16] H. Davenport, *Multiplicative number theory*. Third edition. Revised and with a preface by Hugh L. Montgomery, Graduate Texts in Mathematics, 74, Springer-Verlag, New York, 2000. Zbl 1002.11001 MR 1790423
- [17] D. L. Donoho and B. F. Logan, Signal recovery and the large sieve, *SIAM J. Appl. Math.*, **52** (1992), no. 2, 577–591. Zbl 0768.42001 MR 1154788
- [18] D. L. Donoho and P. Stark, Uncertainty principles and signal recovery, *SIAM J. Appl. Math.*, **49** (1989), no. 3, 906–931. Zbl 0689.42001 MR 997928
- [19] A. Dudek, On the Riemann hypothesis and the difference between primes, *Int. J. Number Theory*, **11** (2015), no. 3, 771–778. Zbl 1317.11095 MR 3327842
- [20] A. Dudek, L. Grenié, and G. Molteni, Primes in explicit short intervals on RH, *Int. J. Number Theory*, **12** (2016), no. 5, 1391–1407. Zbl 1356.11065 MR 3498632
- [21] D. A. Goldston, On a result of Littlewood concerning prime numbers. II, *Acta Arith.*, **43** (1983), no. 1, 49–51. Zbl 0478.10025 MR 730847
- [22] F. Gonçalves, M. Kelly, and J. Madrid, One-sided band-limited approximations of some radial functions, *Bull. Braz. Math. Soc. (N.S.)*, **46** (2015), no. 4, 563–599. Zbl 1339.41026 MR 3436559
- [23] F. Gonçalves, D. Oliveira e Silva, and S. Steinerberger, Hermite polynomials, linear flows on the torus, and an uncertainty principle for roots, *J. Math. Anal. Appl.*, **451** (2017), no. 2, 678–711. Zbl 1373.33018 MR 3624763
- [24] D. V. Gorbachev, A Sharpening of the Taikov lower bound in the inequality between the C - and L - norms for trigonometric polynomials, *Mat. Zametki*, **74** (2003), no. 1, 132–134; translation in *Math. Notes*, **74** (2003), no. 1-2, 123–126. Zbl 1062.41008 MR 2010684
- [25] D. V. Gorbachev, An integral problem of Konyagin and the (C,L) -constants of Nikol'skii, *Trudy Inst. Mat. i Mekh. UrO RAN*, **11** (2005), no. 2, 72–91. Zbl 1126.41016 MR 2200225
- [26] D. R. Heath-Brown, Gaps between primes, and the pair correlation of zeros of the zeta function, *Acta Arith.*, **41** (1982), no. 1, 85–99. Zbl 0414.10044 MR 667711
- [27] D. R. Heath-Brown and D. A. Goldston, A note on the differences between consecutive primes, *Math. Ann.*, **266** (1984), no. 3, 317–320. Zbl 0514.10031 MR 730173

- [28] J. Holt and J. D. Vaaler, The Beurling-Selberg extremal functions for a ball in the Euclidean space, *Duke Math. J.*, **83** (1996), no. 1, 203–247. Zbl 0859.30029 MR 1388849
- [29] L. Hörmander and B. Bernhardsson, An extension of Bohr’s inequality, in *Boundary value problems for partial differential equations and applications*, 179–194, RMA Res. Notes Appl. Math., 29, Masson, Paris, 1993. Zbl 0803.41030 MR 1260445
- [30] H. Iwaniec, On the Brun–Titchmarsh theorem, *J. Math. Soc. Japan*, **34** (1982), no. 1, 95–123. Zbl 0486.10033 MR 639808
- [31] H. Iwaniec and E. Kowalski, *Analytic number theory*, American Mathematical Society Colloquium Publications, 53, American Mathematical Society, Providence, RI, 2004. Zbl 1059.11001 MR 2061214
- [32] M. Kelly, Some Inequalities in Fourier Analysis and Applications, Ph.D. dissertation, The University of Texas at Austin, 2013.
- [33] Y. Lamzouri, X. Li, Xiannan, and K. Soundararajan, Conditional bounds for the least quadratic non-residue and related problems, *Math. Comp.*, **84** (2015), no. 295, 2391–2412. Zbl 1326.11058 MR 3356031
- [34] B. Logan, Properties of high-pass signals, Ph.D. dissertation, Columbia University, New York, 1965.
- [35] H. L. Montgomery and R. C. Vaughan, The large sieve, *Mathematika*, **20** (1973), 119–134. Zbl 0296.10023 MR 374060
- [36] J. H. Mueller, On the difference between consecutive primes, in *Recent progress in analytic number theory. Vol. 1 (Durham, 1979)*, 269–273, Academic Press, London-New York, 1981. Zbl 0457.10022 MR 637352
- [37] B. Sz. Nagy, Über gewisse Extremalfragen bei transformierten trigonometrischen Entwicklungen. II. Nichtperiodischer Fall, *Ber. Math.-Phys. Kl. Sachs Akad. Wiss. Leipzig*, **91** (1939), 3–24. Zbl 0021.40103
- [38] T. Oliveira e Silva, S. Herzog, and S. Pardi, Empirical verification of the even Goldbach conjecture and computation of prime gaps up to $4 \cdot 10^{18}$, *Math. Comp.*, **83** (2014), no. 288, 2033–2060. Zbl 1290.11161 MR 3194140
- [39] O. Ramaré and Y. Saouter, Short effective intervals containing primes, *J. Number Theory*, **98** (2003), no. 1, 10–33. Zbl 1032.11038 MR 1950435
- [40] L. V. Taikov, One scope of extremal problems for trigonometric polynomials, *Uspekhi Mat. Nauk*, **20** (1965), no. 3 (123), 205–211. Zbl 0141.25301 MR 185361
- [41] T. Trudgian, An improved upper bound for the argument of the Riemann zeta-function on the critical line. II, *J. Number Theory*, **134** (2014), 280–292. Zbl 1284.11119 MR 3111568
- [42] J. D. Vaaler, Some extremal functions in Fourier analysis, *Bull. Amer. Math. Soc. (N.S.)*, **12** (1985), no. 2, 183–215. Zbl 0575.42003 MR 776471
- [43] M. Viazovska, The sphere packing problem in dimension 8, *Ann. of Math. (2)*, **185** (2017), no. 3, 991–1015. Zbl 1373.52025 MR 3664816

Received August 14, 2017

E. Carneiro, ICTP – The Abdus Salam International Centre for Theoretical Physics,
Strada Costiera, 11, I - 34151, Trieste, Italy; and

IMPA – Instituto Nacional de Matemática Pura e Aplicada,
Estrada Dona Castorina, 110, Rio de Janeiro, RJ 22460-320, Brazil

E-mail: carneiro@ictp.it; carneiro@impa.br

M. B. Milinovich, Department of Mathematics, University of Mississippi,
University, MS 38677, USA

E-mail: mbmilino@olemiss.edu

K. Soundararajan, Department of Mathematics, Stanford University,
Stanford, CA 94305, USA

E-mail: ksound@stanford.edu