Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 94 (2019)

Heft: 4

Artikel: Aut-invariant norms and Aut-invariant quasimorphisms on free and

surface groups

Autor: Brandenbursky, Michael / Marcinkowski, Michal

DOI: https://doi.org/10.5169/seals-847027

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Aut-invariant norms and Aut-invariant quasimorphisms on free and surface groups

Michael Brandenbursky and Michał Marcinkowski*

Abstract. Let \mathbf{F}_n be the free group on n generators and $\mathbf{\Gamma}_g$ the surface group of genus g. We consider two particular generating sets: the set of all primitive elements in \mathbf{F}_n and the set of all simple loops in $\mathbf{\Gamma}_g$. We give a complete characterization of distorted and undistorted elements in the corresponding Aut-invariant word metrics. In particular, we reprove Stallings theorem and answer a question of Danny Calegari about the growth of simple loops. In addition, we construct infinitely many quasimorphisms on \mathbf{F}_2 that are $\mathrm{Aut}(\mathbf{F}_2)$ -invariant. This answers an open problem posed by Miklós Abért.

Mathematics Subject Classification (2010). 57-xx.

Keywords. Free groups, mapping class groups, quasi-morphisms, invariant norms.

1. Introduction

Let G be a group and let $\operatorname{Aut}(G)$ be the group of all automorphisms of G. A function $|\cdot|: G \to [0, \infty)$ is called a norm if it satisfies the following conditions for all $g, h \in G$:

- |g| = 0 if and only if $g = 1_G$;
- $|g| = |g^{-1}|$;
- |gh| < |g| + |h|.

In this paper we study norms that are $\operatorname{Aut}(G)$ -invariant, i.e. for each $x \in G$ and $\psi \in \operatorname{Aut}(G)$ we have $|x| = |\psi(x)|$. An example of such a norm is the word norm $|\cdot|_S$ defined by $\operatorname{Aut}(G)$ -invariant generating set S, that is:

$$|x|_S = \min\{n : x = s_1, \dots, s_n, \text{ where } s_i \in S \text{ for each } i\}.$$

Under mild assumptions, up to bi-Lipschitz equivalence, there is only one $\operatorname{Aut}(G)$ -invariant word norm. We denote it by $|\cdot|_{\operatorname{Aut}}$ and call it the Aut-norm of G. We focus

^{*}Both authors were partially supported by GIF-Young Grant number I-2419-304.6/2016 and by SFB 1085 "Higher Invariants" funded by DFG.

on two types of groups: surface groups, where the Aut-norm counts the minimal number of simple loops needed to express an element in a group, and free groups, where the Aut-norm counts the minimal number of primitive elements needed to express an element in a group.

Let $x \in G$. Recall that x is *undistorted* with respect to $|\cdot|$ if there exists C>0 such that $|x^n|>Cn$. Otherwise it is *distorted*. Note that this notion depends only on the bi-Lipschitz class of $|\cdot|$. In this paper we characterize distorted and undistorted elements in surface and free groups. The main idea is to find an appropriate quasimorphism on G. More precisely, in order to show that $x \in G$ is undistorted in $|\cdot|_{\text{Aut}}$, it is enough to find a homogeneous quasimorphism which is non-zero on x but is bounded on some Aut(G)-invariant generating set S. This strategy was previously used in the context of conjugation invariant norms, see e.g. [5,6].

We say that G satisfies bq-dichotomy with respect to the Aut-norm, if for every element $x \in G$, either the cyclic group $\langle x \rangle$ is bounded in the Aut-norm, or we can find a homogeneous quasimorphism which does not vanish on x but is bounded on some $\operatorname{Aut}(G)$ -invariant generating set. That is, all undistorted elements are detected by appropriate quasimorphisms.

The main result of this paper is presented below, i.e. we prove the following theorem (see Theorems 3.6 and 4.12 in the text), which, in particular, answers (see Corollary 4.14) a question of Danny Calegari from 2007 (see [9, Question 1.6]) and gives a simple proof of Stallings theorem [25, Theorem 2.4] on Whitehead graphs of separable elements.

Theorem 1. Surface groups and free groups satisfy bq-dichotomy with respect to their Aut-norms. Moreover, in both cases, there is an explicit characterization of undistorted elements.

This theorem has an application to geodesics on closed hyperbolic surfaces. More precisely, we show that for every neighborhood U of a point p that lies on a closed simple non-separating geodesic l, there is another simple closed geodesic l' which passes through U, see Theorem 4.15.

In addition, the methods we use allow us to prove the following theorem (see Corollary 5.6 in the text) which answers a question of Miklós Abért from 2010 (see [1, Question 47]) in the case of \mathbb{F}_2 .

Theorem 2. The space of homogeneous $Aut(\mathbf{F}_2)$ -invariant quasimorphisms on \mathbf{F}_2 is infinite dimensional.

As a corollary we provide an infinite dimensional space of quasimorphisms on \mathbb{F}_2 where each quasimorphism can not be expressed as a finite sum of counting quasimorphisms, see Remark 5.7.

Acknowledgements. We would like to thank Mladen Bestvina for answering our questions, and Danny Calegari and Hugo Parlier for comments on the early draft of our paper.

Part of this work was done during the authors' stay at the University of Regensburg and the second author's stay at the Ben-Gurion University. We wish to express our gratitude to both places for the support and excellent working conditions.

2. Preliminaries

2.1. Aut-invariant norm on free groups. Let \mathbf{F}_n be the free group of rank n. An element b is called *primitive*, if it is an element of some free basis of \mathbf{F}_n . Note that if b is primitive and $\psi \in \operatorname{Aut}(\mathbf{F}_n)$, then $\psi(b)$ is primitive and each base element has a form $\psi(b)$ for some fixed b. In other words, the set of all primitive elements in \mathbf{F}_n is a single orbit of the $\operatorname{Aut}(\mathbf{F}_n)$ action.

Given an element of \mathbf{F}_n where $n \geq 3$, it is difficult to decide if it is primitive or not. In his celebrated papers ([26, 27]), J. H. C. Whitehead provided an algorithmic method to solve this problem. However, the time complexity of this algorithm seems to be ineffective for large n (see [21] for an attempt to find fast algorithms). It is worth to note that in the case of \mathbf{F}_2 the situation is completely different. There is a quadratic in time algorithm which checks whether an element of \mathbf{F}_2 is primitive or not. It can be extracted from [22].

We consider the following norm:

```
|x|_p = \min\{n \mid x = b_1 \dots b_n, \text{ where } b_i \text{ is a primitive element in } \mathbb{F}_n\},\
```

which we call the *primitive norm* of x. A priori it is not clear whether this norm is unbounded. Indeed, for the free group of infinite rank, the norm of any element is at most 2. In case of arbitrary finite rank, the unboundedness was proven in [2] by exhibiting a relevant non-trivial homogeneous quasimorphism. Our Theorem 3.6 may be viewed as a generalization of this result. In the proof we use Whitehead graphs and Whitehead automorphisms, and as a corollary we obtain a result due to Stallings (See Corollary 3.7) on Whitehead graphs of separable elements.

2.2. Aut-invariant norm on surface groups. Let us denote by S_g the oriented closed surface of genus g and let $\star \in S_g$ be a base point. Let $\Gamma_g = \pi_1(S_g, \star)$. An element s of Γ_g is called *simple*, if it can be represented by a based loop with no self-intersection points (such a loop is called simple). Simple elements generate Γ_g , thus we can consider the following norm:

```
|x|_s = \min\{n \mid x = s_1 \dots s_n, \text{ where } s_i \text{ is a simple element in } \Gamma_g\}.
```

We call $|\cdot|_s$ the *simple loops norm*. Danny Calegari proved that every non-simple element $x \in \Gamma_g$ is undistorted in this norm [9], leaving the case of simple elements open. Theorem 4.12 solves this case. In particular, we show that an element which is represented by simple separating closed curve is undistorted.

Recently, Erlandsson considered generating sets consisting of simple loops and obtained interesting results about intersection numbers [11,12]. However, generating sets she considers are finite and we consider infinite generating sets.

Let us put an arbitrary hyperbolic metric on S_g . It is known that in every homotopy class of a based loop there is a unique based closed geodesic. Using Theorem 4.12 we draw some conclusions concerning the behavior of closed geodesic in S_g , see Subsection 4.4.

- **2.3. Generalization:** H-norms. Let G be a group and $H < \operatorname{Aut}(G)$ be a subgroup of the group of automorphisms of G. The group H acts on G from the left. We say that:
- G is H-generated by $S \subset G$ if

$$\bar{S} = \{h(s) : s \in S \cup S^{-1}, h \in H\}$$

generates G.

- H-invariant subset \overline{S} of G is H-finite if \overline{S}/H is finite, i.e. \overline{S} is a sum of finitely many H-orbits.
- G is H-finitely generated if there exists an H-finite subset of G generating G. Equivalently, if there exists a finite set $S \subset G$ such that

$$\overline{S} = \{h(s) : s \in S \cup S^{-1}, h \in H\}$$

generates G.

Having an H-finite set \overline{S} which generates G, we consider the word norm $|\cdot|_{\overline{S}}$ on G defined by \overline{S} . Note, that $|\cdot|_{\overline{S}}$ is H-invariant, i.e. $|h(g)|_{\overline{S}} = |g|_{\overline{S}}$ for $h \in H$ and $g \in G$.

The norm $|\cdot|_{\overline{S}}$ depends on the choice of H-finite set \overline{S} . However, as long as \overline{S} is H-finite, $|\cdot|_{\overline{S}}$ belongs to the same bi-Lipschitz equivalence class. We denote this equivalence class by $|\cdot|_H$. The norm $|\cdot|_H$ is maximal among all H-invariant norms on G, namely: for every H-invariant norm $|\cdot|$, there exists C>0 such that $|x|< C|x|_H$ for every $x\in G$.

Examples include (we always assume that G is H-finitely generated):

- (1) H is trivial, then $|\cdot|_H$ is the standard word norm.
- (2) H is the group of inner automorphisms of G, then $|\cdot|_H$ is the conjugation invariant word norm, see e.g. [5].
- (3) H is the full automorphism group, then $|\cdot|_H$ is called the Aut-norm on G.

Let us show that the primitive norm and the simple loops norm are examples of Aut-norms on free and surface groups. It is clear that the primitive norm is an Aut-norm since the set of primitive elements in \mathbb{F}_n equals to the set

$$\{\psi(b) \mid \psi \in \operatorname{Aut}(\mathbf{F}_n)\},\$$

where b is an arbitrary chosen primitive element. Thus this set is a single $Aut(\mathbf{F}_n)$ orbit. In the case of the simple loops norm we use the Baer–Dehn–Nielsen theorem
which states:

$$\operatorname{Aut}(\Gamma_g) \cong \operatorname{MCG}^{\pm}(S_g, \star).$$

Now it follows from the classification of surfaces, that $\operatorname{Aut}(\Gamma_g)$ -orbit of a simple element $s \in \Gamma_g$ is determined by the homeomorphism type of the surface $S_g \setminus \gamma_s$, where γ_s is the corresponding simple loop. Since there are only finitely many homeomorphism types of such surfaces, the set of simple elements consists of finitely many $\operatorname{Aut}(\Gamma_g)$ -orbits.

2.4. Quasimorphisms and distortion. Let us recall a notion of a quasimorphism. A function $q: G \to \mathbf{R}$ is called a *quasimorphism* if there exists D such that

$$|q(a) - q(ab) + q(b)| < D$$

for all $a, b \in G$. The minimal such D is called the *defect* of q and denoted by D_q . A quasimorphism q is *homogeneous* if $q(x^n) = nq(x)$ for all $n \in \mathbb{Z}$ and all $x \in G$. Homogeneous quasimorphisms are constant on conjugacy classes, i.e. $q(x) = q(yxy^{-1})$ for all $x, y \in G$. We refer to [10] for further details.

Lemma 2.1. Let G be a group and let $H < \operatorname{Aut}(G)$. Assume that $\overline{S} \subset G$ is an H-finite subset which generates G. Let $x \in G$. If there exists a homogeneous quasimorphism $q: G \to \mathbb{R}$ such that $q(x) \neq 0$ and q is bounded on \overline{S} , then x is undistorted in the norm $|\cdot|_H$.

Proof. Let $w \in G$ and C be such that |q(s)| < C for all $s \in \overline{S}$. We can write w in the following form: $w = s_1 \dots s_n$ where $n = |w|_H$ and $s_i \in \overline{S}$. The following inequality holds:

$$|q(w)| \le \sum_{1}^{n} |q(s_i)| + (n-1)D_q \le nC + (n-1)D_q \le (C+D_q)|w|_H.$$

We apply this to x^n and get

$$n|q(x)| = |q(x^n)| \le (C + D_q)|x^n|_H.$$

Since $q(x) \neq 0$, $|x^n|_H$ growths linearly with n.

Remark 2.2. In Lemma 2.1, instead of assuming that q is homogeneous, it is enough to assume that $q(x^n)$ growths linearly.

Corollary 2.3. Assume that q is an H-invariant homogeneous quasimorphism and $q(x) \neq 0$. Then x is undistorted in $|\cdot|_H$.

Let $q: G \to \mathbb{R}$ be a quasimorphism. We define

$$\overline{q}(a) = \lim_{n \to \infty} q(a^n)/n$$
.

Straight-forward computations show that \overline{q} is a homogeneous quasimorphism, see e.g. [10]. The following lemma relates general quasimorphisms with homogeneous quasimorphisms.

Lemma 2.4 ([10]). Let $q: G \to \mathbb{R}$ be a quasimorphism such that $q(x^n)$ growths linearly. Then a homogeneous quasimorphism $\overline{q}: G \to \mathbb{R}$, called the homogenization of q, satisfies $|q(a) - \overline{q}(a)| \le D_q$ for all $q \in G$, and $\overline{q}(x) \ne 0$.

3. Aut-norms on free groups

In this section we use counting quasimorphisms in order to investigate distortion in free groups.

3.1. Whitehead graph. Let **F** be a free group (possibly not finitely generated). Let $w \in \mathbf{F}$ and let B be any free basis of **F**. Here we use the convention that if $b \in B$, then $b^{-1} \notin B$. Suppose that w is a cyclically reduced word with respect to the basis B and $w = w_1 w_2 \dots w_n$ is the reduced expression of w in B. We define the Whitehead graph $\operatorname{Wh}_B(w)$ as follows: for each element $b \in B$ we have two vertices in $\operatorname{Wh}_B(w)$ labeled by b and b^{-1} . For every two consecutive letters w_i, w_{i+1} in w, we draw an edge from w_i to w_{i+1}^{-1} . We regard w_n, w_0 as being consecutive in w, that is, we have an edge from w_n to w_0^{-1} . If w is not reduced with respect to B, then denote by $r_B(w)$ the unique cyclically reduced word in the conjugacy class of w. We define $\operatorname{Wh}_B(w) := \operatorname{Wh}_B(r_B(w))$.




Figure 1. $Wh_{\{a,b\}}(aba^{-1}b^{-1})$.

Theorem 3.1 (Whitehead [26]). If $b \in \mathbf{F}$ is a primitive element and $\operatorname{Wh}_B(b)$ is connected, then $\operatorname{Wh}_B(b)$ has a cut-vertex.

A vertex v of a graph is a *cut-vertex* if after removing v, the graph has more connected components. In the example below a, b, a^{-1} and b^{-1} are cut-vertices.




Figure 2. Wh $\{a,b,c\}$ $(aba^{-1}b^{-1}c)$.

Lemma 3.2. Let B be a free basis of \mathbb{F} and let $b \in \mathbb{F}$ be a cyclically reduced base element. Assume that $x \in \mathbb{F}$ is a cyclically reduced word such that its Whitehead graph $\operatorname{Wh}_B(x)$ is connected and has no cut-vertices. Then the reduced expression of b in the basis B does not contain the reduced expressions of x^2 and of x^{-2} as subwords.

Proof. Let us fist notice, that if a reduced expression of b contains the reduced expression of x^2 , then $\operatorname{Wh}_B(b)$ contains $\operatorname{Wh}_B(x)$ as a subgraph. Indeed, every edge of $\operatorname{Wh}_B(x)$ is given by two cyclically consecutive letters in x. If x^2 is a subword of b, we can find all those edges in $\operatorname{Wh}_B(b)$. $\operatorname{Wh}_B(x)$ has no cut-vertices, it is connected and has the same vertex set as $\operatorname{Wh}_B(b)$. Thus if $\operatorname{Wh}_B(x)$ is a subgraph of $\operatorname{Wh}_B(b)$, then $\operatorname{Wh}_B(b)$ is a connected graph with no cut-vertices. This contradicts the Whitehead Theorem 3.1.

Remark 3.3. In Lemma 3.2, the assumption that b is cyclically reduced can not be omitted. For general b one can show that the number of occurrences of x^2 minus the number of occurrences of x^{-2} is bounded by $2|x|_B$. Here $|\cdot|_B$ is the word metric defined by B. We would like to add that a variation of Lemma 3.2 already appeared in [2].

3.2. Counting quasimorphisms. Let $w \in \mathbb{F}$ and let B be a basis of \mathbb{F} . Having an element $x \in \mathbb{F}$, we can write x and w as reduced expressions in the base B. We define $C_w(x)$ to be the number of occurrences of w as a subword of x (the subwords can overlap). In [7] Brooks proved that the following function

$$Br_w(x) = C_w(x) - C_w(x^{-1})$$

is a quasimorphism. See also [24]. Usually we suppress the basis B from the notation.

Lemma 3.4. Let $x \in \mathbb{F}$ be a cyclically reduced word. Assume that $\operatorname{Wh}_B(x)$ is connected with no cut-vertices. Then x is undistorted in the Aut-norm. Moreover, there exists a homogeneous quasimorphism which is bounded on primitive elements and is non-zero on x.

Proof. Let Br_{x^2} be a counting quasimorphism defined with respect to the basis B. Let $b \in \mathbb{F}$ be a primitive element. By Lemma 3.2 and Remark 3.3 we have that

$$|Br_{x^2}(b)| \le 2|x|_B.$$

Moreover, $|Br_{x^2}(x^n)|$ growths linearly with n, thus by Remark 2.2, x is undistorted in the Aut-norm. In order to obtain a desired quasimorphism, it is enough to consider the homogenization of Br_{x^2} , see Lemma 2.4.

3.3. Separable sets. Let A be a finite subset of \mathbb{F} . We call A separable, if there exist two non-trivial free factors F_1, F_2 such that $\mathbb{F} = F_1 * F_2$, and every element in A can be conjugated into F_1 or F_2 . We always assume that elements in A are cyclically reduced. Note that A is separable if and only if $\phi(A)$ is separable, where $\phi \in \operatorname{Aut}(\mathbb{F})$. The following proposition follows from Proposition 2.2 and Proposition 2.3 in [25].

Proposition 3.5. Let $x \in \mathbb{F}$. If $\operatorname{Wh}_B(x)$ is disconnected, then x is separable. If $\operatorname{Wh}_B(x)$ is connected and has a cut-vertex, then there is $\psi \in \operatorname{Aut}(\mathbb{F})$, such that $|\psi(x)|_B < |x|_B$.

3.4. Distortion of elements in the Aut-norm of free groups.

Theorem 3.6. Let **F** be a free group and let $x \in \mathbf{F}$. Then either:

- (a) x is separable and then the cyclic subgroup generated by x is bounded in the Aut-norm, or
- (b) x is undistorted in the Aut-norm. Moreover, there exists a homogeneous quasimorphism which is bounded on the set of all primitive elements and is non-trivial on x.

Proof. Let $|\cdot|_p$ be the primitive norm, see Subsection 2.1. Suppose that x is separable. It means that $\mathbb{F} = F_1 * F_2$, F_2 is not trivial and x can be conjugated into F_1 . Since $|\cdot|_p$ is invariant under inner automorphisms, we can assume that $x \in F_1$. Let B_1 and B_2 be some bases of F_1 and F_2 respectively. Let $p \in B_2$ and $n \in \mathbb{N}$. Note that the element $x^n p$ is primitive. Indeed, the set $B_1 \cup \{x^n p\} \cup (B_2 \setminus \{p\})$ is a free basis of \mathbb{F} . Thus $|x^n|_p = |x^n pp^{-1}|_p \le 2$.

Now suppose, that x is not separable. Let B be a basis of F. We claim that we can find an element y in the Aut(F)-orbit of x such that $Wh_B(y)$ is connected and has no cut-vertices. Note that $Wh_B(x)$ is connected. Indeed, if it was not connected, then by Proposition 3.5, x would be separable. Let $|\cdot|_B$ be the word norm defined by B. If $Wh_B(x)$ has a cut-vertex, then again by Proposition 3.5 there is an automorphism $\psi \in Aut(F)$ such that $|\psi(x)|_B < |x|_B$.

Now we consider $\operatorname{Wh}_B(\psi(x))$. As before, it is a connected graph. If it has a cutvertex, then we apply Proposition 3.5 again and find a new ψ and further reduce the length of the element. At the end we get an element y with the property that $\operatorname{Wh}_B(y)$ is connected with no cut-vertices. Now consider the cyclical reduction $y' = r_B(y)$.

By definition we have $\operatorname{Wh}_B(y') = \operatorname{Wh}_B(y)$. Lemma 3.4 gives us a homogeneous quasimorphism q which is bounded on primitives and is non-trivial on y'. Since x and y' are in the same Aut-orbit, there exists $\psi \in \operatorname{Aut}(\mathbf{F})$ such that $\psi(x) = y'$. If we define $q'(w) = q(\psi(w))$, then q' is bounded on primitive elements and $q'(x) \neq 0$.

Corollary 3.7. An element x of \mathbf{F} is separable if and only if $\operatorname{Wh}_B(x)$ is not connected or has a cut-vertex for every basis B.

Proof. Assume that x is separable. If there is a basis B such that $Wh_B(x)$ is connected and has no cut-vertices, then by applying Lemma 3.4 for an element $r_B(x)$, we see that x is undistorted in the Aut-norm. Thus x is not separable by Theorem 3.6.

To prove the reversed implication, we apply inductively Proposition 3.5 and obtain an element y in the same $Aut(\mathbf{F})$ -orbit as x, such that $Wh_B(y)$ is not connected. Thus again by Proposition 3.5, an element y, and consequently x, is separable.

Remark 3.8. Corollary 3.7 is not entirely new. It can be deduced from Theorem 2.4 and Proposition 2.3 in [25]. However, we think that our proof of this fact is interesting since it is simpler and shorter than the proof of Stallings.

4. Aut-norm on surface groups

In this section we study distortion in surface groups. Our main tool is the theory of mapping class groups. The principal idea is to embed a surface group in its automorphism group (which is a mapping class group of a surface) via the Birman embedding, and then find appropriate quasimorphisms on this group. In Subsection 4.1 we recall the Nielsen–Thurston normal form of a mapping class. Then in Subsection 4.2 we give the Nielsen–Thurston decomposition of mapping classes which lie in the image of the Birman embedding. Finally, in Subsection 4.3 we use the quasimorphisms defined by Bestvina–Bromberg–Fujiwara to prove Theorem 1 for surface groups.

4.1. Nielsen–Thurston normal form. Let S be a compact oriented surface with finitely many punctures in the interior of S. By MCG(S) we denote the mapping class group of S, that is the group of isotopy classes of orientation preserving homeomorphisms of S. We assume that homeomorphisms and isotopies fix the boundary of S pointwise.

We recall briefly the Nielsen-Thurston normal form of an element in MCG(S). A loop γ in S is called *essential*, if no component of $S \setminus \gamma$ is homeomorphic to a disc, a punctured disc or an annulus. An element $g \in MCG(S)$ is called *reducible* if there exists a non-empty set $C = \{c_1, \ldots, c_n\}$ of isotopy classes of essential simple loops in S such that:

- (1) All elements in C can be represented by pairwise disjoint simple loops.
- (2) The set C is g-invariant.

Such C is called a *reduction system* for g. A reduction system for g is maximal if it is not a proper subset of any other reduction system for g. There may be many maximal reduction systems. However, we can define the unique one by defining the *canonical reduction system* to be the intersection of all maximal reduction systems. Note that the canonical reduction system is not necessary maximal.

Now let us describe the canonical form of an element of a mapping class group. Assume for a moment, that S has no boundary. Let $g \in MCG(S)$ and let $C = \{c_1, \ldots, c_m\}$ be its canonical reduction system. Choose pairwise disjoint representatives of the classes c_i together with pairwise disjoint closed annuli R_1, \ldots, R_m , where R_i is a closed neighborhood of a representative of c_i . Let S_1, \ldots, S_p be the closures of connected components of $S \setminus \bigcup_{i=1}^m R_i$. Then there is a power k and a representative $\psi \in \operatorname{Homeo}^+(S)$ of g^k such that:

- (1) The homeomorphism ψ fixes the subsurfaces, i.e. $\psi(R_i) = R_i$ for $1 \le i \le m$ and $\psi(S_i) = S_i$ for $1 \le i \le p$.
- (2) The restriction of ψ to R_i is a power of a Dehn twist.
- (3) The restriction of ψ to S_i is pseudo-Anosov or the identity.

Thus, up to finite power, any element is described as a commuting product of powers of Dehn twists and pseudo-Anosov homeomorphisms on subsurfaces. If S has a boundary, then we need to add to our collection R_i collar neighborhoods of boundary curves. Then the additional terms which can appear in the decomposition of g^k are powers of Dehn twists along boundary curves. A mapping class which has only one factor in the Nielsen-Thurston decomposition is called *pure*. We have the following characterization of the canonical reduction system.

Proposition 4.1. Assume that the surface S has no boundary and possibly has punctures. The system $C = \{c_1, \ldots, c_s\}$ is the canonical reduction system for g if and only if the following two conditions hold:

- (a) There exists $k \in \mathbb{N}$ and pairwise disjoint loops γ_i representing classes c_i such that g^k restricted to any component of $S \setminus (\gamma_1 \cup \cdots \cup \gamma_s)$ is trivial or pseudo-Anosov.
- (b) The set C is a minimal set with this property.

Dehn twists are not mentioned in Proposition 4.1 since every Dehn twist along some γ_i becomes trivial in the mapping class group of $S \setminus (\gamma_1 \cup \cdots \cup \gamma_s)$.

4.2. Filling curves and the theorem of Kra. Let S be a closed oriented surface of genus g, possibly with punctures, and let $\star \in S$. Assume that S has negative Euler characteristic. We consider the Birman exact sequence:

$$1 \to \pi_1(S, \star) \xrightarrow{\text{Push}} \text{MCG}(S, \star) \xrightarrow{F} \text{MCG}(S) \to 1.$$

By $MCG(S, \star)$ we mean the group of homeomorphisms which fix \star , taken up to isotopies which fix \star at any time. Since fixing a point and removing a point does not make any difference for mapping classes, we have that $MCG(S, \star) = MCG(S \setminus \star)$. The map F is the forgetful map. The map Push is defined as follows: let γ be a based loop which represents an element in $\pi_1(S, \star)$. Let ψ be any homeomorphism which fixes \star , such that: there exists an isotopy $H: [0,1] \times S \to S$ such that $H(0,\cdot) = \mathrm{id}_S$, $H(1,\cdot) = \psi$ and $H(t,\star) = \gamma(t)$. Then $\mathrm{Push}([\gamma]) = [\psi]$. This is a well-defined map. One can imagine that $\mathrm{Push}([\gamma])$ takes \star and pushes it along the loop γ . For a detailed discussion see [13].

The goal of this subsection is to understand the Nielsen–Thurston decomposition of Push($[\gamma]$). In Theorem 4.11 we generalize a theorem of I. Kra [19], for the short proof see [13]. It states that if γ is filling (see Definition 4.6), then Push($[\gamma]$) is pseudo-Anosov.

Example 4.2. Let γ be a simple loop based at \star . We identify a tubular neighborhood of γ with the annulus $S^1 \times [-1, 1]$. Let $\gamma^{\pm} = S^1 \times \{\pm 1\}$. Then $Push([\gamma]) = T_{\gamma^+} T_{\gamma^-}^{-1}$, where $T_{\gamma^{\pm}}$ is the Dehn twist along γ^{\pm} . The surface S is not a torus. Hence γ^- is not homotopic to γ^+ in $S \setminus \{\star\}$, and T_{γ^+} and T_{γ^-} are different elements of $MCG(S, \star)$.

In what follows, a subsurface S' of S always assumed to be closed in S, and the boundary of S' is a union of pairwise disjoint simple loops. A loop is called *primitive* if, as an element of the fundamental group it is not a proper power of any other element. We always assume, that a loop in S is in general position, i.e. it is a smooth immersed loop with only double self-intersections. A loop is in a *minimal position* if it has the minimal number of double points.

Definition 4.3. Assume that S is a closed oriented surface, possibly with punctures. Let $\gamma: \mathbf{S}^1 \to S$ be a loop in a minimal position. We define a *subsurface* S_{γ} in the following way: First we consider a small collar neighborhood

$$N: \mathbf{S}^1 \times [-1, 1] \to S,$$

where N is a smooth immersion and $N(\cdot,0) = \gamma(\cdot)$, such that the image of N retracts onto the image of γ . Then we add to the image of N all the components of $S \setminus \operatorname{im}(N)$ which are disks or punctured disks.

Lemma 4.4. Assume that S is a closed oriented surface, possibly with punctures. Let γ be a loop in a minimal position such that it is not homotopic to a power of a simple loop. Then the boundary components of S_{γ} are essential simple loops in S.

Proof. Assume that one boundary component $\partial_0 S_\gamma$ is not essential in S. Then $\partial_0 S_\gamma$ bounds a disc or a punctured disc D. Thus $S \setminus \partial_0 S_\gamma$ has two connected components: D and $S \setminus D$. Hence γ is contained either in D or in $S \setminus D$. By construction, it is impossible that $\gamma \subset S \setminus D$, because then we would add D to S_γ . Thus $\gamma \subset D$. But in D every loop is homotopic to a power of a simple loop, and we get a contradiction.

Lemma 4.5. Assume that S is a closed oriented surface, possibly with punctures. Let γ be a non-trivial loop in a minimal position such that is not homotopic to a power of a simple loop. Let $\partial_1 S_{\gamma}$ and $\partial_2 S_{\gamma}$ be two homotopic components of ∂S_{γ} . Then they bound an annulus lying outside of the interior of S_{γ} . Moreover, they are not homotopic to any other component of ∂S_{γ} .

Proof. Any two homotopic simple loops bound an annulus in S. Let A be an embedded annulus bounded by $\partial_1 S_{\gamma}$ and $\partial_2 S_{\gamma}$. Since A is a connected component of $S \setminus (\partial_1 S_{\gamma} \cup \partial_2 S_{\gamma})$, the loop γ is either outside A or inside A. The loop γ cannot be inside A, since every loop in A is a power of a simple loop and this is excluded by the assumption.

Now assume that some boundary component $\partial_i S_{\gamma}$ of S_{γ} is homotopic to $\partial_1 S_{\gamma}$. Let $\partial_1' S_{\gamma}$ denote a loop in the interior of A homotopic to $\partial_1 S_{\gamma}$. Let A' be an embedded annulus bounding $\partial_1' S_{\gamma}$ and $\partial_i S_{\gamma}$. If $\partial_i S_{\gamma}$ is not $\partial_1 S_{\gamma}$ or $\partial_2 S_{\gamma}$, then $\partial_1 S_{\gamma}$ or $\partial_2 S_{\gamma}$ are contained in the interior of A'. This is impossible, since we already know that the interior of A' is disjoint from S_{γ} . Thus $\partial_i S_{\gamma}$ equals to $\partial_1 S_{\gamma}$ or $\partial_2 S_{\gamma}$.

In the following definitions and Lemma 4.8, S is an orientable surface, possibly with boundary and punctures.

Definition 4.6. Let γ be a loop on S. We say that γ *fills* S, if every essential simple loop on S has a non-trivial intersection with every curve homotopic to γ .

Definition 4.7. A loop $\gamma: \mathbf{S}^1 \to S$ has an embedded 1-gon if there is a closed arc $\alpha \subset \mathbf{S}^1$, such that γ restricted to the interior of α is an embedding and $\gamma(\partial \alpha)$ is a single point. A loop $\gamma: \mathbf{S}^1 \to S$ has an embedded 2-gon if there are two disjoint closed arcs α , β such that $\gamma(\partial \alpha) = \gamma(\partial \beta)$ and γ restricted to α and β is an embedding.

We need the following lemma.

Lemma 4.8 ([17, Lemma 2.8]). Assume that γ has no embedded 1-gons and 2-gons. Let c be an essential simple loop and assume that c is disjoint from some curve homotopic to γ . Then there is a simple loop c' isotopic to c such that γ is disjoint from c'.

The next two lemmas deal with the surface S_{γ} . Since we defined S_{γ} only for surfaces with no boundary, we assume below that $\partial S = \emptyset$.

Lemma 4.9. Let γ be a loop in a minimal position on a closed orientable surface S, possibly with punctures. Then γ fills S_{γ} .

Proof. By the definition of S_{γ} we see that every connected component of $S_{\gamma} \setminus \gamma$ is a disc or a punctured disc. It follows that every essential simple loop intersects γ non-trivially. If there is an essential simple loop c disjoint from some curve homotopic to γ , then by Lemma 4.8 we get some essential simple loop c' disjoint from γ , which is a contradiction.

Lemma 4.10. Assume that γ is in a minimal position and is not homotopic to a power of a simple loop. Then the Euler characteristic of S_{γ} is negative.

Proof. On annulus, torus, disk, punctured disk or 2-punctured sphere, every loop is homotopic to a power of a simple loop. Thus S_{γ} is none of those. Every other oriented surface has negative Euler characteristic.

Now we prove the main result of this subsection.

Theorem 4.11 (Generalized Kra's theorem). Let S be a closed oriented surface (possibly with punctures) whose Euler characteristic is negative. Let \star be a base point. Then:

- (a) If $x \in \pi_1(S, \star)$ is a power of a simple element represented by γ^n , where γ is a simple essential loop in S, then $\operatorname{Push}(x) = T_{\gamma^+}^n T_{\gamma^-}^{-n}$ is the Nielsen-Thurston decomposition of $\operatorname{Push}(x) \in \operatorname{MCG}(S, \star)$.
- (b) If $x \in \pi_1(S, \star)$ is not a power of a simple element, then the Thurston-Nielsen decomposition of $Push(x) \in MCG(S, \star)$ consists of a single pseudo-Anosov component.

Proof. If x is a power of a simple element, then by Example 4.2,

$$Push(x) = T_{\gamma^+}^n T_{\gamma^-}^{-n}.$$

Since S is not a torus and γ is by assumption essential, γ^+ and γ^- are not homotopic essential disjoint simple loops. If an element of a mapping class group can be represented as a product of powers of Dehn twists along disjoint not homotopic essential simple loops, then this representation is unique. It follows that the set $C = \{\gamma^+, \gamma^-\}$ satisfies the conditions of Proposition 4.1. Indeed, if C is smaller, then Push(x) equals to a power of T_{γ^\pm} , and hence is not equal to $T_{\gamma^+}^n T_{\gamma^-}^{-n}$.

Assume that x is not a power of a simple element. Let γ be a loop in a minimal position which represents x. Then by Lemma 4.10 the Euler characteristic of S_{γ} is negative and by Lemma 4.9 γ fills S_{γ} . By the result of Kra ([19]), Push(x) is pseudo-Anosov on S_{γ} . We have to prove that Push(x), as an element of MCG(S, \star), has the desired decomposition.

Remark. Even though $\operatorname{Push}(x)$ is pseudo-Anosov on S_{γ} , it does not follow automatically that it is pseudo-Anosov on the subsurface $S_{\gamma} \subset S$ regarded as an element of $\operatorname{MCG}(S,\star)$. For example one can easily imagine a homeomorphism ϕ on S which is trivial in $\operatorname{MCG}(S,\star)$ but is pseudo-Anosov on some smaller subsurface containing the support of ϕ . In addition, it also may happen that some boundary loops of S_{γ} are homotopic, and hence the set of boundary loops of S_{γ} cannot be taken as the canonical reduction system.

Let C be a set of connected components of ∂S_{γ} . By Lemma 4.5 and Lemma 4.4, we can write

$$C = \{\partial_1 S_{\gamma}, \partial_1' S_{\gamma}, \dots, \partial_t S_{\gamma}, \partial_t' S_{\gamma}, \partial_{t+1} S_{\gamma}, \partial_{t+2} S_{\gamma}, \dots, \partial_u S_{\gamma}\},\$$

such that for each 0 < i < t + 1 loops $\partial_i S_{\gamma}$ and $\partial'_i S_{\gamma}$ are homotopic, and the set $C^0 = \{\partial_1 S_{\gamma}, \dots, \partial_u S_{\gamma}\}$ consists of pairwise non-homotopic loops.

We prove that C^0 is the canonical reduction system for $\operatorname{Push}(x)$. To do that, it is enough to check the first and the second conditions of Proposition 4.1. Let S'_{γ} be the connected component of $S \setminus C^0$ which contains γ . The surface S'_{γ} is just S_{γ} with annuli attached to some boundary components. Thus if $\operatorname{Push}(x)$ is pseudo-Anosov on S_{γ} , it is pseudo-Anosov on S'_{γ} . On any other component of $S \setminus C^0$, $\operatorname{Push}(x)$ is trivial.

Let us check the second condition. Let $C' \subset C^0$, $\partial_i S_\gamma \not\in C'$ and S' a connected component of $S \setminus C'$ which contains γ . Since $\partial_i S_\gamma \subset S'$, $\operatorname{Push}(x)$ fixes $\partial_t S_\gamma$. All components of the boundary of S' are in C^0 , thus $\partial_t S_\gamma$ is not homotopic to any component of the boundary of S'. Thus $\partial_t S_\gamma$ is essential. Since $\operatorname{Push}(x)$ is nontrivial and fixes an essential curve, it is reducible. It follows, that C' does not satisfy the second condition. Hence C^0 is the canonical reduction system. Thus the decomposition of $\operatorname{Push}(x)$ consists of a single pseudo-Anosov diffeomorphism on S'_γ .

4.3. Distortion of elements in the Aut-norm of surface groups.

Theorem 4.12. Let $x \in \Gamma_g$. Then either:

- (a) x is a power of a simple non-separating loop, then the cyclic subgroup generated by x is bounded in the Aut-norm, or
- (b) x is undistorted in the Aut-norm. Moreover, there exists a homogeneous quasimorphism bounded on the set of all simple elements and is non-trivial on x.

We start with transferring the problem from finding a suitable quasimorphism on Γ_g to finding a quasimorphism on (a finite index subgroup) of MCG(S). Let us consider the general case first. Let G be a group and let Aut(G) be the group of automorphisms of G. Let $c_g(x) = gxg^{-1}$ be the inner automorphism induced by g. Define homomorphism $c: G \to Aut(G)$ by $c(g) = c_g$.

Lemma 4.13. Let G be an H-finitely generated group for some $H < \operatorname{Aut}(G)$, and let $H_{\circ} < H$ be a finite index subgroup such that $\operatorname{im}(c) < H_{\circ}$. Let $x \in G$ and assume that there exists a homogeneous quasimorphism $q: H_{\circ} \to \mathbf{R}$ such that $q(c_x) \neq 0$. Then x is undistorted in $|\cdot|_H$.

Proof. Let \overline{S} be an H-finite generating set of G. Then \overline{S} is H_{\circ} -finite since $\#(\overline{S}/H_{\circ}) = \#(H/H_{\circ})\#(\overline{S}/H)$. Thus \overline{S} can be used to define both norms $|\cdot|_H$

and $|\cdot|_{H_o}$. To prove that x is undistorted in $|\cdot|_H$ it is enough to prove that it is undistorted in $|\cdot|_{H_o}$.

Let \tilde{q} be the pull back of q to G, i.e. $\tilde{q}(w) = q(c_w)$ for $w \in G$. Now we will show, that \tilde{q} is H_{\circ} -invariant. Let $w \in G$ and $\psi \in H_{\circ}$. Note that $\psi c_w \psi^{-1} = c_{\psi(w)}$ and that q is constant on conjugacy classes of H_{\circ} . We have

$$\tilde{q}(\psi(w)) = q(c_{\psi(w)}) = q(\psi c_w \psi^{-1}) = q(c_w) = \tilde{q}(w).$$

We apply Corollary 2.3 and finish the proof.

We will use this lemma in the case of $G = \Gamma_g$ and $H = \operatorname{Aut}(\Gamma_g)$. We need to take a finite index subgroup H_o , because in most cases there is no homogeneous quasimorphism on the whole group $\operatorname{Aut}(\Gamma_g)$ which is non-zero on a given element c_x . We will be able to find such quasimorphism on some H_o and conclude undistortedness of x in $|\cdot|_{\operatorname{Aut}(\Gamma_g)}$.

Proof of Theorem 4.12. Case 1. Let α and γ be the elements shown in the Figure 3. We first prove that the cyclic group generated by α is bounded in $|\cdot|_{\operatorname{Aut}(\Gamma_g)}$. Let $|\cdot|_s$ be the simple loops norm. It is a simple observation, that the loop $\alpha^n \gamma$ is simple for each $n \in \mathbb{Z}$. Thus we have

$$|\alpha^n|_s = |\alpha^n \gamma \gamma^{-1}|_s \le 2.$$

Since $|\cdot|_s$ defines bi-Lipschitz equivalence class $|\cdot|_{\operatorname{Aut}(\Gamma_g)}$, the cyclic subgroup $\langle \alpha \rangle$ is bounded in the Aut-norm.




Figure 3. Loops α and γ .

Now assume that $x \in \Gamma_g$ is a power of a simple non-separating loop. Note that every simple non-separating loop can be mapped to α . Indeed, for every two simple non-separating loops β_1 , β_2 , the surfaces $S \setminus \beta_i$ are homeomorphic, thus all simple non-separating loops are in one $\operatorname{Aut}(\Gamma_g)$ -orbit. It follows that x can be mapped to α^m for some $m \in \mathbb{N}$. Thus the subgroup generated by x is bounded.

Case 2. Assume that $x \in \Gamma_g$ is not a power of a simple non-separating loop. Let S be a closed surface without punctures and let $\star \in \Gamma_g$. Consider the natural map

$$\Psi: MCG(S, \star) \to Aut(\Gamma_g)$$

induced by the action of a homeomorphism on the fundamental group of (S, \star) . Let $\operatorname{Aut}^+(\Gamma_g) := \operatorname{im}(\Psi)$, then $\operatorname{Aut}^+(\Gamma_g)$ has index 2 in $\operatorname{Aut}(\Gamma_g)$. By the Baer–Dehn–Nielsen theorem, Ψ is an embedding.

It is easy to see (at least for simple elements, see Example 4.2) that $\operatorname{Push}(w)$ induces a conjugation on Γ_g by w. Hence instead of working with $\operatorname{Aut}^+(\Gamma_g)$ and homomorphism c, we work with $\operatorname{MCG}(S, \star)$ and Push .

By the work of Bestvina–Bromberg–Fujiwara ([4]) we know that there are plenty of quasimorphisms on mapping class groups. We describe their result in the way that is convenient for us. The group $MCG(S, \star)$ acts on $H_1(S, \mathbb{Z}/3\mathbb{Z})$. Let H_\circ be the subgroup of $MCG(S, \star)$ which contains all elements that act trivially. Hence H_\circ is a finite index subgroup of $MCG(S, \star)$. Since conjugation acts trivially on homology, $Push(\Gamma_g) < H_\circ$. It follows from [4, Corollary 5.3 and Corollary 5.5] that there exists a homogeneous quasimorphism on H_\circ which is non-zero on Push(x) if one of the following holds:

- (a) In the Nielsen–Thurston decomposition of Push(x) there is at least one pseudo-Anosov element.
- (b) In the Nielsen–Thurston decomposition of Push(x) there is at least one non-trivial power of a Dehn twist along some separating curve.

Now we finish the proof. If $x \in \Gamma_g$ is not a power of non-separating simple loop, then by Theorem 4.11 either:

- (a) *x* is not simple, and the Nielsen–Thurston decomposition contains exactly one pseudo-Anosov element, or
- (b) x is a power of a Dehn twist along separating loop, and then the Nielsen–Thurston decomposition is

$$Push(x) = T_{\gamma^+}^n T_{\gamma^-}^n$$

for some separating simple loop γ and $n \in \mathbb{N}$.

In both cases we have a homogeneous quasimorphism

$$q: H_{\circ} \to \mathbf{R}$$

which is non-trivial on Push(x). Since H_o is a finite index subgroup of MCG(S, \star) it can be viewed as finite index subgroup of Aut(Γ_g). Using Lemma 4.13 we conclude the proof of the theorem.

Let S_n be the set of all elements of Γ_g which are represented by curves with crossing-number at most n (see [9, Definition 1.1]) and let S'_n be the set of all primitive elements in S_n . In [9], D. Calegari proved that $x \in \Gamma_g$ is undistorted in $|\cdot|_{S_n}$ if it has a non-zero self-intersection number. He asked (see Question 1.6) whether simple elements are undistorted with respect to the metrics $|\cdot|_{S'_n}$.

First of all, we note that all these metrics are bi-Lipschitz equivalent. Indeed, by [9, Remark 1.4] all sets S'_n are Aut(Γ_g)-finite. The proof is analogous to the proof of

the fact that S_0 , which is the set of all simple elements, is $\operatorname{Aut}(\Gamma_g)$ -finite. It means that all the metrics $|\cdot|_{S'_n}$ define the same bi-Lipschitz equivalence class $|\cdot|_{\operatorname{Aut}(\Gamma_g)}$. Thus the part of Theorem 4.12 that concerns simple elements gives a complete answer to the question of D. Calegari, i.e. we proved the following:

Corollary 4.14. Simple separating elements in Γ_g are undistorted with respect to the $|\cdot|_{S'_n}$ -norm for every n, and simple non-separating elements generate bounded cyclic subgroup.

4.4. More applications and remarks.

Theorem 4.15 (Many fellows property). Let S be a closed hyperbolic surface of genus g and l be a closed simple non-separating geodesic. Then for every $p \in l$ and every neighborhood U of p, there is another simple closed geodesic l' passing through U.

Proof. Assume on the contrary that there exists $p \in l$ and some neighborhood U of p such that every closed geodesic different from l does not pass through U. Let Γ_g be the fundamental group of S. For every $x \in \Gamma_g$ there is a unique closed geodesic l_x in the free homotopy class of x. Two geodesics l_x and $l_{x'}$ are equal if and only if x and x' are conjugated. Moreover, simple elements of Γ_g correspond to simple closed geodesics. Let $l = l_{x_0}$. Let α be a 1-differential form supported on U such that $\int_I \alpha \neq 0$. In [3] Barge–Ghys showed that the following function:

$$q(x) = \int_{l_x} \alpha,$$

is a homogeneous quasimorphism. If x is a simple element not conjugated to x_o , then $l_x \neq l_{x_o}$ and by our assumption l_x does not pass through U. Hence $q(l_x) = 0$. It follows that the only simple elements on which q is non-zero are conjugates of x_0 . Since q is constant on conjugacy classes, it is bounded on the set of all simple elements. By Lemma 2.1 we get that x_0 is undistorted in the simple loops norm which contradicts Theorem 4.12

Remark 4.16. Theorem 4.15 holds for separating closed geodesics as well, see [23, Lemma 5.1]. Note that unlike the set of all closed geodesics, the set of simple closed geodesics is not dense in S and, as suggested by Theorem 4.15, there are some preferred tracks chosen by simple geodesics. For a further discussion of this phenomenon see [8].

Remark 4.17. Let x be a simple non-separating element and let γ be a loop that represents x. Then there is no finite index subgroup of $MCG(S, \star)$ containing Push(x) such that $T_{\gamma+}$ is not conjugated to $T_{\gamma-}$. Indeed, otherwise one could find a quasimorphism on $MCG(S, \star)$ which is non-trivial on Push(x). Using Lemma 4.13 we see that x would be undistorted in the simple loops norm which contradicts Theorem 4.12.

5. $Aut(F_2)$ -invariant quasimorphisms on F_2

Let S_g be a closed surface of genus g. Let $\circ, \star \in S_g$ be two arbitrary points. We shall regard \circ as a puncture and \star as a base point. Let us consider the group $MCG(S_g \setminus \{\circ\}, \star)$ of mapping classes of $S_g \setminus \{\circ\}$ fixing the point \star .

The natural action of a homeomorphism on $\mathbb{F}_{2g} = \pi_1(S_g \setminus \{\circ\}, \star)$ induces a map

$$\pi: MCG(S_g \setminus \{\circ\}, \star) \to Aut(\mathbb{F}_{2g}).$$

We claim that this map is injective. Indeed, the surface $S_g \setminus \{\circ\}$ can be described as a regular 4g-gon with opposite edges identified, such that the point \circ lies in the center and \star is one of the vertices. If $\pi(\psi)$ is the identity for some $\psi \in \text{MCG}(S_g \setminus \{\circ\}, \star)$, then there exists a representation of ψ fixing pointwise the edges of the 4g-gon. Thus we can regard this representation of ψ as a homeomorphism of the punctured disc fixing the boundary. By the Alexander trick ([13, Lemma 2.1]) such an element is isotopic to the identity.

In the next theorem we consider non-simple element $x \in \mathbb{F}_{2g}$ such that for every $\psi \in \mathrm{MCG}(S_g \setminus \{\circ\}, \star)$ we have $x^{-1} \neq \psi(x)$. We postpone the proof of the existence of such elements to the next section (see Lemma 6.5).

Theorem 5.1. Let $x \in \mathbb{F}_{2g}$ such that for every $\psi \in MCG(S_g \setminus \{\circ\}, \star)$ we have $x^{-1} \neq \psi(x)$, and x cannot be represented by a simple loop in $S_g \setminus \{\circ\}$. Then there exists a non-trivial $MCG(S_g \setminus \{\circ\}, \star)$ -invariant homogeneous quasimorphism on \mathbb{F}_{2g} which is non-zero on x.

Proof. We consider the Birman embedding $\mathbb{F}_{2g} \xrightarrow{\operatorname{Push}} \operatorname{MCG}(S_g \setminus \{\circ\}, \star)$. Since x is not simple, it follows from Theorem 4.11 that the Nielsen–Thurston decomposition of $\operatorname{Push}(x)$ consists of one non-trivial pseudo-Anosov pure component. In addition, $\operatorname{Push}(x)$ is not conjugated to its inverse in the group $\operatorname{MCG}(S_g \setminus \{\circ\}, \star)$. Indeed, if there is an element $\psi \in \operatorname{MCG}(S_g \setminus \{\circ\}, \star)$ which conjugates $\operatorname{Push}(x)$ to its inverse, then it implies

$$Push(\psi(x)) = \psi Push(x)\psi^{-1} = Push(x)^{-1} = Push(x^{-1}).$$

Since Push is injective, we conclude that $\psi(x) = x^{-1}$ which contradicts our assumption.

Now we use quasimorphisms constructed in [4]. Note that if an element in a group is conjugated to its inverse, then every homogeneous quasimorphism vanishes on this element. If follows from [4, Theorem 4.2] that for pure elements of $MCG(S_g \setminus \{\circ\}, \star)$, being conjugated to its inverse is the only obstruction to be detected by homogeneous quasimorphisms. Due to Lemma 4.13, Push(x) is pure. Thus there exists a homogeneous quasimorphism q on $MCG(S_g \setminus \{\circ\}, \star)$ which is non-zero on Push(x). The pull-back of q to F_{2g} by Push gives us an $MCG(S_g \setminus \{\circ\}, \star)$ -invariant quasimorphism which does not vanish on x.

Remark 5.2. Let $n \geq 3$. One would like to construct $\operatorname{Aut}(\mathbb{F}_n)$ -invariant quasimorphism on \mathbb{F}_n by restricting a quasimorphism from $\operatorname{Aut}(\mathbb{F}_n)$ to \mathbb{F}_n which is embedded in $\operatorname{Aut}(\mathbb{F}_n)$ via inner automorphisms. However, despite an extensive study of $\operatorname{Aut}(\mathbb{F}_n)$ it is not known if there are quasimorphisms on $\operatorname{Aut}(\mathbb{F}_n)$ which restrict non-trivially to \mathbb{F}_n .

Theorem 5.3. The linear space of $MCG(S_g \setminus \{\circ\}, \star)$ -invariant homogeneous quasi-morphisms on \mathbb{F}_{2g} is infinite dimensional.

Proof. It follows from Lemma 6.6 and Lemma 6.5 that there is an infinite sequence z_1, z_2, \ldots of integers such that the elements $x_k = a^{z_k} b^{2z_k} a^{3z_k} b^{4z_k}$ have the following properties:

- (a) x_k and x_k^{-1} belong to different Aut(\mathbb{F}_{2g})-orbits,
- (b) for $i, j \in \mathbb{Z} \setminus \{0\}$ and $k \neq k'$, elements x_k^i and $x_{k'}^j$ belong to different Aut(\mathbb{F}_{2g})-orbits.

Elements x_k are not simple. Indeed, every element $x \in \mathbb{F}_{2g}$ that can be represented by a simple loop in $MCG(S_g \setminus \{\circ\}, \star)$ is a primitive element of \mathbb{F}_{2g} , and hence is inverted by some automorphism of \mathbb{F}_{2g} .

Recall that by Theorem 4.11, $\operatorname{Push}(x_k)$ is a pure mapping class, whose Nielsen–Thurston decomposition consists only of one pseudo-Anosov component. Moreover, using an argument from the proof of Theorem 5.1, property (a) implies that the element $\operatorname{Push}(x_k)$ is not conjugated to its inverse in the mapping class group, and property (b) implies that for $k \neq k'$ and any non-zero i and j, $\operatorname{Push}(x_k)^i$ is not conjugated to $\operatorname{Push}(x_{k'})^j$. In the language of chiral and achiral classes introduced in [4], it means that the elements $\operatorname{Push}(x_k)$ represent different chiral classes for different k. Let $k \in \mathbb{N}$. It follows from [4, Proposition 4.4] that each function

$$\{\operatorname{Push}(x_1), \dots, \operatorname{Push}(x_k)\} \to \mathbf{R}$$

is a restriction of a homogeneous quasimorphism on $MCG(S_g \setminus \{\circ\}, \star)$. If we pullback these quasimorphisms to \mathbb{F}_{2g} by Push, we obtain that each function

$$\{x_1,\ldots,x_k\}\to\mathbf{R}$$

is a restriction of some $MCG(S_g \setminus \{\circ\}, \star)$ -invariant homogeneous quasimorphism. Consequently, for each $k \in \mathbb{N}$ we constructed a k-dimensional subspace of $MCG(S_g \setminus \{\circ\}, \star)$ -invariant homogeneous quasimorphisms.

For g > 1, $\pi(MCG(S_g \setminus \{\circ\}, \star))$ has infinite index in $Aut(\mathbb{F}_{2g})$. The situation is different for g = 1. For completeness we give a proof of the following lemma.

Lemma 5.4. $\pi(MCG(S_1\setminus\{\circ\},\star))$ has index 2 in Aut(\mathbb{F}_2).

Proof. Let us extend the group $MCG(S_1 \setminus \{\circ\}, \star)$ to $MCG^{\pm}(S_1 \setminus \{\circ\}, \star)$ by allowing orientation reversing homeomorphisms. Then π naturally extends to the map $\pi': MCG^{\pm}(S_1 \setminus \{\circ\}, \star) \to Aut(\mathbb{F}_2)$. By the same argument as in the beginning of this chapter, π' is injective. Let $\{a, b\}$ be a basis of \mathbb{F}_2 . The group $Aut(\mathbb{F}_2)$ is generated by the following automorphisms:

$$a \to a^{-1}$$
 $a \to b$ $a \to ab$
 $b \to b$ $b \to a$ $b \to b$.

Each one of them can be realized by an element of $MCG^{\pm}(S_1 \setminus \{\circ\}, \star)$. Thus π' is onto.

Every automorphism of \mathbf{F}_n acts on its abelianisation which is isomorphic to \mathbf{Z}^n , thus defines a matrix over \mathbf{Z} . Let $\mathrm{Aut}^+(\mathbf{F}_n)$ be the subgroup consisting of all elements which define matrices of determinant 1. In the case of n=2, we have $\mathrm{Aut}^+(\mathbf{F}_2)=\pi(\mathrm{MCG}(S_1\setminus\{\circ\},\star))$. An immediate consequence of Theorem 5.3 is that the linear space of homogeneous $\mathrm{Aut}^+(\mathbf{F}_2)$ -invariant quasimorphisms is infinite dimensional.

Remark 5.5. The fact that the space of homogeneous $\operatorname{Aut}^+(\mathbb{F}_2)$ -invariant quasimorphisms on \mathbb{F}_2 is non-trivial was recently proved in his thesis by Huber in [18]. He showed that certain rotation number quasimorphism is $\operatorname{Aut}^+(\mathbb{F}_2)$ -invariant.

In the next corollary we improve this result to $\text{Aut}(\mathbb{F}_2)$ -invariant homogeneous quasimorphisms.

Corollary 5.6. The linear space of homogeneous $Aut(\mathbf{F}_2)$ -invariant quasimorphisms on \mathbf{F}_2 is infinite dimensional.

Proof. Let a and b be generators of \mathbb{F}_2 . Denote by σ the automorphism defined by $\sigma(a) = a^{-1}, \sigma(b) = b$. Let $\{z_i\}_{i=1}^{\infty}$ be a sequence of integers such that $\gcd(z_i) \neq \gcd(z_j)$ for $i \neq j$. Let $x_k = a^{z_k}b^{2z_k}a^{3z_k}b^{4z_k}$. Consider the set

$$X:=\{x_1,\sigma(x_1),x_2,\sigma(x_2),\ldots\}\subset \mathbf{F}_2.$$

It follows from Lemma 6.5 that no x_k is inverted by an automorphism of \mathbf{F}_2 . The same applies to elements $\sigma(x_k)$. It means that each Push $(\sigma(x_k))$ is chiral.

For every $x, y \in X$, the elements Push(x) and Push(y) define different chiral classes in $MCG(S_1 \setminus \{\circ\}, \star)$, see Lemma 6.9. It follows from [4, Proposition 4.4] that each function

$$\{\operatorname{Push}(x_1),\operatorname{Push}(\sigma(x_1)),\ldots,\operatorname{Push}(x_k),\operatorname{Push}(\sigma(x_k))\}\to\mathbf{R}$$

is a restriction of a homogeneous quasimorphism on $MCG(S_1 \setminus \{\circ\}, \star)$. By pulling back to \mathbb{F}_2 , we obtain that every function

$$\{x_1,\sigma(x_1),\ldots,x_k,\sigma(x_k)\}\to \mathbf{R}$$

is a restriction of a homogeneous $\operatorname{Aut}^+(\mathbb{F}_2)$ -invariant quasimorphism.

Let $Q(\mathbf{F}_2)^{\mathrm{Aut}^+}$ be the space of homogeneous $\mathrm{Aut}^+(\mathbf{F}_2)$ -invariant quasimorphisms on \mathbf{F}_2 . Let $X_k = \{x_1, \sigma(x_1), \dots, x_k, \sigma(x_k)\}$. Denote by \mathbf{R}^{X_k} the set of all functions from X_k to \mathbf{R} . We have the following commutative diagram:

$$Q(\mathbf{F}_{2})^{\operatorname{Aut}^{+}} \xrightarrow{\operatorname{Sym}_{\sigma}} Q(\mathbf{F}_{2})^{\operatorname{Aut}^{+}}$$

$$\downarrow \qquad \qquad \downarrow \qquad$$

where

$$\operatorname{Sym}_{\sigma}(q)(x) = q(x) + q(\sigma(x)),$$

for $q \in \overline{Q}(\mathbb{F}_2)^{\mathrm{Aut}^+}$. We claim that $\mathrm{Sym}_{\sigma}(q)$ is an $\mathrm{Aut}(\mathbb{F}_2)$ -invariant homogeneous quasimorphism. It is clear that $\mathrm{Sym}_{\sigma}(q)$ is a quasimorphism, because q is a quasimorphism and σ is an automorphism of \mathbb{F}_2 . To prove the $\mathrm{Aut}(\mathbb{F}_2)$ invariance, we first note that $\mathrm{Sym}_{\sigma}(q)$ is σ -invariant, which is obvious from the definition. Moreover, for every $\psi \in \mathrm{Aut}^+(\mathbb{F}_2)$ we can find $\psi' \in \mathrm{Aut}^+(\mathbb{F}_2)$ such that $\psi'\sigma = \sigma\psi$. Now

$$\operatorname{Sym}_{\sigma}(q)(\psi(x)) = q(\psi(x)) + q(\sigma\psi(x)) = q(x) + q(\psi'\sigma(x))$$
$$= q(x) + q(\sigma(x)) = \operatorname{Sym}_{\sigma}(q)(x).$$

Thus the quasimorphism $\operatorname{Sym}_{\sigma}(q)$ is σ -invariant and $\operatorname{Aut}^+(\mathbb{F}_2)$ -invariant, and consequently $\operatorname{Aut}(\mathbb{F}_2)$ -invariant.

The map $\widetilde{\operatorname{Sym}}_{\sigma}$ is defined by $\widetilde{\operatorname{Sym}}_{\sigma}(f)(x) = f(x) + f(\sigma(x))$. The vertical epimorphisms in the above diagram are restrictions. We have that

$$\operatorname{Im}(\widetilde{\operatorname{Sym}}_{\sigma}) = \{ f \in \mathbb{R}^{X_k} : f(x_l) = f(\sigma(x_l)) \text{ for each } l \in \{1, \dots, k\} \}$$

is a k-dimensional linear space. Each element of $\operatorname{Im}(\widetilde{\operatorname{Sym}}_{\sigma})$ is a restriction of $\operatorname{Sym}_{\sigma}(q)$ for some $q \in Q(\mathbb{F}_2)^{\operatorname{Aut}^+}$, which is an $\operatorname{Aut}(\mathbb{F}_2)$ -invariant homogeneous quasimorphism. Thus for each $k \in \mathbb{N}$ the space of $\operatorname{Aut}(\mathbb{F}_2)$ -invariant homogeneous quasimorphisms on \mathbb{F}_2 contains a k-dimensional subspace.

Remark 5.7. In his thesis Hase [15, 16] proved that the space of quasimorphisms on \mathbb{F}_2 that are not $\operatorname{Aut}(\mathbb{F}_2)$ -invariant is dense in the space of all homogeneous quasimorphisms on \mathbb{F}_2 . In particular, finite linear combinations of counting quasimorphisms are not $\operatorname{Aut}(\mathbb{F}_2)$ -invariant. Hence a rotation number quasimorphism considered by Huber in [18] is not $\operatorname{Aut}(\mathbb{F}_2)$ -invariant, since it is a linear combination of counting quasimorphisms.

It follows from Corollary 5.6, that there exists an infinite dimensional space of quasimorphisms on \mathbb{F}_2 where each quasimorphism can not be expressed as a finite linear combination of counting quasimorphisms. On the other hand, Grigorchuk [14] showed that every quasimorphism is a linear combination of (possibly infinitely many) counting quasimorphisms.

6. Whitehead algorithm and some elements of F_n

J. H. C. Whitehead [27] described an algorithm that, given two elements $x, x' \in \mathbb{F}_n$, it decides if there is $\psi \in \operatorname{Aut}(\mathbb{F}_n)$ for which $\psi(x) = x'$. Below we recall the Whitehead algorithm.

Definition 6.1. Let X be a basis of \mathbb{F}_n . An element $\psi \in \operatorname{Aut}(\mathbb{F}_n)$ is called:

- (a) *Permutation automorphism* if ψ permutes the set $X \cup X^{-1}$.
- (b) Whitehead automorphism if there is an element $a \in X \cup X^{-1}$ such that $\psi(a) = a$ and $\psi(x) \in \{x, ax, xa^{-1}, axa^{-1}\}$ for each $x \in X \setminus \{a\}$.

Let Ω_n denote the set of all permutation automorphisms. The set Ω_n is a finite subgroup of $\operatorname{Aut}(\mathbb{F}_n)$ which is isomorphic to the extended permutation group.

Theorem 6.2 (Whitehead). Let $x \in \mathbb{F}_n$ and let $m = \min\{|\psi(x)|_X\}$ where the minimum is taken over all $\psi \in \operatorname{Aut}(\mathbb{F}_n)$. If $|x|_X > m$, then there exists a Whitehead automorphism h such that $|h(x)|_X < |x|_X$. If x and x' are in the same Aut-orbit and $|x|_X = |x'|_X = m$, then there exists a sequence of permutation and Whitehead automorphisms t_1, \ldots, t_l such that:

and
$$t_l \dots t_1(x) = x'$$

 $|x|_X = |t_1(x)|_X = |t_2t_1(x)|_X = \dots = |t_l \dots t_1(x)|_X = m.$

Let $x \in \mathbb{F}_n$. We denote by \overline{x} the conjugacy class represented by x. If c is any conjugacy class, we define its length by

$$|c|_X := \min\{|x|_X, \ \overline{x} = c\}.$$

Note that $Aut(\mathbf{F}_n)$ acts on conjugacy classes of \mathbf{F}_n . It is easy to see, that the analogous version of Whitehead algorithm works for conjugacy classes and the norm defined above.

In the following lemmas we consider a sequence $\{x_k\}_{k=2}^{\infty}$ where $x_k \in \mathbb{F}_n$ is of the form

$$x_k = a^k b^{2k} a^{3k} b^{4k}$$
.

Here the elements a and b denote two different generators of \mathbb{F}_n .

Lemma 6.3. Let n = 2 and $X = \{a, b\}$. Let t be a permutation automorphism. Then $t(\overline{x_k}) = \overline{x_k}$ if and only if t is the identity, i.e. the group Ω_2 acts freely on the orbit $\Omega_2(\overline{x_k})$.

Proof. The conjugacy class $t(\overline{x_k})$ is represented by the element of the form $u^k v^{2k} u^{3k} v^{4k}$ for some $u, v \in \{a, b, a^{-1}, b^{-1}\}$. This element represents conjugacy class of $a^k b^{2k} a^{3k} b^{4k}$ if and only if u = a and v = b. It means that t(a) = a and t(b) = b. Thus t is the identity.

Lemma 6.4. Let ψ be a Whitehead automorphism and let $\overline{x} \in \Omega_n(x_k)$. Then either $\psi(\overline{x}) = \overline{x}$ or $|\psi(\overline{x})|_X > |\overline{x}|$.

Proof. The element \overline{x} equals to $\phi(x_k)$ for some $\phi \in \Omega_n$. Thus \overline{x} is represented by the element $u^k v^{2k} u^{3k} v^{4k}$, where $u, v \in X \cup X^{-1}$ and $u \notin \{v, v^{-1}\}$. Consider a Whitehead automorphism ψ . Assume that a from the definition of the Whitehead automorphism is not equal to u, v, u^{-1}, v^{-1} . If $\psi(u) = aua^{-1}$ and $\psi(v) = ava^{-1}$, then $\psi(\overline{x}) = \overline{x}$. In all other cases we have $|\psi(\overline{x})|_x > |\overline{x}|_x$, since in $\psi(\overline{x})$ there always will be some occurrences of the letter a.

If $a \in \{u, v, u^{-1}, v^{-1}\}$, then up to inner automorphisms, there are only 5 different ways a Whitehead automorphism can act on $\{u, v\}$. They are listed below:

$$u \to u$$
 $u \to uv^{\pm}$ $u \to u$
 $v \to v$ $v \to vu^{\pm}$.

Direct computation shows that these automorphisms, except the one which fixes u and v, increase the length of \overline{x} (provided that k > 1). Thus ψ does not increase the length of \overline{x} if and only if ψ fixes \overline{x} .

Lemma 6.5. Elements x_k and x_k^{-1} belong to different $Aut(\mathbf{F}_n)$ -orbits.

Proof. It follows from Theorem 6.2 that $\Omega_n(\overline{x}_k)$ is the set of all conjugacy classes minimizing the norm $|\cdot|_X$ in the $\operatorname{Aut}(\mathbb{F}_n)$ -orbit of \overline{x}_k . Indeed, if some $\overline{y} \in F_n$ minimizes the norm, then there exist permutation or Whitehead automorphisms t_1, \ldots, t_l such that

$$|\overline{x}_k|_X = |t_1(\overline{x}_k)|_X = \dots = |t_l \dots t_1(\overline{x}_k)|_X$$

and $t_1 ldots t_1(\overline{x}_k) = \overline{y}$. Since $|\overline{x}_k|_X = |t_1(\overline{x}_k)|_X$, we conclude that either t_1 is a permutation automorphism, or t_1 is a Whitehead automorphism and by Lemma 6.4 we have $t_1(\overline{x}_k) = \overline{x}_k$. Then we apply the same argument to the element $t_1(\overline{x}_k)$ and the equality

$$|t_2(t_1(\overline{x}_k))|_X = |t_1(\overline{x}_k)|_X$$

to conclude that t_2 is a permutation automorphism or $t_2(t_1(\overline{x}_k)) = t_1(\overline{x}_k)$. It follows that each t_i is a permutation automorphism, or fixes the element $t_{i-1} \dots t_1(\overline{x}_k)$. Hence $\overline{y} \in \Omega_n(\overline{x}_k)$.

If \overline{x}_k^{-1} is in the same $\operatorname{Aut}(\mathbf{F}_n)$ -orbit as \overline{x}_k , then \overline{x}_k^{-1} would minimize the norm. Thus to prove the lemma, it remains to prove that \overline{x}_k^{-1} does not belong to $\Omega_n(\overline{x}_k)$. Note that if $t(\overline{x}_k) = \overline{x}_k^{-1}$ for $t \in \Omega_n$, then necessarily $t(a), t(b) \in \{a, a^{-1}, b, b^{-1}\}$. It is easy to check that for such automorphisms we always have $t(\overline{x}_k) \neq \overline{x}_k^{-1}$.

Lemma 6.6. Let $i, j \in \mathbb{Z} \setminus \{0\}$ and let $k, l \geq 2$ such that $gcd(k) \neq gcd(l)$. Elements $(x_k)^i$ and $(x_l)^j$ belong to different $Aut(\mathbb{F}_n)$ -orbits.

Proof. For every element $x \in \mathbb{F}_n$ there is a unique (up to a sign) number n and a unique (up to taking an inverse) element p which is not a proper power of any other element such that $x = p^n = (p^{-1})^{-n}$. Assume that there exists an automorphism $\psi \in \operatorname{Aut}(\mathbb{F}_n)$ such that $\psi((x_k)^i) = (x_l)^j$. Both x_k and x_l are not proper powers, hence $\psi(x_k) = (x_l)^{\pm 1}$. In what follows we show that this is impossible.

Every automorphism $\psi\colon \mathbb{F}_n o \mathbb{F}_n$ induces the abelianisation automorphism

$$Ab(\psi): \mathbb{Z}^n \to \mathbb{Z}^n$$
.

Elements x_k and $(x_l)^{\pm}$ are mapped to vectors with coordinates equal to 4k, 6k and $\pm 4l$, $\pm 6l$ in the abelianisation. It is enough to show, that these vectors belong to different $\operatorname{Aut}(\mathbf{Z}^n)$ -orbits. Indeed this is the case, since automorphisms of \mathbf{Z}^n preserve the greatest common divisor of coordinates of a vector, and

$$\gcd(4k, 6k) = 2\gcd(k) \neq 2\gcd(l) = \gcd(\pm 4l, \pm 6l).$$

Lemma 6.7. Let n = 2 and $X = \{a, b\}$. Let $\sigma \in \operatorname{Aut}(\mathbb{F}_2)$ be defined by $\sigma(a) = a^{-1}$, $\sigma(b) = b$. Then x_k and $\sigma(x_k)$ belong to different $\operatorname{Aut}^+(\mathbb{F}_2)$ -orbits.

Proof. Let us recall that the group $\operatorname{Aut}^+(\mathbb{F}_2)$ consists of automorphisms ψ for which $\det(\operatorname{Ab}(\psi)) = 1$. We have that

$$\sigma(x_k) = a^{-k} b^{2k} a^{-3k} b^{4k}$$

and $\sigma \in \operatorname{Aut}(\mathbb{F}_2) \setminus \operatorname{Aut}^+(\mathbb{F}_2)$. Suppose that $\sigma(x_k) = \psi(x_k)$ for some automorphism $\psi \in \operatorname{Aut}^+(\mathbb{F}_2)$. It means that $\psi^{-1}\sigma \in \operatorname{Stab}(x_k)$ and $\psi^{-1}\sigma \in \operatorname{Aut}(\mathbb{F}_2) \setminus \operatorname{Aut}^+(\mathbb{F}_2)$. In what follows we show that $\operatorname{Stab}(x_k) < \operatorname{Aut}^+(\mathbb{F}_2)$ which is a contradiction.

Let us consider the stabilizer $\operatorname{Stab}(\overline{x}_k)$ of the conjugacy class \overline{x}_k . Of course $\operatorname{Stab}(x_k) < \operatorname{Stab}(\overline{x}_k)$. We show that $\operatorname{Stab}(\overline{x}_k) < \operatorname{Aut}^+(\mathbf{F}_2)$. We use the construction presented in [20] in order to find a generating set of $\operatorname{Stab}(\overline{x}_k)$. First we define a graph Δ as follows: a vertex of Δ is a conjugacy class of minimal length in the $\operatorname{Aut}(\mathbf{F}_2)$ -orbit of \overline{x}_k . It follows from the proof of Lemma 6.5 that this set equals to $\Omega_2(\overline{x}_k)$. Two vertices $v_1, v_2 \in \Omega_2(\overline{x}_k)$ are connected by a directed edge from v_1 to v_2 if there is a permutation or Whitehead automorphism ψ such that $\psi(v_1) = v_2$. We will consider edge embedded loops in Δ based at \overline{x}_k . The theorem of McCool [20] says, that $\operatorname{Stab}(\overline{x}_k)$ is generated by elements which are products of labels read from all possible loops like this.

It follows from Lemma 6.3 that the subgraph spanned by edges labeled with permutation automorphisms is a complete graph on the set $\Omega_2(\bar{x}_k)$, with no loops. Lemma 6.4 implies that all edges labeled by Whitehead automorphisms are loops.

Let δ be a loop in Δ based at \overline{x}_k . Now we show that the word w which is read from the labels of δ is trivial in $\operatorname{Aut}(\mathbf{F}_2)/\operatorname{Aut}^+(\mathbf{F}_2)\cong \mathbf{Z}/2\mathbf{Z}$. Note that all Whitehead automorphisms belong to $\operatorname{Aut}^+(\mathbf{F}_2)$. Thus we can ignore labels coming from the edges labeled by Whitehead automorphisms. Since the edges labeled by Whitehead

automorphisms are loops, we can assume that δ goes through the edges labeled only by permutation automorphisms. Thus we can assume, that the element read from the labels of δ is just a permutation automorphism which fixes \overline{x}_k . By Lemma 6.3 this element is trivial.

Remark 6.8. More detailed analysis shows that $Stab(x_k)$ is a cyclic subgroup generated by conjugation by x_k . One also can prove the analog of Lemma 6.7 for \mathbf{F}_n by replacing x_k with more complicated elements.

Lemma 6.9. Let $i, j \in \mathbb{Z} \setminus \{0\}$ and let $k, l \geq 2$ such that $\gcd(k \neq \gcd(l))$. Then:

- (a) x_k^i and $\sigma(x_k)^j$ belong to different $\operatorname{Aut}^+(\mathbb{F}_n)$ -orbits.
- (b) $\sigma(x_k)^i$ and $\sigma(x_l)^j$ belong to different Aut(\mathbb{F}_n)-orbits.
- (c) x_k^i and $\sigma(x_l)^j$ belong to different $Aut(\mathbf{F}_n)$ -orbits.

Proof. It follows from the proof of Lemma 6.6 that we can assume i=1 and $j=\pm 1$. Lemma 6.7 implies (a) for j=1. To prove (a) for j=-1 it is enough to note that $\sigma(x_k)^{-1}$ and x_k^{-1} are in the same $\operatorname{Aut}(\mathbb{F}_n)$ -orbit. If x_k and $\sigma(x_k)^{-1}$ are in the same $\operatorname{Aut}^+(\mathbb{F}_n)$ -orbit, then x_k and x_k^{-1} are in the same $\operatorname{Aut}(\mathbb{F}_n)$ -orbit, which contradicts Lemma 6.5. The proof of (b) and (c) is analogous to the proof of Lemma 6.6.

References

- [1] M. Abért, Some questions. Available at: http://www.renyi.hu/~abert/questions.pdf
- [2] V. Bardakov, V. Shpilrain, and V. Tolstykh, On the palindromic and primitive widths of a free group, *J. Algebra*, **285** (2005), no. 2, 574–585. Zbl 1085.20011 MR 2125453
- [3] J. Barge and É. Ghys, Surfaces et cohomologie bornée, *Invent. Math.*, **92** (1988), no. 3, 509–526. Zbl 0641.55015 MR 939473
- [4] M. Bestvina, K. Bromberg, and K. Fujiwara, Stable commutator length on mapping class groups, *Ann. Inst. Fourier (Grenoble)*, **66** (2016), no. 3, 871–898. Zbl 1397.20050 MR 3494163
- [5] M. Brandenbursky, Ś. R. Gal, J. Kędra, and M. Marcinkowski, The cancellation norm and the geometry of bi-invariant word metrics, *Glasg. Math. J.*, 58 (2016), no. 1, 153–176. Zbl 1376.20045 MR 3426433
- [6] M. Brandenbursky and J. Kędra, On the autonomous metric on the group of areapreserving diffeomorphisms of the 2-disc, *Algebr. Geom. Topol.*, 13 (2013), no. 2, 795–816. Zbl 1267.57038 MR 3044593
- [7] R. Brooks, Some remarks on bounded cohomology, in *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, 53–63, Ann. of Math. Stud., 97, Princeton Univ. Press, Princeton, N.J., 1981. Zbl 0457.55002 MR 624804

- [8] P. Buser and H. Parlier, The distribution of simple closed geodesics on a Riemann surface, in *Complex analysis and its applications*, 3–10, OCAMI Stud., 2, Osaka Munic. Univ. Press, Osaka, 2007. Zbl 1151.30031 MR 2405696
- [9] D. Calegari, Word length in surface groups with characteristic generating sets, *Proc. Amer. Math. Soc.*, 136 (2008), no. 7, 2631–2637. Zbl 1146.57001 MR 2390536
- [10] D. Calegari, *scl*, MSJ Memoirs, 20, Mathematical Society of Japan, Tokyo, 2009. Zbl 1187.20035 MR 2527432
- [11] V. Erlandson, A remark on the word length in surface groups, *Trans. Amer. Math. Soc.*, **372** (2019), no. 1, 441–455 Zbl 07076396 MR 3968775
- [12] V. Erlandsson, H. Parlier, and J. Souto, Counting curves, and the stable length of currents, to appear in *J. Amer. Math. Soc.*
- [13] B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, 49, Princeton University Press, 2012. Zbl 1245.57002 MR 2850125
- [14] R. I. Grigorchuk, Some results on bounded cohomology, in *Combinatorial and geometric group theory (Edinburgh, 1993)*, 111–163, London Math. Soc. Lecture Note Ser., 204, Cambridge Univ. Press, Cambridge, 1995. Zbl 0853.20034 MR 1320279
- [15] A. Hase, Dynamics of $Out(F_n)$ on the second bounded cohomology of F_n . arxiv:1805.00366
- [16] A. Hase, The study of $Out(F_n)$ on the second bounded cohomology $H_b^2(F_n, R)$, Master's thesis, Technion, 2016.
- [17] J. Hass and P. Scott, Intersections of curves on surfaces, *Israel J. Math.*, **51** (1985), no. 1-2, 90–120. Zbl 0576.57009 MR 804478
- [18] T. Huber, Rotation quasimorphisms for surfaces, PhD thesis, ETH Z"urich, 2012.
- [19] I. Kra, On the Nielsen-Thurston-Bers type of some self-maps of Riemann surfaces, *Acta Math.*, **146** (1981), no. 3-4, 231–270. Zbl 0477.32024 MR 611385
- [20] J. McCool, Some finitely presented subgroups of the automorphism group of a free group, J. Algebra, 35 (1975), 205–213. Zbl 0325.20025 MR 396764
- [21] A. D. Miasnikov and A. G. Myasnikov, Whitehead method and genetic algorithms, in Computational and experimental group theory, 89–114, Contemp. Math., 349, Amer. Math. Soc., Providence, RI, 2004. Zbl 1076.20013 MR 2077761
- [22] R. P. Osborne and H. Zieschang, Primitives in the free group on two generators, *Invent. Math.*, **63** (1981), no. 1, 17–24. Zbl 0438.20017 MR 608526
- [23] H. Parlier, Hyperbolic polygons and simple closed geodesics, *Enseign. Math.* (2), **52** (2006), no. 3-4, 295–317. Zbl 1114.30044 MR 2300612
- [24] A. H. Rhemtulla, A problem of bounded expressibility in free products, *Proc. Cambridge Philos. Soc.*, **64** (1968), 573–584. Zbl 0159.03001 MR 225889
- [25] J. R. Stallings, Whitehead graphs on handlebodies, in *Geometric group theory down under (Canberra, 1996)*, 317–330, de Gruyter, Berlin, 1999. Zbl 1127.57300 MR 1714852
- [26] J. H. C. Whitehead, On Certain Sets of Elements in a Free Group, *Proc. London Math. Soc.* (2), **41** (1936), no. 1, 48–56. Zbl 0013.24801 MR 1575455
- [27] J. H. C. Whitehead, On equivalent sets of elements in a free group, *Ann. of Math.* (2), **37** (1936), no. 4, 782–800. Zbl 0015.24804 MR 1503309

Received 19 February, 2017

M. Brandenbursky, Department of Mathematics, Ben-Gurion University of the Negev, PO Box 653, Be'er Sheva 8410501, Israel

E-mail: brandens@math.bgu.ac.il

M. Marcinkowski, Ben-Gurion University; Regensburg Universität; and Instytut Matematyczny, Uniwersytet Wrocławski, pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland

E-mail: marcinkow@math.uni.wroc.pl

