

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 94 (2019)
Heft: 4

Artikel: On rational cuspidal plane curves and the local cohomology of Jacobian rings
Autor: Dimca, Alexandru
DOI: <https://doi.org/10.5169/seals-847028>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On rational cuspidal plane curves and the local cohomology of Jacobian rings

Alexandru Dimca*

Abstract. This note gives the complete projective classification of rational, cuspidal plane curves of degree at least 6, and having only weighted homogeneous singularities. It also sheds new light on some previous characterizations of free and nearly free curves in terms of Tjurina numbers. Finally, we suggest a stronger form of Terao’s conjecture on the freeness of a line arrangement being determined by its combinatorics.

Mathematics Subject Classification (2010). 14H50; 14B05, 13D02, 32S22.

Keywords. Jacobian ideal, Tjurina number, free curve, nearly free curve.

1. Introduction

The main result we prove in this note is the following.

Theorem 1.1. *Let C be a rational, cuspidal plane curve of degree $d \geq 6$ such that C has only weighted homogeneous singularities. Then C is projectively equivalent to exactly one of the following $\phi(d)/2$ models*

$$C_{d,k}: y^d + x^k z^{d-k} = 0,$$

where the integer k satisfies $1 \leq k < d/2$ and k is relatively prime to d . Here ϕ denotes the Euler function, with $\phi(d)$ counting the number of integers m , where $1 \leq m \leq d-1$ and m relatively prime to d .

The normal form of the curves above is in fact just the homogenization of the normal form of a simply-connected, irreducible affine plane curve X : by the Lin–Zaidenberg theorem, see [17, 24], any such curve X is given in suitable global coordinates (u, v) on \mathbb{C}^2 by an equation $u^d + v^k = 0$, for some coprime integers $d > k \geq 1$.

*This work has been supported by the French government, through the UCA^{JEDI} Investments in the Future project managed by the National Research Agency (ANR) with the reference number ANR-15-IDEX-01.

A slightly stronger result than Theorem 1.1 is stated and proved below, see Propositions 4.1 and 4.3. For detailed information on the rational, cuspidal plane curves of degree ≤ 5 , we refer to [15, 16] and the references given there. The curves $C_{d,k}$ above are called *binomial cuspidal curves* in [15, Section 7.1]. The necessary restriction $d \geq 6$ in Theorem 1.1 is discussed in Example 4.4 below.

Theorem 1.1 is quite surprising, given that the classification of rational, cuspidal plane curves, even of those having at most two singularities, is rather complicated; see the Sakai–Tono paper [19], or Propositions 3.1 and 3.2 in [10] where their results are quoted. It is known that there are no rational, cuspidal *free* curves of degree $d \geq 6$ having only weighted homogeneous singularities, see [10, Theorem 2.8]. Theorem 1.1 says that if the word *free* is deleted from the above statement, only the binomial cuspidal curves may occur, and they are all nearly free. Note also that a weighted homogeneous cusp has only one Puiseux pair, but the class of plane curve singularities with exactly one Puiseux pair is much larger. This explains the complicated classification of rational unicuspidal curves with a unique Puiseux pair in [14].

To prove Theorem 1.1, we use a number of key results, proved by A. A. du Plessis and C. T. C. Wall in [12, 13], the main one being stated below in Theorem 3.1. To make our note more self-contained, we include a short proof of a similar result to Theorem 3.1, which we explain now.

Let $S = \mathbb{C}[x, y, z]$ be the graded polynomial ring in three variables x, y, z and let $C: f = 0$ be a reduced curve of degree d in the complex projective plane \mathbb{P}^2 . The minimal degree of a Jacobian relation, or Jacobian syzygy, for the polynomial f is the integer $mdr(f)$ defined to be the smallest integer $m \geq 0$ such that there is a nontrivial relation

$$af_x + bf_y + cf_z = 0 \quad (1.1)$$

among the partial derivatives f_x, f_y and f_z of f with coefficients a, b, c in S_m , the vector space of homogeneous polynomials of degree m . When $mdr(f) = 0$, then C is a union of lines passing through one point, a situation easy to analyse. We assume from now on that

$$mdr(f) \geq 1.$$

Denote by $\tau(C)$ the global Tjurina number of the curve C , which is the sum of the Tjurina numbers of the singular points of C . We denote by J_f the Jacobian ideal of f , i.e. the homogeneous ideal in S spanned by f_x, f_y, f_z , and by $M(f) = S/J_f$ the corresponding graded ring, called the Jacobian (or Milnor) algebra of f . Let I_f denote the saturation of the ideal J_f with respect to the maximal ideal $\mathfrak{m} = (x, y, z)$ in S and consider the local cohomology group

$$N(f) = I_f/J_f = H_{\mathfrak{m}}^0(M(f)).$$

It was shown in [7] that the graded S -module $N(f)$ satisfies a Lefschetz type property with respect to multiplication by generic linear forms. This implies in

particular the inequalities

$$0 \leq n(f)_0 \leq n(f)_1 \leq \cdots \leq n(f)_{[T/2]} \geq n(f)_{[T/2]+1} \geq \cdots \geq n(f)_T \geq 0,$$

where $T = 3d - 6$ and $n(f)_k = \dim N(f)_k$ for any integer k . We set

$$v(C) = \max_j \{n(f)_j\}.$$

When $d = 2m$ is even, then the above implies that $n(f)_{3m-3} = v(C)$. When $d = 2m + 1$ is odd, then the above and the self duality of the graded S -module $N(f)$, see [21, 23], implies that

$$n(f)_{3m-2} = n(f)_{3m-1} = v(C).$$

The second main result of this note is the following.

Theorem 1.2. *Let $C: f = 0$ be a reduced plane curve of degree d and let $r = mdr(f)$. Then the following hold.*

(1) *If $r < d/2$, then*

$$v(C) = (d - 1)^2 - r(d - 1 - r) - \tau(C).$$

(2) *If $r \geq (d - 2)/2$, then*

$$v(C) = \left\lceil \frac{3}{4}(d - 1)^2 \right\rceil - \tau(C).$$

Here, for any real number u , $\lceil u \rceil$ denotes the round up of u , namely the smallest integer U such that $U \geq u$. Written down explicitly, this means that for $d = 2m$ is even and $r \geq m - 1$, one has $v(C) = 3m^2 - 3m + 1 - \tau(C)$, while for $d = 2m + 1$ is odd and $r \geq m$, one has $v(C) = 3m^2 - \tau(C)$. For $(d - 2)/2 \leq r < d/2$, both formulas in (1) and (2) apply, and they give the same result for $v(C)$. The relation between Theorem 1.2 and du Plessis-Wall result in Theorem 3.1 is discussed in Remark 3.2 below.

When $C: f = 0$ is a line arrangement, examples due to G. Ziegler show that the invariant $mdr(f)$ is not combinatorially determined, see [1, Example 4.3]. The above result suggests that the following stronger version of H. Terao's conjecture, saying that the freeness of a line arrangement is combinatorially determined, might be true. For more on Terao's conjecture and free hyperplane arrangements we refer to [4].

Conjecture 1.3. *Let $C: f = 0$ be a line arrangement in \mathbb{P}^2 . Then the invariant $v(C)$ is combinatorially determined.*

For a reduced plane curve one may state the following.

Conjecture 1.4. *Let $C: f = 0$ be a reduced plane curve \mathbb{P}^2 , of degree $d \geq 6$. Then the invariant $\nu(C)$ is determined by the degree of C and the list of the analytic types of the isolated singularities of C . Or this version: similar claim, but with the additional restriction that all the singularities of C are weighted homogeneous, and we use now the list of topological type of singularities.*

Note that it is the second version in Conjecture 1.4 which generalizes Conjecture 1.3. Some cases where these conjectures hold are described in Example 4.4 (i) (where the corresponding invariant $mdr(f)$ is not determined by the data stated in Conjecture 1.4), and in Proposition 3.5. The need of the assumption $d \geq 6$ in Conjecture 1.4 is explained by looking at a pair of quintic curves with two cusps of type A_4 , and respectively E_8 , in Example 4.4 (iii). This pair of quintic curves occurs already in [2, Remark 3.10], as a possible source of counter-examples to a generalized Terao's conjecture.

The examples of conic-line arrangements \mathcal{A} and \mathcal{A}' in [20], having the same combinatorics but different values for the invariant $\nu(C)$, do not give counter-examples to Conjecture 1.4, since, as implied by [20, Concluding remarks (2)], the list of the analytic type of singularities in the two conic-line arrangements \mathcal{A} and \mathcal{A}' are not the same, and some of the singularities are not weighted homogeneous.

Acknowledgements. We thank the referee for his many valuable suggestions, in particular for drawing our attention on the two quintic curves described in Example 4.4 (iii) below.

2. The proof of Theorem 1.2

Consider the graded S -submodule $AR(f) \subset S^3$ of *all relations* involving the partial derivatives of f , namely

$$\rho = (a, b, c) \in AR(f)_m$$

if and only if $af_x + bf_y + cf_z = 0$ and a, b, c are in S_m . We set $ar(f)_k = \dim AR(f)_k$ for any integer k .

Consider the rank two vector bundle $T\langle C \rangle = \text{Der}(-\log C)$ of logarithmic vector fields along C , which is the coherent sheaf associated to the graded S -module $AR(f)(1)$. Here we use the *shift notation*: for a graded S -module M and any integer k , the shifted graded S -module $M(k)$ satisfies $M(k)_j = M_{k+j}$ for any j . Using the results in the third section of [8], for any integer k one has

$$\chi(T\langle C \rangle(k)) = 3 \binom{k+3}{2} - \binom{d+k+2}{2} + \tau(C). \quad (2.1)$$

Moreover, one has the following for $E = T\langle C \rangle$ and any integer k , see [8, 21].

$$\begin{aligned} h^0((E(k))) &= ar(f)_{k+1}, \\ h^1((E(k))) &= n(f)_{d+k}, \quad \text{and} \quad h^2((E(k))) = ar(f)_{d-5-k}. \end{aligned} \quad (2.2)$$

Assume that we are in the case $d = 2m$ and apply the formulas (2.1) and (2.2) for $k = m - 3$. We get

$$2ar(f)_{m-2} - v(C) = \tau(C) - 3m^2 + 3m - 1.$$

Let $r = mdr(f)$ and note that $ar(f)_{m-2} = 0$ if $r \geq m - 1 = [(d - 1)/2]$ and

$$ar(f)_{m-2} = \dim S_{m-2-r} = \binom{m-r}{2} \text{ if one has } r \leq m - 2.$$

Indeed, it follows from [22, Lemma 1.4] that the S -module $AR(f)$ cannot have two independent elements of degree d_1 and d_2 satisfying $d_1 + d_2 < d - 1$. Assume next that we are in the case $d = 2m + 1$ and apply the formulas (2.1) and (2.2) again for $k = m - 3$. We get

$$ar(f)_{m-2} - v(C) + ar(f)_{m-1} = \tau(C) - 3m^2.$$

As above, if $r = mdr(f)$, we have $ar(f)_{m-2} = ar(f)_{m-1} = 0$ if $r \geq m = [(d - 1)/2]$ and

$$ar(f)_{m-2} = \dim S_{m-2-r} = \binom{m-r}{2}$$

$$\text{and} \quad ar(f)_{m-1} = \dim S_{m-1-r} = \binom{m-r+1}{2}$$

if $r \leq m - 1$, with the convention $\binom{1}{2} = 0$. These formulas, plus some simple computation, prove Theorem 1.2 stated in the introduction.

Remark 2.1. An alternative proof of Theorem 1.2 can be obtained using the results in [1]. Indeed, Proposition 3.2 in [1] implies that $d_1^{L_0} = r$ for $r < (d - 2)/2$ and $d_1^{L_0} = \lfloor (d - 1)/2 \rfloor$ for $r \geq (d - 2)/2$. Here $(d_1^{L_0}, d_2^{L_0})$, with $d_1^{L_0} \leq d_2^{L_0}$, denotes the splitting type of the vector bundle $T\langle C \rangle(-1)$ along a generic line L_0 in \mathbb{P}^2 . Then [1, Theorem 1.1] says that

$$(d - 1)^2 - d_1^{L_0} d_2^{L_0} = \tau(C) + v(C).$$

Since $d_1^{L_0} + d_2^{L_0} = d - 1$ by [1, Proposition 3.1], these facts give a new proof of Theorem 1.2.

In view of this remark, Conjecture 1.3 may be restated as follows.

Conjecture 2.2. *Let $C: f = 0$ be a line arrangement in \mathbb{P}^2 . Then the generic splitting type $(d_1^{L_0}, d_2^{L_0})$ of the vector bundle $T\langle C \rangle(-1)$ is combinatorially determined.*

This conjecture, in the form of a question, was made for the first time in [3, Question 7.12]. Note that we can restate Conjecture 1.4 in a similar way, see Example 4.4 (iii) below for an interesting pair of quintic curves.

3. Some related results and direct applications

We start this section by recalling the following result due to du Plessis and Wall, see [12, Theorem 3.2].

Theorem 3.1. *For positive integers d and r , define two new integers by*

$$\tau(d, r)_{\min} = (d - 1)(d - r - 1) \quad \text{and} \quad \tau(d, r)_{\max} = (d - 1)^2 - r(d - r - 1).$$

Then, if $C: f = 0$ is a reduced curve of degree d in \mathbb{P}^2 and $r = mdr(f)$, one has

$$\tau(d, r)_{\min} \leq \tau(C) \leq \tau(d, r)_{\max}.$$

Moreover, for $r = mdr(f) \geq d/2$, the stronger inequality

$$\tau(C) \leq \tau(d, r)_{\max} - \binom{2r + 2 - d}{2}$$

holds.

Remark 3.2. Let $C: f = 0$ be a reduced curve of degree d in \mathbb{P}^2 and $r = mdr(f)$. Note that the function $\tau(d, r)_{\max}$, regarded as a function of r , occurs also in Theorem 1.2 (1), which can be restated as

$$\tau(C) + \nu(C) = \tau(d, r)_{\max}, \tag{3.1}$$

for $r = mdr(f) < d/2$. The inequality $\tau(C) \leq \tau(d, r)_{\max}$ in Theorem 3.1 is made more precise, when $r < d/2$, by the result in Theorem 1.2 (1).

On the other hand, note that $\tau(d, r)_{\max}$ is a decreasing (resp. increasing) function of r on the interval $[0, (d - 1)/2]$ (resp. on the interval $[(d - 1)/2, +\infty)$), and the number $\lceil \frac{3}{4}(d - 1)^2 \rceil$ that occurs in Theorem 1.2 (2) is precisely the minimum of the values of the function $\tau(d, r)_{\max}$ for $r \in [0, (d - 1)/2] \cap \mathbb{N}$. Hence Theorem 1.2 (2) implies only

$$\tau(C) \leq \left\lceil \frac{3}{4}(d - 1)^2 \right\rceil,$$

for $r \geq d/2$, which is weaker than the inequality in Theorem 3.1, though enough for many applications.

At the end of the proof of Theorem 3.1, in [12], the authors state the following very interesting consequence (of the proof, not of the statement) of Theorem 3.1.

Corollary 3.3. *Let $C: f = 0$ be a reduced curve of degree d in \mathbb{P}^2 and $r = mdr(f)$. One has*

$$\tau(C) = \tau(d, r)_{\max}$$

if and only if $C: f = 0$ is a free curve, and then $r < d/2$.

Since a plane curve C is free if and only if $\nu(C) = 0$, this characterization of free curves follows also from Theorem 1.2, as explained in Remark 3.2.

In the paper [5], we have given an alternative proof of Corollary 3.3 and have shown that a plane curve C is nearly free, which can be defined by the property $\nu(C) = 1$, if and only if a similar property holds. Namely, one has the following result, an obvious consequence of Theorem 1.2.

Proposition 3.4. *Let $C: f = 0$ be a reduced curve of degree d in \mathbb{P}^2 and $r = mdr(f)$. One has*

$$\tau(C) = \tau(d, r)_{\max} - 1$$

if and only if $C: f = 0$ is a nearly free curve, and then $r \leq d/2$.

Concerning Conjecture 1.3 in Introduction we have the following.

Proposition 3.5. *Let C be a plane curve of degree d such that, for $d = 2m$ (resp. $d = 2m + 1$) one has*

$$\tau(C) < \tau(d, m - 2)_{\min} = (m + 1)(2m - 1)$$

(resp. $\tau(C) < \tau(d, m - 1)_{\min} = 2m(m + 1)$). Then

$$\nu(C) = \left\lceil \frac{3}{4}(d - 1)^2 \right\rceil - \tau(C).$$

In particular, if C be a line arrangement (resp. a reduced plane curve) satisfying these inequalities, then Conjecture 1.3 (resp. Conjecture 1.4) holds for C .

An example of quintic curves showing that the upper bounds on $\tau(C)$ given above are sharp is given in Example 4.4 (iii).

Proof. We give the proof only in the case $d = 2m$ even. The function $\tau(d, r)_{\min}$ is a strictly decreasing function of r on \mathbb{R} , so Theorem 3.1 implies that $r = mdr(f) \geq m - 1$. But then Theorem 1.2 (2) implies that $\nu(C)$ is determined by d and $\tau(C)$. When C is a line arrangement, d and $\tau(C)$ are both combinatorially determined. For a plane curve in general, $\tau(C)$ is determined by the analytic type of the local singularities, but not by their topological type. \square

Example 3.6. Let $C = \mathcal{A}: f = 0$ be a line arrangement consisting of d lines. For $d = 6$, there are 10 possibilities for the intersection lattice $L(\mathcal{A})$, see for details [6, Section 5.6]. Out of them, 8 satisfy the inequality $\tau(\mathcal{A}) < 20$ from Proposition 3.5. For $d = 6$, $r = mdr(f)$ is determined by the intersection lattice $L(\mathcal{A})$, and there are exactly 8 lattices with $r \geq 2$.

For $d = 7$, there are 23 possibilities for the intersection lattice $L(\mathcal{A})$. Out of them, only 4 satisfy the inequality $\tau(\mathcal{A}) < 24$ from Proposition 3.5. For $d = 7$, $r = mdr(f)$ is again determined by the intersection lattice $L(\mathcal{A})$, and there are exactly 19 lattices with $r \geq 3$, for which $\nu(C) = \nu(\mathcal{A})$ is given by the formula in Proposition 3.5.

It would be interesting to find a lower bound for $\tau(\mathcal{A})$ in terms of d and r , in the case of line arrangements, which is better than the bound $\tau(r)_{\min}$ given by Theorem 3.1.

4. The proof of Theorem 1.1

The following result is related to the main conjecture in [11], namely that a rational, cuspidal plane curve is either free or nearly free. It is also the first step in proving Theorem 1.1.

Recall that a plane curve C has only weighted homogeneous singularities if and only if $\mu(C) = \tau(C)$, where $\mu(C)$ denotes the sum of all the Milnor numbers of the singularities of C , see [18]. In general one has the obvious inequality $\tau(C) \leq \mu(C)$.

Proposition 4.1. *Let $C: f = 0$ be an irreducible curve in \mathbb{P}^2 , of degree $d \geq 6$. Then the following properties are equivalent.*

- (1) $\tau(C) \geq d^2 - 4d + 8$.
- (2) $r = mdr(f) = 1$.
- (3) C is a rational, cuspidal plane curve such that $\mu(C) = \tau(C)$, i.e. C has only weighted homogeneous singularities.

If any of these properties hold, then C is a nearly free curve.

Note that the condition (1) implies $d \geq 6$. As the proof below shows, the implication (2) \Rightarrow (3) hold for any $d \geq 2$, and the inequality $d \geq 6$ is essential for the implication (3) \Rightarrow (1).

Proof. To prove that (1) implies (2), recall that the function $\tau(d, r)_{\max}$ is a decreasing function of r on the interval $[1, (d-1)/2]$ and note that $\tau(d, r)_{\max} = \tau(d, r')_{\max}$ for $r + r' = d - 1$. For $r = 2$ we get $\tau(d, 2)_{\max} = d^2 - 4d + 7$. Then (1) implies that $r = mdr(f)$ has to be 1. Indeed, the value $r = d - 2$ is excluded using the stronger final inequality in Theorem 3.1. Note that Theorem 1.2 (2) is also enough for this purpose, as discussed in Remark 3.2. To prove that (2) implies (3), note that

$$\tau(d, 1)_{\min} = d^2 - 3d + 2 \leq \tau(C) \leq \tau(d, 1)_{\max} = d^2 - 3d + 3.$$

The genus $g(C)$ is given by

$$2g(C) = (d-1)(d-2) - \sum_p (\mu(C, p) + r(C, p) - 1),$$

where p runs through the singular points of C , $\mu(C, p)$ is the Milnor number of the singularity (C, p) and $r(C, p)$ is the number of its branches. It follows that

$$\mu(C) = \sum_p \mu(C, p) \leq (d-1)(d-2),$$

with equality if and only if C is a rational cuspidal curve. Then the inequality $\tau(C) \leq \mu(C) \leq (d-1)(d-2)$ forces the equality

$$\tau(d, 1)_{\min} = d^2 - 3d + 2 = \tau(C)$$

and we see that C is nearly free using Proposition 3.4. The fact that (3) implies (1) is obvious as soon as $d \geq 6$, since then

$$\tau(C) = \mu(C) = (d-1)(d-2) \geq d^2 - 4d + 8. \quad \square$$

Remark 4.2. Irreducible, cuspidal nearly free curves satisfying $\mu(C) = \tau(C)$, but not rational have been constructed in [2]. For instance, for any odd integer $k \geq 1$, it is shown that the irreducible curve

$$C_{2k}: f = x^{2k} + y^{2k} + z^{2k} - 2(x^k y^k + x^k z^k + y^k z^k) = 0,$$

has $3k$ singular points of type A_{k-1} as singularities, it is a nearly free curve with $mdr(f) = k$ and has genus

$$g(C_{2k}) = \frac{(k-1)(k-2)}{2}.$$

For $k = 3$, we have in this case $\tau(C) = 18 < \tau(d, 2)_{\max} + 1 = 20$.

The following result completes the proof of Theorem 1.1.

Proposition 4.3. *Let $C: f = 0$ be an irreducible plane curve of degree $d \geq 6$ such that $mdr(f) = 1$. Then C is projectively equivalent to exactly one of the following $\phi(d)/2$ models*

$$C_{d,k}: y^d + x^k z^{d-k} = 0,$$

where the integer k satisfies $1 \leq k < d/2$ and k is relatively prime to d .

Proof. Using Proposition 4.1 and [13, Proposition 1.1], we see that C has a 1-dimensional symmetry, i.e. it admits a 1-dimensional algebraic subgroup H' of $PGL_2(\mathbb{C})$ as automorphism group. This group lifts uniquely to an algebraic 1-parameter subgroup H of $GL_3(\mathbb{C})$, not contained in the center and preserving not

only the curve C , but also its defining equation f . Such a subgroup H may be either semi-simple or nilpotent, and [13, Proposition 3.1] tells us that only the semi-simple case can occur for our situation. Indeed, as in the proof above, we know that

$$\tau(C) = (d-1)(d-2) = d^2 - 3d + 2.$$

Note that the statement of [13, Proposition 3.1] should be slightly corrected, namely the part “if y^d has non-zero coefficient in f ” is to be replaced by “if y^d has zero coefficient in f .” Once we know that H is semi-simple, we can assume that $H = \mathbb{C}^*$ acting on \mathbb{P}^2 via

$$t \cdot [x : y : z] = [t^{w_1}x : t^{w_2}y : t^{w_3}z],$$

for some integers w_j . Then using the discussion and the notation on page 120 in [13], we see that the only possibilities to get irreducible curves correspond to line segments $A\alpha_k$ for $1 \leq k < d/2$ and k relatively prime to d . \square

Example 4.4. (i) Consider the following two rational cuspidal quartics

$$C: f = y^4 - xz^3 = 0 \text{ and } C': f' = y^4 - xz^3 - y^3z.$$

Then both curves have an E_6 -singularity located at $[1 : 0 : 0]$, hence they have only weighted homogeneous singularities. However, as noticed in [15, Section 7.1], in the final part on semi-binomial curves, C and C' are not projectively equivalent. This follows there from the fact that C (resp. C') has one (resp. two) inflection points. From our point of view, the difference between C and C' is that $mdr(f) = 1$, while $mdr(f') = 2$, as a direct computation shows. Note that both curves C and C' are nearly free, since one has

$$\tau(C) = \tau(C') = 6 = \tau(4, 1)_{\max} - 1 = \tau(4, 2)_{\max} - 1.$$

Similar semi-binomial rational cuspidal curves $C: f = 0$ of arbitrary degree d , with a unique Puiseux pair and $\tau(C) < \mu(C)$, and with all the possible values for $r = mdr(f) \in [2, d/2]$, are discussed in [9].

(ii) Up to projective equivalence, all rational cuspidal quintic curves were described and found by Namba in [16, Theorem 2.3.10, pp. 179–182]. In particular, any rational cuspidal quintic curve with 4 cusps is projectively equivalent to

$$C: f = 16x^4y + 128x^2y^2z - 4x^3z^2 + 256y^3z^2 - 144xyz^3 + 27z^5 = 0,$$

see also the discussion of the curve $C_8 = [(2_3), (2), (2), (2)]$ in [15, Section 6.3]. This curve has three A_2 cusps and one A_6 cusp, hence only weighted homogeneous singularities. A direct computation shows that $mdr(f) = 2$ and that $\tau(C) = \tau(5, 2)_{\max}$, and therefore C is a free curve.

(iii) There are essentially two quintic curves with two cusps, one cusp of type A_4 and the other of type E_8 . The first curve, denoted by C_{3A} in [15, Section 6.3], has the parametrization $(s^5 : s^3t^2 : t^5)$, and hence it is just the curve $C_{5,2}$ from Theorem 1.1, if we forget the restriction $d \geq 6$ there. In particular, the curve C_{3A} is nearly free, and hence $\nu(C_{3A}) = 1$. The second curve, denoted by C_{3B} in [15, Section 6.3], has the parametrization $(s^5 : s^3t^2 : st^4 + t^5)$, and hence it can be given by the equation

$$f_{3B} = y^5 - xy^4 + 2x^2y^2z - x^3z^2 = 0.$$

It follows that this curve is free, and hence $\nu(C_{3B}) = 0$. Note that both curves have $\tau(C) = 12$, and hence the inequality in Proposition 3.5, which for $d = 5$ is $\tau(C) < 12$, fails just by a unit. This shows that the bounds given on $\tau(C)$ in Proposition 3.5 are sharp. This pair of quintic curves has been considered also in [2, Remark 3.10]. Using the above computations and Remark 2.1, we see that the generic splitting type of the vector bundle $T\langle C_{3A} \rangle(-1)$ (resp. $T\langle C_{3B} \rangle(-1)$) is $(d_1^{L_0}, d_2^{L_0}) = (1, 3)$ (resp. $(d_1^{L_0}, d_2^{L_0}) = (2, 2)$).

It is surprising that the condition $d \geq 6$ in our results above avoids complicated situations as described in this example.

References

- [1] T. Abe and A. Dimca, On the splitting types of bundles of logarithmic vector fields along plane curves, *Internat. J. Math.*, **29** (2018), no. 8, 1850055, 20pp. Zbl 1394.14020 MR 3835732
- [2] E. Artal Bartolo, L. Gorrochategui, I. Luengo, and A. Melle-Hernández, On some conjectures about free and nearly free divisors, in *Singularities and Computer Algebra. Festschrift for Gert-Martin Greuel on the Occasion of his 70th Birthday*, 1–19, Springer, Cham, 2017. Zbl 1386.14119 MR 3675719
- [3] D. Cook II, B. Harbourne, J. Migliore, and U. Nagel, Line arrangements and configuration of points with an unusual geometric property, *Compos. Math.*, **154** (2018), no. 10, 2150–2194. Zbl 07036902 MR 3867298
- [4] A. Dimca, *Hyperplane Arrangements. An Introduction*, Universitext, Springer, Cham, 2017. Zbl 1362.14001 MR 3618796
- [5] A. Dimca, Freeness versus maximal global Tjurina number for plane curves, *Math. Proc. Cambridge Phil. Soc.*, **163** (2017), 161–172. Zbl 1387.14080 MR 3656354
- [6] A. Dimca, D. Ibadula, and A. Măcinic, Numerical invariants and moduli spaces for line arrangements, to appear in *Osaka J. Math.* arXiv:1609.06551
- [7] A. Dimca and D. Popescu, Hilbert series and Lefschetz properties of dimension one almost complete intersections, *Comm. Algebra*, **44** (2016), no. 10, 4467–4482. Zbl 06621098 MR 3508312
- [8] A. Dimca and E. Sernesi, Syzygies and logarithmic vector fields along plane curves, *J. Éc. polytech. Math.*, **1** (2014), 247–267. Zbl 1327.14049 MR 3322789

- [9] A. Dimca and G. Sticlaru, On the exponents of free and nearly free projective plane curves, *Rev. Mat. Complut.*, **30** (2017), no. 2, 259–268. Zbl 06802578 MR 3642034
- [10] A. Dimca and G. Sticlaru, Free divisors and rational cuspidal plane curves, *Math. Res. Lett.*, **24** (2017), no. 4, 1023–1042. Zbl 1390.14083 MR 3723802
- [11] A. Dimca and G. Sticlaru, Free and nearly free curves vs. rational cuspidal plane curves, *Publ. Res. Inst. Math. Sci.*, **54** (2018), no. 1, 163–179. Zbl 1391.14057 MR 3749348
- [12] A. A. du Plessis and C. T. C. Wall, Application of the theory of the discriminant to highly singular plane curves, *Math. Proc. Cambridge Phil. Soc.*, **126** (1999), no. 2, 259–266. Zbl 0926.14012 MR 1670229
- [13] A. A. du Plessis and C. T. C. Wall, Curves in $P^2(\mathbb{C})$ with 1-dimensional symmetry, *Rev. Mat. Complut.*, **12** (1999), no. 1, 117–132. Zbl 0969.14022 MR 1698902
- [14] J. Fernández de Bobadilla, I. Luengo, A. Melle-Hernández, and A. Némethi, Classification of rational unicuspidal projective curves whose singularities have one Puiseux pair, in *Real and complex singularities (São Carlos, 2004)*, 31–45, Trends in Mathematics, Birkhäuser, Basel, 2007. Zbl 1120.14019 MR 2280130
- [15] T. K. Moe, Rational Cuspidal Curves, Masters thesis, University of Oslo, Norway, 2008, 139pp. arXiv:1511.02691
- [16] M. Namba, *Geometry of projective algebraic curves*, Monographs and Textbooks in Pure and Applied Mathematics, 88, Marcel Dekker Inc., New York, 1984. Zbl 0556.14012 MR 768929
- [17] K. Palka, A new proof of the theorems of Lin–Zaidenberg and Abhyankar–Moh–Suzuki, *J. Algebra Appl.*, **14** (2015), no. 9, 1540012, 15pp. Zbl 1326.14073 MR 3368264
- [18] K. Saito, Quasihomogene isolierte Singularitäten von Hyperflächen, *Invent. Math.*, **14** (1971), 123–142. Zbl 0224.32011 MR 294699
- [19] F. Sakai and K. Tono, Rational cuspidal curves of type $(d, d-2)$ with one or two cusps, *Osaka J. Math.*, **37** (2000), 405–415. Zbl 0969.14020 MR 1772840
- [20] H. Schenck and S. O. Tohăneanu, Freeness of conic-line arrangements in \mathbb{P}^2 , *Comment. Math. Helv.*, **84** (2009), no. 2, 235–258. Zbl 1183.52014 MR 2495794
- [21] E. Sernesi, The local cohomology of the jacobian ring, *Doc. Math.*, **19** (2014), 541–565. Zbl 1314.14089 MR 3218781
- [22] A. Simis and S. O. Tohăneanu, Homology of homogeneous divisors, *Israel J. Math.*, **200** (2014), no. 1, 449–487. Zbl 1327.13051 MR 3219587
- [23] D. van Straten and T. Warmt, Gorenstein duality for one-dimensional almost complete intersections—with an application to non-isolated real singularities, *Math. Proc. Cambridge Phil. Soc.*, **158** (2015), no. 2, 249–268. Zbl 1371.14006 MR 3310244
- [24] M. G. Zaidenberg and V. Ya. Lin, An irreducible, simply connected algebraic curve in \mathbb{C}^2 is equivalent to a quasihomogeneous curve, *Dokl. Akad. Nauk SSSR*, **271** (1983), no. 5, 1048–1052. Zbl 0564.14014 MR 722017

Received July 28, 2017

A. Dimca, Université Côte d’Azur, CNRS, LJAD,
06108 Nice CEDEX 2, France
E-mail: alexandru.dimca@unice.fr