Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 94 (2019)

Heft: 4

Artikel: Characterization of generic projective space bundles and algebraicity of

foliations

Autor: Araujo, Carolina / Druel, Stéphane

DOI: https://doi.org/10.5169/seals-847032

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Characterization of generic projective space bundles and algebraicity of foliations

Carolina Araujo and Stéphane Druel

Abstract. In this paper we consider various notions of positivity for distributions on complex projective manifolds. We start by analyzing distributions having big slope with respect to curve classes, obtaining characterizations of generic projective space bundles in terms of movable curve classes. We then apply this result to investigate algebraicity of leaves of foliations, providing a lower bound for the algebraic rank of a foliation in terms of invariants measuring positivity. We classify foliations attaining this bound, and describe those whose algebraic rank slightly exceeds this bound.

Mathematics Subject Classification (2010). 14M22, 14J40, 37F75.

Keywords. Holomorphic foliations, rationally connected varieties.

1. Introduction

The existence of sufficiently positive subsheaves of the tangent bundle of a complex projective manifold X imposes strong restrictions on X. In particular, several special varieties can be characterized by positivity properties of their tangent bundle. Early results in this direction include Kobayashi and Ochiai's characterizations of projective spaces and hyperquadrics [23] and Mori's characterization of projective spaces [26]. There are many ways of measuring positivity of a torsion free sheaf. One way is to consider slopes with respect to movable curve classes $\alpha \in \text{Mov}(X)$. We refer to Section 2.3 for the notion of slope and its properties. When α is an ample class, we have the following characterization of projective spaces due to Höring.

Theorem 1.1. Let X be an n-dimensional complex normal projective variety, \mathcal{L} an ample line bundle on X, and $\mathcal{F} \subseteq T_X$ a distribution. Set $\alpha := [\mathcal{L}^{n-1}] \in \text{Mov}(X)$. If $\mu_{\alpha}(\mathcal{F} \otimes \mathcal{L}^*) > 0$, then

- $(X, \mathcal{L}) \cong (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$ [22, Theorem 1.1], and
- $\mathcal{F} = T_{\mathbb{P}^n}$ [5, Theorem 1.3].

Projective space bundles provide counter-examples to the statement of Theorem 1.1 if we replace the ample class $\alpha = [\mathcal{L}^{n-1}]$ with more general movable curve

classes. More precisely, let Y be a complex projective manifold, and \mathcal{E} an ample vector bundle of rank $r+1 \geq 2$ on Y. Consider the projectivization $X = \mathbb{P}_Y(\mathcal{E})$, with tautological line bundle $\mathcal{O}_X(1)$ and natural morphism $\pi\colon X\to Y$. Let $\alpha\in \operatorname{Mov}(X)$ be the class of a line contained in a fiber of π . Then

$$\mu_{\alpha}(T_{X/Y}\otimes\mathcal{O}_X(-1))=\frac{1}{r}>0.$$

So there is an open neighborhood $U \subset \operatorname{Mov}(X)$ of α such that $\mu_{\beta}(T_{X/Y} \otimes \mathcal{O}_X(-1)) > 0$ for every $\beta \in U$. The following characterization of generic \mathbb{P}^r -bundles shows that these are all the new examples that arise when the ample class $\alpha = [\mathcal{L}^{n-1}]$ is replaced with an arbitrary movable class $\alpha \in \operatorname{Mov}(X)$ in Theorem 1.1. By a generic \mathbb{P}^r -bundle we mean an almost proper dominant map $X \dashrightarrow Y$ to a normal projective variety Y with general fibers isomorphic to \mathbb{P}^r . See Section 2.3 for the notion of μ_{α}^{\max} .

Theorem 1.2. Let X be a normal \mathbb{Q} -factorial complex projective variety, and \mathcal{L} an ample line bundle on X. If $\mu_{\alpha}^{\max}(T_X \otimes \mathcal{L}^*) > 0$ for some movable curve class $\alpha \in \text{Mov}(X)$, then X is a generic \mathbb{P}^r -bundle for some positive integer r.

Theorem 1.2 follows from the more refined statement in Theorem 3.1.

Corollary 1.3. Let X be a normal \mathbb{Q} -factorial complex projective variety, and suppose that X is not a generic \mathbb{P}^r -bundle for any positive integer r. Let \mathcal{L} be an ample line bundle on X. Then, for any positive integer m and any torsion-free quotient $(\Omega^1_X \otimes \mathcal{L})^{\otimes m} \to \mathcal{Q}$ of positive rank, $\det(\mathcal{Q})$ is pseudo-effective.

Next we apply these results to investigate algebraicity of leaves of holomorphic foliations on complex projective manifolds.

A central problem in the theory of holomorphic foliations is to find conditions that guarantee the existence of algebraic leaves. Algebraic leaves of holomorphic foliations correspond to algebraic solutions of complex differential equations. It has been noted that positivity of foliations on complex projective varieties tend to improve algebraicity of leaves ([1–4, 6, 7, 10–12, 14, 15, 25]). In order to measure algebraicity of leaves, we introduce the *algebraic rank*. The algebraic rank $r^a(\mathcal{F})$ of a holomorphic foliation \mathcal{F} on a complex algebraic variety X is the maximum dimension of an algebraic subvariety through a general point of X that is tangent to \mathcal{F} . These maximal algebraic subvarieties tangent to \mathcal{F} are the leaves of a subfoliation $\mathcal{F}^a \subseteq \mathcal{F}$, the *algebraic part* of \mathcal{F} (see Definition 2.4).

In a series of papers, we have addressed $Fano\ foliations\ ([1-4])$. These are holomorphic foliations $\mathcal F$ on complex projective varieties with ample anti-canonical class $-K_{\mathcal F}$. For a Fano foliation $\mathcal F$ on a complex projective manifold X, a rough measure of positivity is the $index\ \iota(\mathcal F)$, which is the largest integer dividing $-K_{\mathcal F}$ in Pic(X). Our works on Fano foliations with high index indicated that the larger is the index, the larger is the algebraic rank of the Fano foliation. Now we investigate this relation between positivity and algebraicity of leaves for a wider class of foliations than Fano foliations.

We first introduce a new invariant measuring positivity for foliations with big anticanonical class.

Definition 1.4. Let X be a complex projective manifold, and \mathcal{F} a holomorphic foliation on X with big anti-canonical class $-K_{\mathcal{F}}$. The *generalised index* of \mathcal{F} is defined as follows (see Lemma 4.1):

$$\hat{\iota}(\mathcal{F}) := \max \{ t \in \mathbb{R} \mid -K_{\mathcal{F}} \equiv tA + E, \text{ where } A \text{ is an ample divisor}$$
 and $E \text{ is a pseudo-effective } \mathbb{R}\text{-divisor} \}.$

We provide a lower bound for the algebraic rank in terms of the generalised index, and classify foliations attaining this bound. We refer to Definition 2.4 for the notions of pull-back and purely transcendental foliations.

Theorem 1.5. Let X be an n-dimensional complex projective manifold, and $\mathcal{F} \subsetneq T_X$ a foliation on X with big anti-canonical class. Then the algebraic rank and the generalised index of \mathcal{F} satisfy

$$r^a(\mathcal{F}) \geqslant \hat{\iota}(\mathcal{F}).$$

Moreover, equality holds if and only if $X \cong \mathbb{P}^n$ and \mathcal{F} is the linear pull-back of a purely transcendental foliation on $\mathbb{P}^{n-r^a(\mathcal{F})}$ with zero canonical class.

The following are immediate consequences of Theorem 1.5.

Corollary 1.6. Let X be an n-dimensional complex projective manifold, and $\mathcal{F} \subsetneq T_X$ a Fano foliation of index $\iota(\mathcal{F})$ on X. Then the algebraic rank of \mathcal{F} satisfies the following inequality:

$$r^a(\mathcal{F}) \geqslant \iota(\mathcal{F}).$$

Moreover, equality holds if and only if $X \cong \mathbb{P}^n$ and \mathcal{F} is the linear pull-back of a purely transcendental foliation on $\mathbb{P}^{n-r^a(\mathcal{F})}$ with zero canonical class.

Corollary 1.7. Let X be an n-dimensional complex projective manifold, L an ample divisor on X, and $\mathcal{F} \subsetneq T_X$ a foliation of rank r on X. Suppose that $-K_{\mathcal{F}} - rL$ is pseudo-effective. Then $(X, \mathcal{O}_X(L)) \cong (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$, and \mathcal{F} is induced by a linear projection of \mathbb{P}^n .

Finally, we address foliations \mathcal{F} whose algebraic rank and generalised index satisfy $r^a(\mathcal{F}) \leq \hat{\iota}(\mathcal{F}) + 1$.

Theorem 1.8. Let X be a complex projective manifold, and $\mathcal{F} \subsetneq T_X$ a foliation on X with big anti-canonical class. Suppose that the algebraic rank and the generalised index of \mathcal{F} satisfy $r^a(\mathcal{F}) \leqslant \hat{\iota}(\mathcal{F}) + 1$. Then the closure of a general leaf of \mathcal{F}^a is rationally connected.

The following is an immediate consequence of Theorem 1.8.

Corollary 1.9. Let X be a complex projective manifold, and $\mathcal{F} \subsetneq T_X$ a Fano foliation of index $\iota(\mathcal{F})$ on X. Suppose that the algebraic rank of \mathcal{F} satisfies $r^a(\mathcal{F}) = \iota(\mathcal{F}) + 1$. Then the closure of a general leaf of \mathcal{F}^a is rationally connected.

We remark that codimension 1 Fano foliations \mathcal{F} of index n-3 on n-dimensional projective manifolds were classified in [4]. When $X \ncong \mathbb{P}^n$, then either \mathcal{F} is algebraically integrable or $r^a(\mathcal{F}) = \iota(\mathcal{F}) + 1$.

Notation and conventions. We always work over the field \mathbb{C} of complex numbers. Varieties are always assumed to be irreducible. We denote by X_{ns} the nonsingular locus of a variety X. When X is a normal variety, and \mathcal{F} is a quasi-coherent sheaf of generic rank r on X, we denote by T_X the sheaf $(\Omega_X^1)^*$, and by $\det(\mathcal{F})$ the sheaf $(\wedge^r \mathcal{F})^{**}$. If \mathcal{E} is a vector bundle on a variety X, we denote by $\mathbb{P}_X(\mathcal{E})$ the Grothendieck projectivization $\operatorname{Proj}_X(S^{\bullet}\mathcal{E})$.

Acknowledgements. This project was supported by the Brazilian-French Network in Mathematics. Much of this work was developed during the authors' visits to IMPA and Institut Fourier. We would like to thank both institutions for their support and hospitality.

The authors would also like to thank the anonymous referees for their helpful and very detailed reports.

The first named author was partially supported by CNPq and Faperj Research Fellowships. The second named author was partially supported by the ALKAGE project (ERC grant Nr 670846, 2015–2020) and by the project Foliage of Agence Nationale de la Recherche (ANR-16-CE40-0008-01).

2. Foliations

2.1. Basic notions. Let *X* be a normal variety.

Definition 2.1. A distribution on X is a coherent saturated subsheaf $\mathcal{F} \subset T_X$. By saturated we mean that the quotient T_X/\mathcal{F} is torsion-free.

The rank r of \mathcal{F} is the generic rank of \mathcal{F} . The codimension of \mathcal{F} is defined as $q := \dim X - r$.

The *normal sheaf* of \mathcal{F} is the sheaf $\mathcal{N}_{\mathcal{F}} := (T_X/\mathcal{F})^{**}$.

The *canonical class* $K_{\mathcal{F}}$ of \mathcal{F} is any Weil divisor on X such that $\mathcal{O}_X(-K_{\mathcal{F}}) \cong \det(\mathcal{F})$. We say that \mathcal{F} is \mathbb{Q} -Gorenstein if $K_{\mathcal{F}}$ is a \mathbb{Q} -Cartier divisor.

Definition 2.2. A *foliation* on X is a distribution $\mathcal{F} \subset T_X$ that is closed under the Lie bracket.

Let $X^{\circ} \subset X_{\rm ns}$ be the maximal open subset such that $\mathcal{F}_{|X_{\rm ns}}$ is a subbundle of $T_{X_{\rm ns}}$. By Frobenius' theorem, through any point of X° there is a maximal connected and immersed holomorphic submanifold $L \subset X^{\circ}$ such that $T_L = \mathcal{F}_{|L}$. We say that such L is a *leaf* of \mathcal{F} . A leaf is called *algebraic* if it is open in its Zariski closure.

2.3. To a codimension q distribution \mathcal{F} on a normal variety X, one naturally associates a unique (up to scaling) twisted q-form $\omega_{\mathcal{F}} \in H^0(X, \Omega_X^q \otimes \det(\mathcal{N}_{\mathcal{F}}))$.

This form does not vanish in codimension 1, and completely determines the distribution \mathcal{F} . (See for instance [3, Paragraph 3.5] for details.)

Definition 2.4 (The algebraic and transcendental parts of a foliation [2, Definition 2]). Let \mathcal{F} be a foliation of rank r on a normal variety X. There exist a normal variety Y, unique up to birational equivalence, a dominant rational map with connected fibers $\varphi: X \dashrightarrow Y$, and a foliation \mathcal{G} on Y such that the following conditions hold (see [24, Section 2.3]).

- (1) \mathcal{G} is purely transcendental, i.e. there is no positive-dimensional algebraic subvariety through a general point of Y that is tangent to \mathcal{G} .
- (2) \mathcal{F} is the pull-back of \mathcal{G} via φ . This means the following. Let $X^{\circ} \subset X$ and $Y^{\circ} \subset Y$ be smooth open subsets such that φ restricts to a smooth morphism $\varphi^{\circ}: X^{\circ} \to Y^{\circ}$. Then $\mathcal{F}_{|X^{\circ}} = (d\varphi^{\circ})^{-1}(\mathcal{G}_{|Y^{\circ}})$. In this case we write $\mathcal{F} = \varphi^{-1}\mathcal{G}$.

The foliation \mathcal{F}^a on X induced by φ is called the *algebraic part* of \mathcal{F} , and its rank is the *algebraic rank* of \mathcal{F} , which we denote by r^a . When $r^a = r$, we say that \mathcal{F} is *algebraically integrable*. The foliation $\mathcal{G} \subset T_Y$ is called the *transcendental part* of \mathcal{F} .

Next we relate the canonical class of a foliation with those of its algebraic and transcendental parts. For that we introduce some notation.

Definition 2.5. Let $\pi: X \to Y$ be a dominant morphism of normal varieties.

Let D be a Weil \mathbb{Q} -divisor on Y. If $\pi\colon X\to Y$ is equidimensional, we define the pull-back π^*D of D to be the unique \mathbb{Q} -divisor on X whose restriction to $\pi^{-1}(Y_{\rm ns})$ is $(\pi_{|\pi^{-1}(Y_{\rm ns})})^*D_{|\pi^{-1}(Y_{\rm ns})}$. This agrees with the usual pull-back if D is \mathbb{Q} -Cartier.

We define the *ramification divisor* $R(\pi)$ of π as follows. Let $Y^{\circ} \subset Y$ be a dense open subset such that $\operatorname{codim}(Y \setminus Y^{\circ}) \geq 2$ and π restricts to an equidimensional morphism π° : $X^{\circ} = \pi^{-1}(Y^{\circ}) \to Y^{\circ}$. Set

$$R(\pi^{\circ}) = \sum_{D^{\circ}} ((\pi^{\circ})^* D^{\circ} - ((\pi^{\circ})^* D^{\circ})_{\text{red}}),$$

where D° runs through all prime divisors on Y° . Then $R(\pi)$ is the Zariski closure of $R(\pi^{\circ})$ in X.

Assume that either K_Y is \mathbb{Q} -Cartier, or that π is equidimensional. We define the relative canonical divisor of X over Y as $K_{X/Y} := K_X - \pi^* K_Y$.

2.6 (The canonical class of a pull-back foliation). Let $\pi: X \to Y$ be a dominant morphism with connected fibers between normal varieties, \mathcal{G} a foliation on Y, and $\mathcal{F} = \pi^{-1}\mathcal{G} \subset T_X$ its pull-back via π , as in Definition 2.4 (2). Assume that π is equidimensional. Let $\{B_i\}_{i\in I}$ be the (possibly empty) set of prime divisors on Y contained in the set of critical values of π and invariant by \mathcal{G} . A straightforward computation shows that

$$K_{\mathcal{F}} = \pi^* K_{\mathcal{G}} + K_{X/Y} - \sum_{i \in I} (\pi^* B_i - (\pi^* B_i)_{\text{red}}).$$
 (2.1)

In particular, if \mathcal{F} is induced by π , then (2.1) reads

$$K_{\mathcal{F}} = K_{X/Y} - R(\pi). \tag{2.2}$$

Remark 2.7. Let $\varphi: X \dashrightarrow Y$ be a dominant rational map with connected fibers between normal varieties, and \mathcal{F} a foliation on X. Suppose that the general fiber of φ is tangent to \mathcal{F} . This means that, for a general point x on a general fiber F of φ , the linear subspace $\mathcal{F}_x \subset T_x X$ determined by the inclusion $\mathcal{F} \subset T_X$ contains $T_x F$. Then, by [1, Lemma 6.7], there is a foliation \mathcal{G} on Y such that $\mathcal{F} = \varphi^{-1}\mathcal{G}$. We remark that this is not true in general if \mathcal{F} is just a distribution.

2.8 (Restricting foliations to subvarieties). Let X be a smooth variety, and \mathcal{F} a codimension q foliation on X. Let Z be a smooth subvariety with dim $Z \geqslant q$. Suppose that Z is generically transverse to \mathcal{F} . This means that the associated twisted q-form $\omega_{\mathcal{F}} \in H^0(X, \Omega_X^q \otimes \det(\mathcal{N}_{\mathcal{F}}))$ restricts to a nonzero twisted q-form on Z, and so \mathcal{F} induces a foliation \mathcal{F}_Z of codimension q on Z. Then there is an effective divisor B on Z such that \mathcal{F}_Z corresponds to a twisted q-form in $H^0(Z, \Omega_Z^q \otimes \det(\mathcal{N}_{\mathcal{F}})_{|Z}(-B))$ non vanishing in codimension 1. A straightforward computation shows that

$$K_{\mathcal{F}_Z} + K_{X|Z} = K_{\mathcal{F}|Z} + K_Z - B.$$

When Z is a general hyperplane section, we show that B = 0.

Lemma 2.9. Let $X \subset \mathbb{P}^N$ be a smooth projective variety, and \mathcal{F} a foliation of codimension $q \leq \dim X - 2$ on X. Let $H \subset \mathbb{P}^N$ be a general hyperplane. Then \mathcal{F} induces a foliation $\mathcal{F}_{X \cap H}$ of codimension q on $X \cap H$ with

$$K_{\mathcal{F}_{X\cap H}} = (K_{\mathcal{F}} + H)_{|X\cap H}.$$

Proof. Let $H \subset \mathbb{P}^N$ be a general hyperplane, and set $Z := X \cap H$. Notice that Z is smooth and that Z is generically transverse to \mathcal{F} . Let B be the effective divisor on Z introduced in 2.8. If B_1 is a prime divisor on Z, then $B_1 \subset \operatorname{Supp}(B)$ if and only if \mathcal{F} is tangent to H at a general point of B_1 .

Let $X^{\circ} \subset X$ be the open subset where \mathcal{F} is a subbundle of T_X , and consider the incidence subset

$$I^{\circ} = \{(x, H) \in X^{\circ} \times (\mathbb{P}^N)^* \mid x \in H \text{ and } \mathcal{F}_x \subset T_x H\}.$$

An easy dimension count gives that

$$\dim I^{\circ} = \dim X + N - (\dim X - q) - 1 \leqslant \dim X + N - 3.$$

It follows that if H is general, then B = 0.

2.2. Algebraically integrable foliations. Throughout this subsection, we let X be a normal projective variety, and \mathcal{F} a \mathbb{Q} -Gorenstein algebraically integrable foliation on X.

Definition 2.10 ([2, Definition 3.11]). Let $i: F \to X$ be the normalization of the closure of a general leaf of \mathcal{F} . There is a canonically defined effective \mathbb{Q} -divisor Δ_F on F such that $K_F + \Delta_F \sim_{\mathbb{Q}} i^* K_{\mathcal{F}}$. The pair (F, Δ_F) is called a *general log leaf* of \mathcal{F} .

In the setup of Definition 2.10, we often write $L_{|F}$ for the pull-back i^*L of a Cartier divisor L on X.

2.11 (The family of log leaves of \mathcal{F} [2, Lemma 3.9 and Remark 3.12]). There is a unique proper subvariety Y' of the Chow variety of X whose general point parametrizes the closure of a general leaf of \mathcal{F} (viewed as a reduced and irreducible cycle in X). Let Y be the normalization of Y', and $Z \to Y' \times X$ the normalization of the universal cycle, with induced morphisms:

$$Z \xrightarrow{\nu} X$$

$$\pi \downarrow \qquad (2.3)$$

$$Y$$

Then $\nu: Z \to X$ is birational and, for a general point $y \in Y$, $\nu(\pi^{-1}(y)) \subset X$ is the closure of a leaf of \mathcal{F} . We refer to the diagram (2.3) as the *family of leaves* of \mathcal{F} .

Let \mathcal{F}_Z be the foliation on Z induced by \mathcal{F} (or π). By (2.2), $K_{\mathcal{F}_Z} = K_{Z/Y} - R(\pi)$. Moreover, there is a canonically defined effective Weil \mathbb{Q} -divisor Δ on Z such that

$$K_{\mathcal{F}_Z} + \Delta = K_{Z/Y} - R(\pi) + \Delta \sim_{\mathbb{Q}} \nu^* K_{\mathcal{F}}.$$
 (2.4)

Note that Δ is ν -exceptional since $\nu_* K_{\mathcal{F}_Z} = K_{\mathcal{F}}$.

Let $y \in Y$ be a general point, set $Z_y := \pi^{-1}(y)$ and $\Delta_y := \Delta_{|Z_y|}$. Then (Z_y, Δ_y) coincides with the general log leaf (F, Δ_F) from Definition 2.10.

We will need the following observation.

Lemma 2.12. Let X be a smooth projective variety, and \mathcal{F} an algebraically integrable foliation on X. In the setup of Paragraph 2.11 above, we have

$$\operatorname{Supp}(\Delta_y) = \operatorname{Exc}(v) \cap Z_y.$$

In particular, the singular locus of Z_{ν} is contained in Supp (Δ_{ν}) .

Proof. Since Δ is ν -exceptional and $\operatorname{Supp}(\Delta_y) = \operatorname{Supp}(\Delta) \cap Z_y$, we must have $\operatorname{Supp}(\Delta_y) \subset \operatorname{Exc}(\nu) \cap Z_y$.

To prove that $\operatorname{Exc}(\nu) \cap Z_{\nu} \subset \operatorname{Supp}(\Delta_{\nu})$, we first reduce to the case when \mathcal{F} has rank 1. Suppose that \mathcal{F} has rank $r \geq 2$, and consider an embedding $X \subset \mathbb{P}^N$.

Let $L \subset \mathbb{P}^N$ be a general linear subspace of codimension r-1. By Lemma 2.9, \mathcal{F} induces a foliation by curves $\mathcal{F}_{X \cap L}$ on $X \cap L$ with

$$K_{\mathcal{F}_{X\cap L}} = \left(K_{\mathcal{F}} + (r-1)H\right)_{|X\cap L}, \qquad (2.5)$$

where H is a hyperplane in \mathbb{P}^N . Notice that the closure of a general leaf of $\mathcal{F}_{X\cap L}$ is $\nu(Z_y)\cap L$ for a general point $y\in Y$. Moreover, the general log leaf of $\mathcal{F}_{X\cap L}$ is $(Z_y\cap \nu^{-1}(L),\Delta_y\cap \nu^{-1}(L))$. This follows from the definition of general log leaf, (2.5), and the usual adjunction formula. Let E be an irreducible component of $\mathrm{Exc}(\nu)$ meeting Z_y . Since the restriction of ν to $E\cap Z_y$ is finite, we have $\dim \nu(E)\geqslant r-1$. In particular, $\nu(E)\cap L\neq\emptyset$. Moreover, $\mathrm{Exc}(\nu)\cap Z_y\subset \mathrm{Supp}(\Delta_y)$ if and only if $\mathrm{Exc}(\nu)\cap Z_y\cap \nu^{-1}(L)\subset \mathrm{Supp}(\Delta_y)\cap \nu^{-1}(L)$. Since $\mathrm{Exc}(\nu)\cap \nu^{-1}(L)=\mathrm{Exc}(\nu|_{\nu^{-1}(L)})$, we may assume that \mathcal{F} has rank 1.

When \mathcal{F} has rank 1, \mathcal{F} is a line bundle on X by [1, Remark 2.3] and [20, Proposition 1.9]. If E is an irreducible component of $\operatorname{Exc}(\nu)$ that dominates Y, then $\nu(E)$ is contained in the singular locus of \mathcal{F} . Apply [13, Lemme 1.2] (see also [1, Lemma 5.6]) to conclude that $E \cap Z_y \subset \operatorname{Supp}(\Delta_y)$, completing the proof of the lemma.

Corollary 2.13. Let X be a smooth projective variety, and $\mathcal{F} \subsetneq T_X$ an algebraically integrable foliation on X, with general log leaf (F, Δ_F) . Suppose that either $\rho(X) = 1$, or \mathcal{F} is a Fano foliation. Then $\Delta_F \neq 0$.

Proof. Let the notation be as in Paragraph 2.11, and suppose that $\Delta_F = 0$. It follows from Lemma 2.12 that no irreducible component of $\operatorname{Exc}(v)$ dominates Y. Hence, \mathcal{F} is induced by a rational map $X \dashrightarrow Y$ that restricts to a smooth proper morphism on a dense open subset of X. This is impossible if $\rho(X) = 1$. Moreover, F is smooth, and in particular log canonical. On the other hand, an algebraically integrable Fano foliation whose general log leaf is log canonical has the special property that there is a common point in the closure of a general leaf (see [1, Proposition 5.3] and [3, Proposition 3.13]). We conclude that $\Delta_F \neq 0$.

We end this subsection with a consequence of [21, Lemma 2.14] (see also [10,11,15] for related results). Recall that a Weil \mathbb{Q} -divisor D on a normal projective variety X is said to be *pseudo-effective* if, for any ample divisor L on X and any rational number $\varepsilon > 0$, there exists an effective Weil \mathbb{Q} -divisor E such that $D + \varepsilon L \sim_{\mathbb{Q}} E$.

Lemma 2.14. Let X be a normal projective variety, L an ample divisor on X, and \mathcal{F} a \mathbb{Q} -Gorenstein algebraically integrable foliation on X. Let F be the normalization of the closure of a general leaf of \mathcal{F} , and let $v_F \colon \widehat{F} \to F$ be a resolution of singularities. If $K_{\widehat{F}} + v_F^* L_{|F}$ is pseudo-effective, then so is $K_{\mathcal{F}} + L$.

Proof. Consider the family of leaves of \mathcal{F} as in Paragraph 2.11:

$$Z \xrightarrow{\nu} X$$

$$\psi \downarrow \qquad \qquad Y.$$

By [15, Lemma 4.2], there exists a finite surjective morphism $\mu_1: Y_1 \to Y$ with Y_1 normal and connected satisfying the following property. If Z_1 denotes the normalization of the product $Y_1 \times_Y Z$, then the induced morphism $\psi_1: Z_1 \to Y_1$ has reduced fibers over codimension one points in Y_1 . Let $\nu_1: Z_1 \to Z$ be the natural morphism. Let $\mu_2: Y_2 \to Y_1$ be a resolution of singularities, and Z_2 a resolution of singularities of the product $Y_2 \times_{Y_1} Z_1$, with natural morphism $\nu_2: Z_2 \to Z_1$. We have a commutative diagram:

$$Z_{2} \xrightarrow{\nu_{2}, \text{ birational}} Z_{1} \xrightarrow{\nu_{1}, \text{ finite}} Z \xrightarrow{\nu, \text{ birational}} X$$

$$\downarrow \psi_{2} \qquad \qquad \downarrow \psi_{1} \qquad \qquad \downarrow \psi$$

$$Y_{2} \xrightarrow{\mu_{2}, \text{ birational}} Y_{1} \xrightarrow{\mu_{1}, \text{ finite}} Y.$$

Let $F_1 \cong F$ be a general fiber of ψ_1 , and set $L_1 := (\nu \circ \nu_1)^* L$. Let F_2 be the fiber of $\psi_2 : Z_2 \to Y_2$ mapping to F_1 , and set $L_2 := \nu_2^* L_1$.

By Paragraph 2.11, there is a canonically defined ν -exceptional effective $\mathbb Q$ -Weil divisor Δ on Z such that

$$K_{Z/Y} - R(\psi) + \Delta \sim_{\mathbb{Q}} \nu^* K_{\mathcal{F}},$$
 (2.6)

where $R(\psi)$ denotes the ramification divisor of ψ . Moreover, a straightforward computation shows that

$$\nu_1^* (K_{Z/Y} - R(\psi)) = K_{Z_1/Y_1}. \tag{2.7}$$

Suppose that $K_{\widehat{F}} + \nu_F^* L_{|F}$ is pseudo-effective. We may assume without loss of generality that the restriction of $\nu_2 \colon Z_2 \to Z_1$ to F_2 factors through $\widehat{F} \to F \cong F_1$. This implies that $K_{F_2} + L_{2|F_2}$ is pseudo-effective as well. By [21, Lemma 2.14], $K_{Z_2/Y_2} + L_2$ is also pseudo-effective. This implies that $(\nu \circ \nu_1 \circ \nu_2)_* (K_{Z_2/Y_2} + L_2)$ is too. Set $Z_1^\circ := Z_1 \setminus \nu_2(\operatorname{Exc}(\nu_2))$ and $Y_1^\circ := Y_1 \setminus \mu_2(\operatorname{Exc}(\mu_2))$. Since ψ_1 is equidimensional, we have $\psi_1(Z_1^\circ) \subset Y_1^\circ$.

Notice that codim $Y_1 \setminus Y_1^{\circ} \ge 2$, codim $Z_1 \setminus Z_1^{\circ} \ge 2$, and that ν_2 (respectively, μ_2) induces an isomorphism

$$v_2^{-1}(Z_1 \setminus Z_1^{\circ}) \cong Z_1 \setminus Z_1^{\circ}$$

(respectively, $\mu_2^{-1}(Y_1 \setminus Y_1^{\circ}) \cong Y_1 \setminus Y_1^{\circ}$). It follows that

$$(\nu_2)_*(K_{Z_2/Y_2} + L_2) = K_{Z_1/Y_1} + L_1.$$

From (2.6) and (2.7), we conclude that

$$(\nu \circ \nu_1 \circ \nu_2)_*(K_{Z_2/Y_2} + L_2) = \deg(\nu_1)(K_{\mathcal{F}} + L)$$

is pseudo-effective, completing the proof of the lemma.

2.3. Foliations defined by stability conditions. Let X be a normal projective variety, and $\mathcal{F} \subset T_X$ a foliation on X. Harder–Narasimhan filtrations of \mathcal{F} allow one to construct subfoliations of \mathcal{F} that inherit some of the positivity properties of \mathcal{F} (see for instance [1, Section 7]). However, the classical notion of slope-stability with respect to an ample line bundle is not flexible enough to be applied in many situations in birational geometry. The papers [9] and [17] extend a number of known results from the classical case to the setting where stability conditions are given by movable curve classes.

Definition 2.15. A curve class $\alpha \in N_1(X)_{\mathbb{R}}$ is *movable* if $D \cdot \alpha \geq 0$ for all effective Cartier divisors D on X. The set of movable classes is a closed convex cone $\text{Mov}(X) \subset N_1(X)_{\mathbb{R}}$, called the *movable cone* of X. If X is smooth, then Mov(X) is the closure of the convex cone in $N_1(X)_{\mathbb{R}}$ generated by classes of curves whose deformations cover a dense subset of X by [8].

2.16 (The Harder–Narasimhan filtration with respect to a movable curve class). Let X be a normal, \mathbb{Q} -factorial, projective variety, $\alpha \in N_1(X)_{\mathbb{R}}$ a movable curve class, and \mathcal{F} a torsion-free sheaf of positive rank on X.

The slope of \mathcal{F} with respect to α is the real number

$$\mu_{\alpha}(\mathcal{F}) = \frac{\det(\mathcal{F}) \cdot \alpha}{\operatorname{rank}(\mathcal{F})}.$$

The sheaf \mathcal{F} is α -semistable if, for any subsheaf $\mathcal{E} \neq 0$ of \mathcal{F} , one has $\mu_{\alpha}(\mathcal{E}) \leq \mu_{\alpha}(\mathcal{F})$.

The maximal and minimal slopes of \mathcal{F} with respect to α are defined by

$$\begin{split} &\mu_{\alpha}^{\max}(\mathcal{F}) := \sup \big\{ \mu_{\alpha}(\mathcal{E}) \mid 0 \neq \mathcal{E} \subseteq \mathcal{F} \text{ is a coherent subsheaf} \big\}, \\ &\mu_{\alpha}^{\min}(\mathcal{F}) := \inf \big\{ \mu_{\alpha}(\mathcal{Q}) \mid \mathcal{Q} \neq 0 \text{ is a torsion-free quotient of } \mathcal{F} \big\}. \end{split}$$

By [17, Corollary 2.26], there exists a unique filtration of $\mathcal F$ by saturated subsheaves

$$0 = \mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \cdots \subseteq \mathcal{F}_k = \mathcal{F}$$
,

with α -semistable quotients $Q_i = \mathcal{F}_i/\mathcal{F}_{i-1}$ such that $\mu_{\alpha}(Q_1) > \mu_{\alpha}(Q_2) > \cdots > \mu_{\alpha}(Q_k)$. This filtration is called the *Harder–Narasimhan filtration* of \mathcal{F} .

Using the basic properties of slopes and the Harder–Narasimhan filtration, one can check that, for $1 \le i \le k$,

$$\mu_{\alpha}^{\min}(\mathcal{F}_i) = \mu_{\alpha}(\mathcal{Q}_i) = \mu_{\alpha}^{\max}(\mathcal{F}/\mathcal{F}_{i-1}). \tag{2.8}$$

The sheaf \mathcal{F}_1 is called the *maximal destabilizing subsheaf* of \mathcal{F} .

Suppose that $\mu_{\alpha}^{\max}(\mathcal{F}) > 0$, and set $s := \max\{1 \le i \le k \mid \mu_{\alpha}(\mathcal{Q}_i) > 0\} \ge 1$. The positive part of \mathcal{F} with respect to α is the sheaf $\mathcal{F}_{\alpha}^+ := \mathcal{F}_s$.

The following is a useful criterion for a subsheaf of a foliation to be a foliation. We include a proof for the reader's convenience.

Lemma 2.17 ([10, Lemma 4.10]). Let X be a normal, \mathbb{Q} -factorial, projective variety and $\mathcal{F} \subset T_X$ a foliation on X. Let $\mathcal{G} \subset \mathcal{F}$ be a saturated subsheaf and suppose that, for some movable curve class $\alpha \in \mathbb{N}_1(X)_{\mathbb{R}}$,

$$2\mu_{\alpha}^{\min}(\mathcal{G}) > \mu_{\alpha}^{\max}(\mathcal{F}/\mathcal{G}).$$

Then \mathcal{G} is also a foliation on X.

Proof. Note that \mathcal{G} is saturated in T_X . Integrability of \mathcal{G} is equivalent to the vanishing of the map $\wedge^2 \mathcal{G} \to \mathcal{F}/\mathcal{G}$ induced by the Lie bracket. This vanishing follows from the inequality

$$\mu_{\alpha}^{\min}(\wedge^2 \mathcal{G}/\operatorname{Tors}) > \mu_{\alpha}^{\max}(\mathcal{F}/\mathcal{G}).$$

So it is enough to prove that

$$\mu_{\alpha}^{\min}(\wedge^2 \mathcal{G}/\operatorname{Tors}) \geqslant 2\mu_{\alpha}^{\min}(\mathcal{G}).$$
 (2.9)

Observe $(\wedge^2 \mathcal{G}/\text{Tors})^{**} \cong (\wedge^2 \mathcal{G})^{**}$ is a direct summand of $(\mathcal{G} \otimes \mathcal{G})^{**}$. Therefore, we have

$$\begin{split} \mu_{\alpha}^{\min}(\wedge^2\mathcal{G}/\operatorname{Tors}) &= \mu_{\alpha}^{\min}((\wedge^2\mathcal{G})^{**}) \\ &\geqslant \mu_{\alpha}^{\min}((\mathcal{G}\otimes\mathcal{G})^{**}) \\ &= 2\mu_{\alpha}^{\min}(\mathcal{G}) \qquad \text{(by [17, Theorem 4.2]),} \end{split}$$

proving (2.9).

Corollary 2.18. In the setup of Paragraph 2.16, suppose that $\mathcal{F} \subset T_X$ is a foliation on X with $\mu_{\alpha}^{\max}(\mathcal{F}) \geq 0$. Then $\mathcal{F}_i \subset T_X$ is also a foliation on X whenever $\mu_{\alpha}(\mathcal{Q}_i) \geq 0$.

We end this subsection with a remarkable result of Campana and Păun concerning algebraic integrability of foliations.

Theorem 2.19. Let X be a normal \mathbb{Q} -factorial projective variety, $\alpha \in N_1(X)_{\mathbb{R}}$ a movable curve class, and $\mathcal{F} \subset T_X$ a foliation on X. Suppose that $\mu_{\alpha}^{\min}(\mathcal{F}) > 0$. Then \mathcal{F} is algebraically integrable, and the closure of a general leaf is rationally connected.

In particular, if \mathcal{F} is purely transcendental, then $K_{\mathcal{F}}$ is pseudo-effective.

Proof. Let $\nu: \hat{X} \to X$ be a resolution of singularities. By [17, Proposition 2.7 and Remark 2.8], we have $\mu_{\alpha}^{\min}(\mathcal{F}) = \mu_{\nu^*\alpha}^{\min}(\nu^{-1}\mathcal{F})$, where $\nu^*\alpha \in \text{Mov}(\hat{X})$ is the numerical pull-back of α . The claim now follows from [10, Theorem 4.7] applied to $\nu^{-1}\mathcal{F}$.

The following is an immediate consequence of Corollary 2.18 and Theorem 2.19. **Corollary 2.20.** In the setup of Paragraph 2.16, suppose that $\mathcal{F} \subset T_X$ is a foliation on X with $\mu_{\alpha}^{\max}(\mathcal{F}) > 0$. Then, for $1 \leq i \leq s$, $\mathcal{F}_i \subset T_X$ is an algebraically integrable foliation, and the closure of a general leaf is rationally connected.

3. Characterization of generic projective space bundles

Theorem 3.1. Let X be a normal \mathbb{Q} -factorial projective variety, and \mathcal{L} an ample line bundle on X. Suppose that $\mu_{\alpha}^{\max}(T_X \otimes \mathcal{L}^*) > 0$ for some movable curve class $\alpha \in \text{Mov}(X)$, and let \mathcal{T}_1 be the maximal destabilizing subsheaf of T_X with respect to α .

- (1) Then \mathcal{T}_1 is induced by a generic \mathbb{P}^{r_1} -bundle structure $\pi_1: X \dashrightarrow Y_1$ on X, and \mathcal{L} restricts to $\mathcal{O}_{\mathbb{P}^{r_1}}(1)$ on a general fiber of π_1 . Moreover, $\mu_{\alpha}^{\max}((T_X/\mathcal{T}_1) \otimes \mathcal{L}^*) \leq 0$.
- (2) If \mathcal{F} is a foliation on X and $\mu_{\alpha}(\mathcal{F} \otimes \mathcal{L}^*) > 0$, then $\mathcal{T}_1 \subseteq \mathcal{F}$, and equality holds if and only if \mathcal{F} is algebraically integrable.

Proof. Step 1. Suppose first that \mathcal{F} is an algebraically integrable foliation of rank r, and $\mu_{\alpha}(\mathcal{F} \otimes \mathcal{L}^*) > 0$ for some movable curve class $\alpha \in \text{Mov}(X)$. We show that \mathcal{F} is induced by a generic \mathbb{P}^r -bundle structure $\pi\colon X \dashrightarrow Y$ on X and that \mathcal{L} restricts to $\mathcal{O}_{\mathbb{P}^r}(1)$ on a general fiber of π .

Let F be the normalization of the closure of a general leaf of \mathcal{F} , and let $\nu_F: \widehat{F} \to F$ be a resolution of singularities. Let L be a divisor on X such that $\mathcal{O}_X(L) \cong \mathcal{L}$. If $K_{\widehat{F}} + \dim F \cdot \nu_F^* L_{|F}$ is pseudo-effective, then so is $K_{\mathcal{F}} + \dim F \cdot L$ by Lemma 2.14. This is absurd since $-r\mu_{\alpha}(\mathcal{F} \otimes \mathcal{L}^*) = (K_{\mathcal{F}} + \dim F \cdot L) \cdot \alpha < 0$ by assumption, proving that $K_{\widehat{F}} + \dim F \cdot \nu_F^* L_{|F}$ is not pseudo-effective. Apply [22, Lemma 2.5] to conclude that $(F, \mathcal{L}_{|F}) \cong (\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(1))$. Consider the family of leaves of \mathcal{F} as in Paragraph 2.11:

$$Z \xrightarrow{\nu} X$$

$$\psi \downarrow \qquad \qquad Y.$$

By [2, Proposition 4.10], we have

$$(Z, \nu^* \mathcal{L}) \cong (\mathbb{P}_Y(\mathcal{E}), \mathcal{O}_{\mathbb{P}_Y(\mathcal{E})}(1)),$$

where $\mathcal{E} := \psi_* \nu^* \mathcal{L}$.

Suppose that the exceptional set $\operatorname{Exc}(\nu)$ dominates Y under ψ . Consider a resolution of singularities $\mu\colon \widehat{Y} \to Y$, and set $\widehat{Z} = \mathbb{P}_{\widehat{Y}}(\mu^*\mathcal{E}) \cong \widehat{Y} \times_Y Z$ with natural morphisms $\widehat{\psi}\colon \widehat{Z} \to \widehat{Y}$ and $\widehat{\nu}\colon \widehat{Z} \to X$. The exceptional set $\operatorname{Exc}(\widehat{\nu})$ also dominates \widehat{Y} under $\widehat{\psi}$. Note that $\operatorname{Exc}(\widehat{\nu})$ has pure codimension one since X is \mathbb{Q} -factorial. Let \widehat{E} be an irreducible component of $\operatorname{Exc}(\widehat{\nu})$ dominating \widehat{Y} . Then

$$\mathcal{O}_{\widehat{Z}}(-\widehat{E}) \cong \mathcal{O}_{\mathbb{P}_{\widehat{Y}}(\mu^*\mathcal{E})}(-k) \otimes \widehat{\psi}^*\mathcal{J}$$

for some positive integer k, and some line bundle \mathcal{J} on \widehat{Y} . In particular, $h^0(\widehat{Y}, S^k \mu^* \mathcal{E} \otimes \mathcal{J}^*) \geq 1$. Let $\widehat{\xi}$ be a divisor on \widehat{Z} such that

$$\mathcal{O}_{\widehat{Z}}(\widehat{\xi}) \cong \mathcal{O}_{\mathbb{P}_{\widehat{Y}}(\mu^*\mathcal{E})}(1).$$

Notice that $\mathcal{O}_{\mathbb{P}_{\widehat{Y}}(\mu^*\mathcal{E})}(1)$ is the pull-back of $\mathcal{O}_{\mathbb{P}_Y(\mathcal{E})}(1) \cong \nu^*\mathcal{L}$ under the natural morphism $\mathbb{P}_{\widehat{Y}}(\mu^*\mathcal{E}) \to \mathbb{P}_Y(\mathcal{E}) \cong Z$, and hence it is semi-ample. From Lemma 3.2 below, we conclude that $m_0(K_{\widehat{Z}/\widehat{Y}} + r\widehat{\xi} + s\widehat{E})$ is effective for some positive integer m_0 and every sufficiently large integer s. This in turn implies that

$$\widehat{\nu}_*(K_{\widehat{Z}/\widehat{Y}} + r\widehat{\xi} + s\widehat{E}) = K_{\mathcal{F}} + rL$$

is pseudo-effective. Thus $\mu_{\alpha}(\mathcal{F} \otimes \mathcal{L}^*) \leq 0$, yielding a contradiction. This proves that the map $\pi\colon X \dashrightarrow Y$ induced by ψ is almost proper, and so \mathcal{F} is induced by a generic \mathbb{P}^r -bundle structure on X.

Step 2. We prove statement (1).

Let $0 = \mathcal{T}_0 \subsetneq \mathcal{T}_1 \subsetneq \cdots \subsetneq \mathcal{T}_k = T_X$ be the Harder–Narasimhan filtration of T_X with respect to α , and set $\mathcal{Q}_i = \mathcal{T}_i/\mathcal{T}_{i-1}$. By Corollary 2.20, \mathcal{T}_1 is an algebraically integrable foliation on X. Moreover, $\mu_{\alpha}(\mathcal{T}_1 \otimes \mathcal{L}^*) > 0$. From Step 1 applied to \mathcal{T}_1 , we conclude that \mathcal{T}_1 is induced by a generic \mathbb{P}^{r_1} -bundle structure $\pi_1 \colon X \dashrightarrow Y_1$ on X and \mathcal{L} restricts to $\mathcal{O}_{\mathbb{P}^{r_1}}(1)$ on a general fiber of π_1 .

Suppose that

$$\mu_{\alpha}^{\max}((T_X/\mathcal{T}_1)\otimes\mathcal{L}^*)=\mu_{\alpha}(\mathcal{Q}_2\otimes\mathcal{L}^*)>0.$$

By Corollary 2.20, \mathcal{T}_2 is an algebraically integrable foliation on X. By Step 1, \mathcal{T}_2 is induced by a generic \mathbb{P}^{r_2} -bundle structure on X. This is impossible since \mathbb{P}^{r_2} cannot admit a generic \mathbb{P}^{r_1} -bundle structure with $r_1 < r_2$. Thus $\mu_{\alpha}(\mathcal{Q}_2 \otimes \mathcal{L}^*) \leq 0$, proving (1).

Step 3. Suppose that \mathcal{F} is a foliation on X, and $\mu_{\alpha}(\mathcal{F} \otimes \mathcal{L}^*) > 0$ for some movable curve class $\alpha \in \text{Mov}(X)$. Let $\mathcal{F}_1 \subset \mathcal{F}$ be the maximal destabilizing subsheaf of \mathcal{F} with respect to α . By Corollary 2.20, \mathcal{F}_1 is an algebraically integrable foliation on X. Moreover, we have $\mu_{\alpha}(\mathcal{F}_1 \otimes \mathcal{L}^*) > 0$. By Step 1, \mathcal{F}_1 is induced by generic \mathbb{P}^{s_1} -bundle structure on X. Let $\mathcal{T}_1 \subset \mathcal{T}_X$ be as in Step 2. By [17, Corollary 2.17], $\mathcal{F}_1 \subset \mathcal{T}_1$ since

$$\mu_{\alpha}^{\min}(\mathcal{F}_1 \otimes \mathcal{L}^*) = \mu_{\alpha}(\mathcal{F}_1 \otimes \mathcal{L}^*) > 0 \quad \text{and} \quad \mu_{\alpha}^{\max}((T_X/T_1) \otimes \mathcal{L}^*) \leq 0$$

by Step 2. Hence, we must have $\mathcal{F}_1 = \mathcal{T}_1$.

If \mathcal{F} is algebraically integrable, then from Step 1, we conclude that $\mathcal{F}_1 = \mathcal{T}_1 = \mathcal{F}$.

Recall that a vector bundle \mathcal{E} on a normal projective variety Y is said to be *semi-ample* if the line bundle $\mathcal{O}_{\mathbb{P}_Y(\mathcal{E})}(m)$ on $\mathbb{P}_Y(\mathcal{E})$ is generated by its global sections for some positive integer m.

Lemma 3.2. Let Y be a normal projective variety, \mathcal{M} a line bundle on Y, and \mathcal{E} a semi-ample vector bundle on Y. Suppose that $h^0(Y, S^k \mathcal{E} \otimes \mathcal{M}) \geq 1$ for some integer $k \geq 1$. Then there exists an integer $m_0 \geq 1$ such that

$$h^0(Y, S^{m_0(sk-1)}\mathcal{E} \otimes \det(\mathcal{E})^{\otimes m_0} \otimes \mathcal{M}^{\otimes m_0s}) \geq 1$$

for every sufficiently large integer s.

Proof. Let \mathcal{E} be a semi-ample vector bundle on Y. By [16, Corollary 1], the vector bundle $\mathcal{E}^* \otimes \det(\mathcal{E})$ is semi-ample. Let m_0 be a positive integer such that the line bundle $\mathcal{O}_Z(m_0)$ on $Z := \mathbb{P}_Y(\mathcal{E}^* \otimes \det(\mathcal{E}))$ is generated by its global sections. Set

$$V := H^0(Y, S^{m_0}(\mathcal{E}^* \otimes \det(\mathcal{E}))) \cong H^0(Z, \mathcal{O}_Z(m_0)),$$

and consider the exact sequence

$$0 \to \mathcal{K} \otimes \mathcal{O}_{Z}(m) \to V \otimes \mathcal{O}_{Z}(m) \to \mathcal{O}_{Z}(m_{0}+m) \to 0.$$

Pick a point y on Y, and denote by Z_y the fiber over y of the natural morphism $Z \to Y$. Let m_1 be a positive integer such that $h^1(Z_y, \mathcal{K}_{|Z_y} \otimes \mathcal{O}_{Z_y}(m)) = 0$ for $m \ge m_1$. Then the map

$$V \otimes H^0(Z_y, \mathcal{O}_{Z_y}(m)) \to H^0(Z_y, \mathcal{O}_{Z_y}(m_0 + m))$$

is surjective for every $m \ge m_1$, and thus the morphism

$$V \otimes S^{m}(\mathcal{E}^{*} \otimes \det(\mathcal{E})) \to S^{m_{0}+m}(\mathcal{E}^{*} \otimes \det(\mathcal{E}))$$

is generically surjective. This yields, for every $m \ge m_1$, an injective map of sheaves

$$S^{m_0+m}(\mathcal{E}) \hookrightarrow V^* \otimes S^m(\mathcal{E}) \otimes \det(\mathcal{E})^{\otimes m_0}. \tag{3.1}$$

Let \mathcal{M} be a line bundle on Y, and k a positive integer such that $h^0(Y, S^k \mathcal{E} \otimes \mathcal{M}) \geq 1$. Let s be an integer such that $m := m_0(sk-1) \geq m_1$. The m_0s -th power of a nonzero global section of $S^k \mathcal{E} \otimes \mathcal{M}$ is a nonzero global section of $S^{m_0sk} \mathcal{E} \otimes \mathcal{M}^{\otimes m_0s}$. Then (3.1) yields

$$1 \leq h^{0}(Y, S^{m_{0}sk} \mathcal{E} \otimes \mathcal{M}^{\otimes m_{0}s})$$

$$\leq \dim V \cdot h^{0}(Y, S^{m_{0}(sk-1)} \mathcal{E} \otimes \det(\mathcal{E})^{\otimes m_{0}} \otimes \mathcal{M}^{\otimes m_{0}s}),$$

completing the proof of the lemma.

Proof of Corollary 1.3. Suppose that there is a positive integer m, an ample line bundle \mathcal{L} , a torsion-free quotient $(\Omega_X^1 \otimes \mathcal{L})^{\otimes m} \twoheadrightarrow \mathcal{Q}$ of positive rank, and a movable class $\alpha \in \operatorname{Mov}(X)$ such that $\mu_{\alpha}(\mathcal{Q}) < 0$. Then $\mu_{\alpha}^{\min}((\Omega_X^1 \otimes \mathcal{L})^{\otimes m}) < 0$. This implies that $\mu_{\alpha}^{\max}((T_X \otimes \mathcal{L}^*)^{\otimes m}) > 0$. Thus, by [17, Theorem 4.2] (see also [9, Proposition 6.1]), $\mu_{\alpha}^{\max}(T_X \otimes \mathcal{L}^*) > 0$. Theorem 3.1 then implies that X is a generic \mathbb{P}^r -bundle for some positive integer r.

4. Bounding the algebraic rank

We start this section by justifying the definition of generalised index in Definition 1.4.

Lemma 4.1. Let X be a complex projective manifold, and D a big divisor on X. Set

$$\hat{\iota}(D) := \sup \{ t \in \mathbb{R} \mid D \equiv tA + E \text{ where } A \text{ is an ample divisor}$$

and $E \text{ is a pseudo-effective } \mathbb{R}\text{-divisor} \}.$

Then there exists an ample divisor A_0 and a pseudo-effective \mathbb{R} -divisor E_0 on X such that $D \equiv \hat{\iota}(D)A_0 + E_0$.

Proof. Set $t_0 = \frac{\widehat{\iota}(D)}{2} < \infty$. Pick a real number $t \ge t_0$, an ample divisor A_t and a pseudo-effective \mathbb{R} -divisor E_t on X such that $D \equiv tA_t + E_t$. Then $\frac{1}{t_0}D - A_t = (\frac{t}{t_0} - 1)A_t + \frac{1}{t_0}E_t$ is pseudo-effective. In order to prove the lemma, it is enough to show that there are finitely many classes of integral effective divisors B on X such that $\frac{1}{t_0}D - B$ is pseudo-effective. Let C_1, \ldots, C_m be movable curves on X such that $[C_1], \ldots, [C_m]$ is a basis of $N_1(X)_{\mathbb{R}}$. If B is an effective divisor on X such that $\frac{1}{t_0}D - B$ is pseudo-effective, then we have $0 \le B \cdot C_i \le \frac{1}{t_0}D \cdot C_i$. These inequalities define a compact set $\Delta \subset N^1(X)_{\mathbb{R}}$. Since the set of classes of effective divisors is discrete in $N^1(X)_{\mathbb{R}}$, the compact set Δ contains at most finitely many of these classes. □

Next we show that the algebraic rank of a foliation is bounded from below by its generalised index, and classify foliations attaining this bound (Theorem 1.5). We fix the notation to be used in what follows.

Notation 4.2. Let $\mathcal{F} \subseteq T_X$ be a foliation with big anti-canonical class and generalised index $\hat{\iota}$. Denote by \mathcal{F}^a the algebraic part of \mathcal{F} , and by r^a its algebraic rank. By Lemma 4.1, there is an ample divisor L and a pseudo-effective \mathbb{R} -divisor E such that $-K_{\mathcal{F}} \equiv \hat{\iota}L + E$. Set $\mathcal{L} := \mathcal{O}_X(L)$, and $\alpha_0 = [\mathcal{L}^{n-1}] \in \text{Mov}(X)$. For any movable curve class $\alpha \in \text{Mov}(X)$, we denote by \mathcal{F}^+_{α} the positive part of \mathcal{F} with respect to α , as defined in Paragraph 2.16, and by $r_{\alpha} = \text{rank}(\mathcal{F}^+_{\alpha})$ its rank.

Proof of Theorem 1.5. We follow Notation 4.2, and assume that $\mathcal{F} \neq T_X$. For any movable curve class $\alpha \in \text{Mov}(X)$ such that $\mu_{\alpha}(\mathcal{F}_{\alpha}^+ \otimes \mathcal{L}^*) \leq 0$, we have

$$\widehat{\iota}\mu_{\alpha}(\mathcal{L}) \leq \det(\mathcal{F}) \cdot \alpha$$

$$\leq \det(\mathcal{F}_{\alpha}^{+}) \cdot \alpha \quad (\text{since } \mu_{\alpha}(\mathcal{F}/\mathcal{F}_{\alpha}^{+}) \leq 0)$$

$$\leq r_{\alpha}\mu_{\alpha}(\mathcal{L}) \quad (\text{since } \mu_{\alpha}(\mathcal{F}_{\alpha}^{+} \otimes \mathcal{L}^{*}) \leq 0)$$

$$\leq r^{a}\mu_{\alpha}(\mathcal{L}) \quad (\text{since } \mathcal{F}_{\alpha}^{+} \subset \mathcal{F}^{a} \text{ by Theorem 2.19}).$$

$$(4.1)$$

By Theorem 1.1, we have $\mu_{\alpha_0}(\mathcal{F}_{\alpha_0}^+ \otimes \mathcal{L}^*) \leq 0$ for the ample class α_0 , and (4.1) gives that $\hat{\iota} \leq r^a$.

Suppose from now on that $\hat{\iota} = r^a$. Then (4.1) shows that, for any movable curve class $\alpha \in \text{Mov}(X)$ such that $\mu_{\alpha}(\mathcal{F}_{\alpha}^{+} \otimes \mathcal{L}^{*}) \leq 0$ (in particular for $\alpha = \alpha_{0}$), we have $\mathcal{F}^{a} = \mathcal{F}_{\alpha}^{+}$, and $\mu_{\alpha}(\mathcal{F}^{a} \otimes \mathcal{L}^{*}) = 0$. Moreover $E \cdot \alpha_{0} = 0$, and thus E = 0 and $-K_{\mathcal{F}} \equiv \hat{\iota}L$.

Next we show that $\mu_{\alpha}(\mathcal{F}^a \otimes \mathcal{L}^*) = 0$ for every $\alpha \in \operatorname{Mov}(X)$. Suppose that this is not the case. Since the ample class α_0 lies in the interior of $\operatorname{Mov}(X)$ and $\mu_{\alpha_0}(\mathcal{F}^a \otimes \mathcal{L}^*) = 0$, there must exist a class $\alpha \in \operatorname{Mov}(X)$ such that $\mu_{\alpha}(\mathcal{F}^a \otimes \mathcal{L}^*) > 0$. By Theorem 3.1, there is a dense open subset $X^{\circ} \subset X$ and a \mathbb{P}^{r^a} -bundle structure $\pi^{\circ} \colon X^{\circ} \to Y^{\circ}$ such that $\mathcal{F}^a_{|X^{\circ}} = T_{X^{\circ}/Y^{\circ}}$, and \mathcal{L} restricts to $\mathcal{O}_{\mathbb{P}^{r^a}}(1)$ on any fiber of π° . By Remark 2.7, there is a foliation \mathcal{G} on Y° such that $\mathcal{F}_{|X^{\circ}} = (\pi^{\circ})^{-1}\mathcal{G}$. By shrinking Y° if necessary, we may assume that \mathcal{G} is locally free, and $\mathcal{F}/T_{X^{\circ}/Y^{\circ}} \cong \phi^*\mathcal{G}$. Therefore, for a general fiber F of π° , we have

$$\mathcal{O}_{\mathbb{P}^{r_a}}(\widehat{\iota}) \cong \det(\mathcal{F})_{|F} \cong \det(T_{X^{\circ}/Y^{\circ}})_{|F} \cong \mathcal{O}_{\mathbb{P}^{r_a}}(r^a+1),$$

contradicting our assumption that $\hat{\iota} = r^a$. We conclude that $\mu_{\alpha}(\mathcal{F}^a \otimes \mathcal{L}^*) = 0$ for every $\alpha \in \text{Mov}(X)$.

This implies that $-K_{\mathcal{F}^a} \equiv r^a L$, and hence \mathcal{F}^a is an algebraically integrable Fano foliation. Let (F, Δ_F) be a general log leaf of \mathcal{F}^a , so that $-(K_F + \Delta_F) \equiv r^a L_{|F}$. By Corollary 2.13, $\Delta_F \neq 0$, and thus $K_F + r^a L_{|F} \equiv -\Delta_F$ is not pseudo-effective. By [22, Lemma 2.5],

$$(F, \mathcal{L}, \mathcal{O}_F(\Delta_F)) \cong (\mathbb{P}^{r^a}, \mathcal{O}_{\mathbb{P}^{r^a}}(1), \mathcal{O}_{\mathbb{P}^{r^a}}(1)).$$

In particular, (F, Δ_F) is log canonical. By [1, Proposition 5.3], there is a common point x in the closure of a general leaf of \mathcal{F}^a . This implies that X is covered by rational curves passing through x having degree 1 with respect to \mathcal{L} . It follows that X is a Fano manifold with Picard number $\rho(X) = 1$, and hence $-K_{\mathcal{F}^a} \equiv r^a L$ implies that $-K_{\mathcal{F}^a} \sim r^a L$. By [5, Theorem 1.1], $X \cong \mathbb{P}^n$ and \mathcal{F}^a is induced by a linear projection $\pi \colon \mathbb{P}^n \dashrightarrow \mathbb{P}^{n-r_a}$. By Remark 2.7, there is a foliation \mathcal{G} on \mathbb{P}^{n-r_a} such that $\mathcal{F} = \pi^{-1}\mathcal{G}$. By (2.1) and (2.2), outside the center of the linear projection $\pi \colon \mathbb{P}^n \dashrightarrow \mathbb{P}^{n-r_a}$, we have $\pi^*K_{\mathcal{G}} \sim K_{\mathcal{F}} - K_{\mathcal{F}^a} \sim 0$, and thus $K_{\mathcal{G}} \sim 0$.

Finally, we describe foliations \mathcal{F} whose algebraic rank slightly exceeds the generalised index, namely $\hat{\iota}(\mathcal{F}) < r^a(\mathcal{F}) \leqslant \hat{\iota}(\mathcal{F}) + 1$. We start by addressing foliations \mathcal{F} with $K_{\mathcal{F}} \equiv 0$ and algebraic rank $r^a = 1$.

Lemma 4.3. Let X be a complex projective manifold with Picard number $\rho(X) = 1$, and let \mathcal{F} be a foliation on X. Suppose that $K_{\mathcal{F}} \equiv 0$ and that \mathcal{F} has algebraic rank 1. Then the closure of a general leaf of the algebraic part of \mathcal{F} is a rational curve.

Proof. There exist a normal projective variety Y, a dominant rational map $\pi: X \dashrightarrow Y$ of relative dimension 1, and a purely transcendental foliation \mathcal{G} on Y such that $\mathcal{F} = \pi^{-1}\mathcal{G}$. Denote by \mathcal{C} the foliation induced by π , i.e. the algebraic part of \mathcal{F} .

After replacing Y with a birationally equivalent variety, we may assume that Y is the family of leaves of \mathcal{C} . Let $X^{\circ} \subset X$ be an open subset with complement of codimension at least 2 such that π restricts to an equidimensional morphism $\pi_{|X^{\circ}}\colon X^{\circ} \to Y$. By (2.1) and (2.2) applied to $\pi_{|X^{\circ}}$, there is an effective divisor R on X such that, on X° , we have

$$-K_{\mathcal{C}} \equiv \pi^* K_{\mathcal{G}} + R.$$

By Theorem 2.19, $K_{\mathcal{C}}$ is pseudo-effective, and hence so is $-K_{\mathcal{C}}$. If $K_{\mathcal{C}} \not\equiv 0$, then the lemma follows from Theorem 2.19. Suppose from now on that $K_{\mathcal{C}} \equiv 0$, and let (F, Δ_F) be a general log leaf of \mathcal{C} , so that $K_F + \Delta_F \equiv 0$. By Corollary 2.13, $\Delta_F \neq 0$. Thus $\deg(K_F) < 0$, and hence $F \cong \mathbb{P}^1$.

Proof of Theorem 1.8. We follow Notation 4.2, and assume that $\hat{\iota} < r^a \le \hat{\iota} + 1$. We will show that the closure of a general leaf of \mathcal{F}^a is rationally connected.

Step 1. Let $\alpha \in \text{Mov}(X)$ be a movable curve class. If $\mu_{\alpha}(\mathcal{F}_{\alpha}^{+} \otimes \mathcal{L}^{*}) < 0$, then (4.1) gives that $r^{a} - 1 \leq \hat{\iota} < r_{\alpha} \leq r^{a}$. Hence, $\mathcal{F}_{\alpha}^{+} = \mathcal{F}^{a}$. By Theorem 2.19, the closure of a general leaf of \mathcal{F}^{a} is rationally connected. If $\mu_{\alpha}(\mathcal{F}_{\alpha}^{+} \otimes \mathcal{L}^{*}) = 0$, then (4.1) gives that

$$r^a - 1 \le \hat{\iota} \le r_\alpha \le r^a$$
,

and hence \mathcal{F}_{α}^{+} has codimension at most 1 in \mathcal{F}^{a} . If $\mathcal{F}_{\alpha}^{+} = \mathcal{F}^{a}$, then, as before, the closure of a general leaf of \mathcal{F}^{a} is rationally connected. Suppose that \mathcal{F}_{α}^{+} has codimension 1 in \mathcal{F}^{a} . Then $\hat{\imath} = r^{a} - 1$. Let $\beta \in \operatorname{Mov}(X)$ be another movable curve class. If $\mathcal{F}_{\beta}^{+} \not\subset \mathcal{F}_{\alpha}^{+}$, then $\mathcal{F}_{\alpha}^{+} + \mathcal{F}_{\beta}^{+} = \mathcal{F}^{a}$ and Theorem 2.19 applied to both α and β gives that the closure of a general leaf of \mathcal{F}^{a} is rationally connected.

Therefore, from now on we may assume the following.

Additional assumption. For any $\alpha \in \text{Mov}(X)$, $\mu_{\alpha}(\mathcal{F}_{\alpha}^{+} \otimes \mathcal{L}^{*}) \geq 0$. If $\alpha \in \text{Mov}(X)$ is such that $\mu_{\alpha}(\mathcal{F}_{\alpha}^{+} \otimes \mathcal{L}^{*}) = 0$, then \mathcal{F}_{α}^{+} has codimension 1 in \mathcal{F}^{a} , and $\mathcal{F}_{\beta}^{+} \subset \mathcal{F}_{\alpha}^{+}$ for any $\beta \in \text{Mov}(X)$.

By Theorem 1.1, we have $\mu_{\alpha_0}(\mathcal{F}_{\alpha_0}^+ \otimes \mathcal{L}^*) \leq 0$ for the ample class α_0 . The additional assumption and (4.1) for $\alpha = \alpha_0$ give that $r_{\alpha_0} = r^a - 1 = \hat{\iota}$, E = 0 and $-K_{\mathcal{F}} \equiv \hat{\iota}L$.

Step 2. Suppose that $\mu_{\alpha}(\mathcal{F}_{\alpha}^{+}\otimes\mathcal{L}^{*})=0$ for every α in a nonempty open subset U of $\operatorname{Mov}(X)$. From the additional assumption, it follows that $\mathcal{F}_{\alpha}^{+}=\mathcal{F}_{\alpha_{0}}^{+}$ and $\mu_{\alpha}(\mathcal{F}_{\alpha_{0}}^{+}\otimes\mathcal{L}^{*})=0$ for every $\alpha\in U$. Thus

$$-K_{\mathcal{F}_{\alpha_0}}\equiv r_{\alpha_0}L.$$

By Theorem 1.5, $X \cong \mathbb{P}^n$, and $\mathcal{F}_{\alpha_0}^+$ is induced by a linear projection $\pi: \mathbb{P}^n \longrightarrow \mathbb{P}^{n-r_{\alpha_0}}$. Since $X \cong \mathbb{P}^n$, we must have

$$-K_{\mathcal{F}} \sim \hat{\iota}L \sim -K_{\mathcal{F}_{\alpha_0}}.$$

By Remark 2.7, there is a foliation \mathcal{G} on $\mathbb{P}^{n-r_{\alpha_0}}$ such that $\mathcal{F}=\pi^{-1}\mathcal{G}$, and $K_{\mathcal{G}}\sim 0$. Moreover, since $\mathcal{F}_{\alpha_0}^+$ has codimension 1 in \mathcal{F}^a , \mathcal{G} has algebraic rank 1. By Lemma 4.3, the closure of a general leaf of the algebraic part of \mathcal{G} is a rational curve, and thus the closure of a general leaf of \mathcal{F}^a is rationally connected.

Step 3. By Step 2, we may assume that $\mu_{\alpha}(\mathcal{F}_{\alpha}^{+} \otimes \mathcal{L}^{*}) > 0$ for some $\alpha \in \text{Mov}(X)$. By Theorem 3.1, \mathcal{F}_{α}^{+} is induced by a generic $\mathbb{P}^{r_{\alpha}}$ -bundle structure $\pi_{\alpha} \colon X \dashrightarrow Y_{\alpha}$ on X, and \mathcal{L} restricts to $\mathcal{O}_{\mathbb{P}^{r_{\alpha}}}(1)$ on general fibers of π_{α} . If necessary, we can replace α by the class of a line on a fiber of π_{α} . As in (4.1), we have

$$(r^a - 1) = \hat{\iota} = \det(\mathcal{F}) \cdot \alpha$$

 $\leq \det(\mathcal{F}_{\alpha}^+) \cdot \alpha$
 $= (r_{\alpha} + 1),$

and thus $r_{\alpha} \geq r^a - 2$. Moreover, by Corollary 1.7, $-K_{\mathcal{F}_{\alpha}^+} - r_{\alpha}L$ is not pseudoeffective. Thus, there exists a nonempty open subset V of Mov(X) such that, for every $\beta \in V$, $\mu_{\beta}(\mathcal{F}_{\alpha}^+ \otimes \mathcal{L}^*) < 0$. The additional assumption implies in particular that $\mathcal{F}_{\beta}^+ \neq \mathcal{F}_{\alpha}^+$ for every $\beta \in V$.

By Step 2, we may assume that $\mu_{\beta}(\mathcal{F}_{\beta}^{+}\otimes\mathcal{L}^{*})>0$ for some $\beta\in V$. As before, \mathcal{F}_{β}^{+} is induced by a generic $\mathbb{P}^{r_{\beta}}$ -bundle structure $\pi_{\beta}\colon X\dashrightarrow Y_{\beta}$ on X, \mathcal{L} restricts to $\mathcal{O}_{\mathbb{P}^{r_{\beta}}}(1)$ on general fibers of π_{β} , and $r_{\beta}\geqslant r^{a}-2$. Notice that $\mathcal{F}_{\alpha}^{+}\cap\mathcal{F}_{\beta}^{+}=0$ and $\mathcal{F}_{\alpha}^{+}+\mathcal{F}_{\beta}^{+}\subset\mathcal{F}^{a}$. If $\mathcal{F}_{\alpha}^{+}+\mathcal{F}_{\beta}^{+}=\mathcal{F}^{a}$, then we conclude as before that the closure of a general leaf of \mathcal{F}^{a} is rationally connected. So we may assume that $\mathcal{F}_{\alpha}^{+}+\mathcal{F}_{\beta}^{+}\subsetneq\mathcal{F}^{a}$. This can only happen if $(r_{\alpha},r_{\beta},r^{a})=(1,1,3)$.

Let H_{α} and H_{β} be the dominating unsplit families of rational curves on X whose general members correspond to lines on fibers of π_{α} and π_{β} , respectively. Denote by $\pi^{\circ}: X^{\circ} \to T^{\circ}$ the (H_{α}, H_{β}) -rationally connected quotient of X (see for instance [1, 6.4] for this notion). By [5, Lemma 2.2], we may assume that codim $X \setminus X^{\circ} \ge 2$, T° is smooth, and π° has irreducible and reduced fibers. Applying [19, Corollaires 14.4.4 et 15.2.3], we see that π° is flat. This in turn implies that its fibers are Cohen–Macaulay since both X and T° are smooth. By [1, Lemma 6.9], there is an

inclusion $T_{X^{\circ}/T^{\circ}} \subset \mathcal{F}^{a}_{|X^{\circ}}$. If $T_{X^{\circ}/T^{\circ}} = \mathcal{F}^{a}_{|X^{\circ}}$, then the closure of a general leaf of \mathcal{F}^{a} is rationally connected. So we may assume that dim $T^{\circ} = \dim X^{\circ} - 2$. By Remark 2.7, there is a foliation \mathcal{G}° on T° such that $\mathcal{F}_{|X^{\circ}} = \pi^{\circ -1} \mathcal{G}^{\circ}$. By (2.1), we have

$$-K_{X^{\circ}/T^{\circ}} = -K_{\mathcal{F}|X^{\circ}} + \pi^{\circ *}K_{\mathcal{G}^{\circ}}.$$

Since codim $X \setminus X^{\circ} \ge 2$, a general complete intersection curve in X is contained in $X \setminus X^{\circ}$. Let $C \to T^{\circ}$ be the normalization of a complete curve passing through a general point, denote by X_C the fiber product $C \times_{T^{\circ}} X^{\circ}$, and by $\pi_C \colon X_C \to C$ the natural morphism. By [18, Corollaire 5.12.4], X_C satisfies Serre's condition S_2 . On the other hand, X_C is smooth in codimension 1 since the fibers of π_C are reduced. It follows that X_C is a normal variety by Serre's criterion for normality. We have

$$-K_{X_C/C} = -K_{\mathcal{F}|X_C} + \pi_C^* K_{\mathcal{G}^{\circ}|C}.$$

If $K_{\mathcal{G}^{\circ}} \cdot C \ge 0$, then $-K_{X_C/C}$ is ample, contradicting [5, Theorem 3.1]. Hence, $K_{\mathcal{G}^{\circ}} \cdot C < 0$, and [1, Proposition 7.5] implies that the closure of a general leaf of \mathcal{F} is rationally connected.

Question 4.4. Is there a foliation \mathcal{F} with $\widehat{\iota}(\mathcal{F}) \notin \mathbb{N}$ and $\widehat{\iota}(\mathcal{F}) < r^a(\mathcal{F}) < \widehat{\iota}(\mathcal{F}) + 1$? From the proof of Theorem 1.8, we see that in this case we must have $\mathcal{F}_{\alpha}^+ = \mathcal{F}^a$ for every $\alpha \in \text{Mov}(X)$ such that $\mu_{\alpha}(\mathcal{F}_{\alpha}^+ \otimes \mathcal{L}^*) \leq 0$.

References

- [1] C. Araujo and S. Druel, On Fano foliations, *Adv. Math.*, **238** (2013), 70–118. Zbl 1282.14085 MR 3033631
- [2] C. Araujo and S. Druel, On codimension 1 del Pezzo foliations on varieties with mild singularities, *Math. Ann.*, **360** (2014), no. 3-4, 769–798. Zbl 1396.14035 MR 3273645
- [3] C. Araujo and S. Druel, On Fano foliations. II, in *Foliation theory in algebraic geometry*. *Proceedings of the conference, (New York, NY, September 3–7, 2013)*, 1–20, Springer, Cham, 2016. Zbl 1382.14012 MR 3644241
- [4] C. Araujo and S. Druel, Codimension 1 Mukai foliations on complex projective manifolds, *J. Reine Angew. Math.*, **727** (2017), 191–246. Zbl 1395.32016 MR 3652251
- [5] C. Araujo, S. Druel, and S. J. Kovács, Cohomological characterizations of projective spaces and hyperquadrics, *Invent. Math.*, 174 (2008), no. 2, 233–253. Zbl 1162.14037 MR 2439607
- [6] F. Bogomolov and M. McQuillan, Rational curves on foliated varieties, in *Foliation theory in algebraic geometry*, 21–51, Simons Symp., Springer, Cham, 2016. Zbl 1337.14041 MR 3644242
- [7] J.-B. Bost, Algebraic leaves of algebraic foliations over number fields, *Publ. Math. Inst. Hautes Études Sci.*, (2001), no. 93, 161–221. Zbl 1034.14010 MR 1863738

- [8] S. Boucksom, J.-P. Demailly, M. Păun, and T. Peternell, The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension, J. Algebraic Geom., 22 (2013), no. 2, 201–248. Zbl 1267.32017 MR 3019449
- [9] F. Campana and T. Peternell, Geometric stability of the cotangent bundle and the universal cover of a projective manifold. With an appendix by Matei Toma, *Bull. Soc. Math. France*, **139** (2011), no. 1, 41–74. Zbl 1218.14030 MR 2815027
- [10] F. Campana and M. Păun, Foliations with positive slopes and birational stability of orbifold cotangent bundles, *Publ. Math. Inst. Hautes Études Sci.*, 129 (2019), 1–49. Zbl 1423.14109 MR 3949026
- [11] F. Campana and M. Păun, Orbifold generic semi-positivity: an application to families of canonically polarized manifolds, *Ann. Inst. Fourier (Grenoble)*, **65** (2015), no. 2, 835–861. Zbl 1338.14012 MR 3449168
- [12] D. Cerveau and A. Lins Neto, Irreducible components of the space of holomorphic foliations of degree two in CP(n), $n \ge 3$, Ann. of Math. (2), **143** (1996), no. 3, 577–612. Zbl 0855.32015 MR 1394970
- [13] S. Druel, Caractérisation de l'espace projectif, *Manuscripta Math.*, **115** (2004), no. 1, 19–30. Zbl 1070.14012 MR 2092774
- [14] S. Druel, A decomposition theorem for singular spaces with trivial canonical class of dimension at most five, *Invent. Math.*, 211 (2018), no. 1, 245–296. Zbl 1419.14063 MR 3742759
- [15] S. Druel, On foliations with nef anti-canonical bundle, *Trans. Amer. Math. Soc.*, **369** (2017), no. 11, 7765–7787. Zbl 1388.14056 MR 3695844
- [16] T. Fujiwara, Varieties of small Kodaira dimension whose cotangent bundles are semiample, *Compositio Math.*, **84** (1992), no. 1, 43–52. Zbl 0763.14015 MR 1183561
- [17] D. Greb, S. Kebekus, and T. Peternell, Movable curves and semistable sheaves, *Internat. Math. Res. Notices*, (2016), no. 2, 536–570. Zbl 1342.14022 MR 3493425
- [18] A. Grothendieck, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II, *Inst. Hautes Études Sci. Publ. Math.*, (1965), no. 24, 231pp. Zbl 0135.39701 MR 199181
- [19] A. Grothendieck, Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III, *Inst. Hautes Études Sci. Publ. Math.*, (1966), no. 28, 255pp. Zbl 0144.19904 MR 217086
- [20] R. Hartshorne, Stable reflexive sheaves, Math. Ann., 254 (1980), no. 2, 121–176.
 Zbl 0431.14004 MR 597077
- [21] A. Höring, On a conjecture of Beltrametti and Sommese, *J. Algebraic Geom.*, **21** (2012), no. 4, 721–751. Zbl 1253.14007 MR 2957694
- [22] A. Höring, Twisted cotangent sheaves and a Kobayashi-Ochiai theorem for foliations, *Ann. Inst. Fourier (Grenoble)*, **64** (2014), no. 6, 2465–2480. Zbl 1348.14054 MR 3331171
- [23] S. Kobayashi and T. Ochiai, Characterizations of complex projective spaces and hyperquadrics, *J. Math. Kyoto Univ.*, **13** (1973), 31–47. Zbl 0261.32013 MR 316745
- [24] F. Loray, J. V. Pereira, and F. Touzet, Singular foliations with trivial canonical class, *Invent. Math.*, **213** (2018), no. 3, 1327–1380. Zbl 06933334 MR 3842065

- [25] F. Loray, J. V. Pereira, and F. Touzet, Foliations with trivial canonical bundle on Fano 3-folds, *Math. Nachr.*, **286** (2013), no. 8-9, 921–940. Zbl 1301.37032 MR 3066408
- [26] S. Mori, Projective manifolds with ample tangent bundles, *Ann. of Math.* (2), **110** (1979), no. 3, 593–606. Zbl 0423.14006 MR 554387

Received November 28, 2017

- C. Araujo, IMPA, Estrada Dona Castorina 110, Rio de Janeiro, 22460-320, Brazil E-mail: caraujo@impa.br
- S. Druel, Institut Camille Jordan, Université Lyon, CNRS, Université Claude Bernard, Lyon 1, UMR 5208, 69622 Villeurbanne, France

E-mail: stephane.druel@math.cnrs.fr