Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 94 (2019)

Heft: 4

Artikel: Rigidity of Busemann convex Finsler metrics

Autor: Ivanov, Sergei / Lytchak, Alexander

DOI: https://doi.org/10.5169/seals-847033

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Rigidity of Busemann convex Finsler metrics

Sergei Ivanov and Alexander Lytchak

Abstract. We prove that a Finsler metric is nonpositively curved in the sense of Busemann if and only if it is affinely equivalent to a Riemannian metric of nonpositive sectional curvature. In other terms, such Finsler metrics are precisely Berwald metrics of nonpositive flag curvature. In particular in dimension 2 every such metric is Riemannian or locally isometric to that of a normed plane. In the course of the proof we obtain new characterizations of Berwald metrics in terms of the so-called linear parallel transport.

Mathematics Subject Classification (2010). 53B40, 53C60, 53C23.

Keywords. Non-positive curvature, Busemann convex space, Berwald metric.

1. Introduction

The notion of nonpositive curvature in Riemannian geometry has two famous generalizations to metric geometry due to Alexandrov and Busemann, respectively. Alexandrov's generalization is nowadays known as locally CAT(0) spaces. We refer to [1,4] and the bibliography therein for a vast literature on the subject. Busemann nonpositively curved spaces, also known as *locally convex* spaces, are a larger class of metric spaces defined as follows (see e.g. [16]):

Definition 1.1. A geodesic metric space (X, d) is *Busemann convex* if for every pair of constant-speed geodesics $\gamma_{1,2}$: $[a,b] \to X$ the function $t \mapsto d(\gamma_1(t), \gamma_2(t))$ is convex on [a,b].

A metric space (X, d) is nonpositively curved in the sense of Busemann (Busemann NPC for short) if every point in X has a Busemann convex neighborhood.

In more geometric terms, a geodesic metric space (X, d) is Busemann convex if and only if for every geodesic triangle $\triangle abc$ in X, the distance between the midpoints of its sides [ab] and [ac] is no greater than $\frac{1}{2}d(b,c)$.

Contrary to Alexandrov's definition of nonpositive curvature, Busemann's one is satisfied by all normed vector spaces with strictly convex norms. Thus it can be sensibly applied to Finsler metrics. In fact, Finsler metrics are one of the main motivations in Busemann's work [5] where the definition is introduced. It is natural

to ask how this class of Finsler metrics can be characterized in differential geometric terms. For discussions of this question, see [5, 8, §13], the introduction in [10, 11] and Problem 35 in Z. Shen's problem list [19].

In this paper we settle this question. It turns out that the Busemann NPC condition for (smooth and strictly convex) Finsler metrics has surprising rigidity implications and very few metrics satisfy it.

For Riemannian manifolds the Busemann NPC condition is equivalent to non-positive sectional curvature. Hence one can construct an open set of Riemannian examples by perturbing any negatively curved metric. Indeed, if a perturbed metric tensor is sufficiently C^2 -close to the original one, then it is also negatively curved and hence Busemann NPC. One might expect that in the Finsler case a similar property holds and that one can construct many examples of Busemann NPC Finsler metrics by varying negatively curved Riemannian ones. These expectations turn out to be wrong as our main theorem shows:

Theorem 1. A Finsler manifold (M, F) is Busemann NPC if and only if there exists a nonpositively curved Riemannian metric g on M whose Levi-Civita connection preserves the Finsler norm F.

The Finsler norm F in Theorem 1 is not assumed to be reversible. Strictly speaking, non-reversible Finsler manifolds are not metric spaces since the distance lacks symmetry. Nevertheless Definition 1.1 applies just as well, cf. [11, Section 5].

In the language of Finsler geometry our result reads as follows:

Theorem 2. A Finsler manifold (M, F) is Busemann NPC if and only if it is a Berwald manifold of nonpositive flag curvature.

The definitions of Berwald metrics and flag curvature are given in Section 2. Theorem 2 is essentially a reformulation of Theorem 1, their equivalence easily follows from Szabó's metrization theorem [20]. The "if" direction of Theorem 2 is proved by Kristály, Varga and Kozma [11], see also [9]. Results of [9] also imply that if (M, F) is Berwald and Busemann NPC, then it has nonpositive flag curvature.

We now discuss some implications of Theorems 1 and 2 and well-known features of Berwald metrics. First, Theorem 1 implies that in a connected Busemann NPC Finsler manifold all tangent spaces are isometric as normed vector spaces. This is already a strong restriction on a Finsler metric.

Furthermore, if (M, F) and g are as in Theorem 1 then the Finsler norm at every point is invariant under the holonomy group of (M, g). Hence, if the holonomy group acts transitively on the unit sphere then the Finsler structure must be Riemannian. In dimension 2 the local holonomy group is either transitive or trivial, thus we have the following (cf. [6, Corollary 4.3.5]):

Corollary 1.2. If a 2-dimensional connected Finsler manifold is Busemann NPC, then it is Riemannian or locally isometric to a normed plane.

In general, recall that the Riemannian holonomy group is transitive unless the metric locally splits as a product or is locally symmetric [3]. This leads to a classification of Busemann NPC Finsler metrics exactly as that of Berwald metrics (see [20] or [6, Theorem 4.3.4]) with additional requirements that the symmetric spaces involved are of non-compact type and the Riemannian factors are nonpositively curved.

A Finsler norm F is preserved by the Levi-Civita parallel transport of a Riemannian metric g if and only if F and g are affinely equivalent, that is if they have the same geodesics up to affine reparametrizations (see e.g. [6, Theorem 4.1.3]). Thus we have the following short reformulation of Theorem 1:

Corollary 1.3. A Finsler metric is Busemann NPC if and only if it is affinely equivalent to a nonpositively curved Riemannian metric.

We mention that Finsler manifolds affinely equivalent to nonpositively curved Riemannian symmetric spaces have recently turned up and played a prominent role in a series of papers of B. Leeb, M. Kapovich, and J. Porti, see [7] and references therein.

Characterizations of Berwald spaces. Due to known results mentioned above, in order to prove Theorem 1 it would suffice to show that every Busemann NPC Finsler space is Berwald. However, we prove the "only if" part of Theorem 1 directly, bypassing Szabo's metrization theorem. The proof requires very little background from Finsler geometry, summarized in Section 2.

The proof consists of two steps. In the first step, contained in Section 3, we observe that in a Busemann NPC Finsler space the linear parallel transport along any geodesic preserves the Finsler norm. This follows from analysis of Jacobi fields near points where they vanish.

The above mentioned *linear parallel transport* is defined in Section 2. Note that this notion is different from the more commonly used "canonical parallel transport", which is usually non-linear. See e.g. [6, Chapter 4] where the two parallel transports are considered together. The non-linear parallel transport is not used in this paper beyond the comparison remarks in this introduction.

In the second step, contained in Proposition 4.2 (see Section 4), we prove the following general fact about Finsler metrics: If the linear parallel transport of a Finsler manifold (M, F) preserves F, then F is Berwald. In the proof (see also Proposition 4.1) we construct a Riemannian metric affinely equivalent to F. This Riemannian metric is then used in the proof of Theorem 1 in Section 5.

Remark. We suggest the reader to compare Proposition 4.2 with well-known facts about the canonical (non-linear) parallel transport: It always preserves the Finsler norm, and it is linear if and only if the metric is Berwald. Proposition 4.2 "mirrors" the last mentioned characterization with the linear parallel transport, which is by

definition linear but does not, in general, preserve the norm. Note that for Berwald metrics the two parallel transports coincide, see [6, §4.3].

As a by-product, we obtain another characterization of Berwald spaces in terms of the *linear holonomy group* naturally defined via the linear parallel transport. Namely in Proposition 4.3 we prove the following: *The closure of the linear holonomy group of a Finsler metric is compact if and only if the metric is Berwald.*

We mention that similar questions for the non-linear holonomy group (defined via the non-linear parallel transport) were recently studied in [15].

Beyond smoothness. We work only with smooth Finsler structures and the smoothness is essential in our proofs. It would be interesting to extend some of the results to non-smooth Finsler structures and more general metric spaces. In particular Corollary 1.3 suggests the following questions.

Question 1. Given a Busemann convex metric space, is there a CAT(0) space closely and naturally related to it?

Question 2. Let (X, d) be a geodesic space affinely equivalent to a Busemann convex space (X, d_0) . Is it true that (X, d) must be Busemann convex as well?

In the slightly more general class of spaces with convex bicombings the first question turns out to be of interest in relation with the theory of Gromov hyperbolic groups, see [12]. Some examples showing that one cannot expect affine equivalence, as in Corollary 1.3, are discussed in Section 6.

Metric spaces affinely equivalent to Riemannian manifolds are characterized in [13] in a way similar to Szabó's metrization theorem. In fact, every such space is a limit of smooth Finsler metrics affinely equivalent to the same Riemannian one, see Theorem 1.4 and Lemma 4.1 in [13]. This and Corollary 1.3 imply an affirmative answer to the second question above if (X, d_0) is a smooth Finsler manifold.

Acknowledgements. The authors thank David Bao, Martin Kell, Alexandru Kristaly, Hans-Bert Rademacher, Jozsef Szilasi for helpful comments. Special thanks are due to Urs Lang for a discussion which led to the examples in Section 6. The first author was supported in part by the RFBR grant 17-01-00128-a and by the Presidium of the Russian Academy of Sciences grant PRAS-18-01. The second author was supported in part by the DFG grants SFB TRR 191 and SPP 2026.

2. Preliminaries

In this section we recall some basics in Finsler geometry and prove some auxiliary facts. We follow the presentation in [17], where most concepts are developed from Riemannian point of view. We refer to [6] and [2] for more exhaustive exposition

of Finsler geometry. We include proofs of some standard facts to keep the paper accessible to readers not familiar with Finsler geometry.

Let (M, F) be a Finsler manifold, with the usual smoothness and strict convexity assumptions on the Finsler norm $F: TM \to \mathbb{R}_+$. These assumptions ensure that F determines a smooth geodesic flow on $TM \setminus \{0\}$. As mentioned in the introduction we do not assume that F is reversible. We denote by d_F the (non-symmetric) distance function induced by F on M.

By a *geodesic* we always mean an affine geodesic, i.e., a constant-speed one. Constant paths are not regarded as geodesics. Since all our considerations are local, we may always assume that the manifold M is narrowed down to a small open region where all geodesics are embedded.

For a nonzero $v \in TM$, we denote by γ_v the unique geodesic with initial data $\dot{\gamma}_v(0) = v$. Note that $\gamma_v(t)$ depends smoothly on v and t.

A *Jacobi field* along a geodesic γ is a variation field of a smooth family of geodesics. We denote by $\mathcal{J}^F(\gamma)$ the set of all Jacobi fields along γ .

We make use of several notions from Finsler geometry. They all can be defined by means of osculating Riemannian metrics, see below.

2.1. Osculating Riemannian metrics. For every $p \in M$ and $v \in T_pM \setminus \{0\}$ there is a unique positive definite quadratic form g_v on T_pM such that g_v and $F^2|_{T_pM}$ agree to second order at v. If V is a non-vanishing vector field on an open set $U \subset M$, the family $\{g_{V(p)}\}_{p \in U}$ of quadratic forms defines a Riemannian metric g_V on U. If γ is an embedded geodesic and V extends the velocity field of γ to a neighborhood of γ , we call g_V an osculating Riemannian metric for γ and denote it by g^{γ} . Note that g^{γ} is uniquely determined at every point on γ but the extension to a neighborhood is not unique.

We need the following property (see Lemma 4.4 and Proposition 5.1 in [17] or Chapter 8 in [18]):

Lemma 2.1. Let γ be an embedded geodesic of (M, F) and $g = g^{\gamma}$ an osculating Riemannian metric for γ . Then γ is a geodesic of g with the same space of Jacobi fields: $\mathcal{J}^g(\gamma) = \mathcal{J}^F(\gamma)$.

We comment that Lemma 2.1 is essentially a consequence of the fact that respective equations from calculus of variations involve only first and second derivatives of a Lagrangian with respect to the velocity variable.

- **2.2.** Linear parallel transport and Jacobi operator. An osculating Riemannian metric g^{γ} for a geodesic γ defines the following structures, which are in fact independent on the choice of g^{γ} (see [17] or Lemma 2.3 below):
- The covariant derivative of a vector field W along γ , denoted by $D_{\gamma}W$:

$$D_{\gamma}W(t) = \frac{\nabla^{\gamma}}{dt}W(t) \tag{2.1}$$

where ∇^{γ} is the Levi-Civita connection of g^{γ} .

- The linear parallel transport along γ associated to this connection. It is a family of non-degenerate linear maps between tangent spaces $T_{\gamma(t)}M$.
- The Jacobi operator R^{γ} , which is a family of linear operators on the tangent spaces $T_{\gamma(t)}M$ satisfying the Jacobi equation

$$D_{\gamma}D_{\gamma}J(t) = -R^{\gamma}(J(t)) \tag{2.2}$$

for all Jacobi fields J along γ . Namely $R^{\gamma}(w) = R_{g^{\gamma}}(w, \dot{\gamma}(t))\dot{\gamma}(t)$ for all $w \in T_{\gamma(t)}M$, where $R_{g^{\gamma}}$ is the Riemannian curvature tensor of g^{γ} .

Our notation has γ rather than $\dot{\gamma}$ in indices (cf. [17]) since we are only interested in vector fields along geodesics.

The following simple computation is used several times in the paper.

Lemma 2.2. Let $\gamma = \gamma_v$ be a geodesic of (M, F) and $w \in T_{\gamma(0)}M$. Define a Jacobi field J along γ by

$$J(t) = \frac{d}{ds} \Big|_{s=0} \gamma_{v+sw}(t), \qquad (2.3)$$

and let W be a D_{γ} -parallel vector field along γ (with respect to any osculating Riemannian metric) with W(0) = w. Then

$$J(t) = t \cdot W(t) + O(t^3), \quad t \to 0.$$
 (2.4)

Proof. Fix an osculating Riemannian metric g^{γ} for (2.1) and (2.2). For brevity, in this proof we use notation X' for the covariant derivative $D_{\gamma}X$ of a vector field X along γ . From (2.3) we have J(0) = 0,

$$J'(0) = \frac{\nabla^{\gamma}}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \gamma_{v+sw}(t) = \frac{d}{ds} \Big|_{s=0} \frac{d}{dt} \Big|_{t=0} \gamma_{v+sw}(t) = w ,$$

and $J''(0) = -R^{\gamma}(0) = 0$. The vector field $X(t) := t \cdot W(t)$ has the same value and the same first and second covariant derivatives at t = 0, namely X(0) = 0, X'(0) = W(0) = w and X''(0) = 2W'(0) = 0 since W is D_{γ} -parallel. Therefore $J(t) - X(t) = O(t^3)$ as $t \to 0$.

Lemma 2.3. Let g_1, g_2 be two Riemannian metrics on M. Suppose that a path γ is a geodesic for both g_1 and g_2 and the two metrics induce the same Jacobi fields along γ , i.e. $\mathcal{J}^{g_1}(\gamma) = \mathcal{J}^{g_2}(\gamma)$. Then g_1 and g_2 induce the same covariant derivative and the same Jacobi operator along γ .

In particular, for a Finsler manifold (M, F) and an F-geodesic γ , the operators D_{γ} and R^{γ} are independent of the choice of the osculating metric g^{γ} .

Proof. We apply the computation from Lemma 2.2 to the Riemannian metrics g_i . The formula (2.4) determines the first order jet of W(t) at t=0 in terms of the Jacobi field J. Hence the space $\mathcal{J} = \mathcal{J}^{g_i}(\gamma)$ determines (independently of the metric) the set of vector fields W along γ such that $D_{\gamma}W(0) = 0$. Indeed, a vector field satisfies

 $D_{\gamma}W(0)=0$ if and only if there exists $J\in\mathcal{J}$ such that (2.4) holds. Similarly, for every $t_0\in\mathbb{R}$ the set of vector fields W along γ satisfying $D_{\gamma}W(t_0)=0$ is determined by \mathcal{J} .

Thus the set of parallel fields along γ is determined by $\mathcal J$ and hence it is the same set for g_1 and g_2 . The parallel fields determine the covariant derivative D_{γ} . The Jacobi equation (2.2), $\mathcal J$ and D_{γ} together determine the Jacobi operator R^{γ} . This proves the first statement of the lemma. The second one follows by Lemma 2.1. \square

With Lemma 2.3 at hand, one can define the covariant derivative D_{γ} and the Jacobi operator R^{γ} along a Finsler geodesic γ as those of any osculating Riemannian metric associated to γ . Note that with these definitions the first part of Lemma 2.3 applies to Finsler metrics as well. In particular, affinely equivalent Finsler metrics induce the same covariant derivatives and the same Jacobi operators.

Following [6], we use the term *linear parallel transport* for the parallel transport operator induced by D_{γ} and refer to D_{γ} -parallel vectors fields as *linearly parallel* vector fields along γ .

2.3. Flag curvature. The *flag curvature* $K_F(v,\sigma)$ of a Finsler manifold (M,F) is a function of a point $p \in M$, a plane (i.e., two-dimensional linear subspace) $\sigma \subset T_pM$, and a direction $v \in \sigma \setminus \{0\}$ in that plane. It can be defined as the sectional curvature $K_g(\sigma)$ of an osculating Riemannian metric $g = g^{\gamma}$ for the geodesic $\gamma = \gamma_v$. From the Riemannian Jacobi equation one sees that

$$K_F(v,\sigma) = K_g(\sigma) = \frac{g(R^{\gamma}(w), w)}{|v \wedge w|_{\sigma}^2}$$
 (2.5)

for any vector $w \in \sigma$ such that v and w are linearly independent, where

$$|v \wedge w|_g^2 = g(v, v)g(w, w) - g(v, w)^2.$$

The formula (2.5) and Lemma 2.3 imply that the definition does not depend on the choice of the osculating metric g. If the metric F is Riemannian, then $K_F(v,\sigma)$ equals its sectional curvature at σ , in particular it does not depend on v.

Consider the Jacobi operator R^{γ} at $p=\gamma(0)$. By symmetries of the Riemannian curvature tensor, this operator on T_pM is symmetric with respect to the inner product induced by $g=g^{\gamma}$. Hence it has only real eigenvalues. By (2.5), the flag curvature of (M,F) is nonpositive if and only if these eigenvalues are all nonpositive for every p and γ . Since Jacobi operators of affinely equivalent metrics coincide, this implies the following:

Lemma 2.4. Let F_1 and F_2 be affinely equivalent Finsler metrics and the flag curvature of F_1 is nonpositive. Then the flag curvature of F_2 is nonpositive as well.

2.4. Berwald metrics. Berwald metrics are a special class of Finsler metrics that can be defined in many equivalent ways, see [2, Ch. 10] and [6, Ch. 4]. A definition convenient for our purposes is the following. A Finsler manifold (M, F) is Berwald if and only if there exists a symmetric affine connection on M such that the geodesics of F are also geodesics of this connection.

By Szabó's metrization theorem [20], this affine connection can be realized as the Levi-Civita connection of a Riemannian metric. Thus a Finsler metric is Berwald if and only if it is affinely equivalent to a Riemannian metric.

Finally, we need the fact that if the Levi-Civita connection of a Riemannian metric g preserves a Finsler norm F, then F and g are affinely equivalent (and hence F is Berwald). See e.g. [6, Lemma 4.1.5] for a proof. Alternatively, one can verify directly that in this case the Euler-Lagrange equation for Finsler geodesics is satisfied by Riemannian geodesics. This is easy to see in the Riemannian exponential coordinates at the point in question.

3. An implication of Busemann convexity

The aim of this section is the following:

Lemma 3.1. Let (M, F) be a Busemann NPC Finsler manifold and $\gamma: (a, b) \to M$ a geodesic. Then the linear parallel transport along γ preserves F, i.e., for every linearly parallel vector field W along γ the function $t \mapsto F(W(t))$ is constant.

Proof. The statement is local, thus we may assume that γ is contained in a Busemann convex open subset of M.

First observe that for every Jacobi field J along γ , its norm F(J(t)) is a convex function of t. Indeed, let $\{\gamma_s\}$ be a geodesic variation of γ whose variation field is J. By Busemann convexity, for every s the function $f_s(t) = d_F(\gamma(t), \gamma_s(t))$ is convex. Hence so is the limit function $\lim_{s\to 0} \frac{f_s(t)}{s} = F(J(t))$.

Now let $W \neq 0$ be a linearly parallel vector field along γ . Note that F(W(t)) is a smooth function of t. Assuming that $0 \in (a,b)$, Lemma 2.3 implies that there is a Jacobi field J along γ such that

$$J(t) = t \cdot W(t) + O(t^3), \quad t \to 0.$$

For this Jacobi field J and t > 0 we have

$$F(J(t)) = F(t \cdot W(t)) + O(t^3) = t \cdot F(W(t)) + O(t^3), \quad t \to 0_+.$$
 (3.1)

In particular $F(J(t)) = t \cdot F(W(0)) + o(t)$ as $t \to 0_+$. Since F(J(t)) is a convex function, it follows that $F(J(t)) \ge t \cdot F(W(0))$ for all $t \ge 0$. This and (3.1) imply that

$$F(W(t)) \ge F(W(0)) - O(t^2), \quad t \to 0_+,$$

hence $F(W(t))'_{t=0} \ge 0$.

Similarly, for the Jacobi field -J and t < 0 we have

$$F(-J(t)) = -t \cdot F(W(t)) + O(t^3), \quad t \to 0_-,$$

and therefore $F(-J(t)) \ge -t \cdot F(W(0))$ for all $t \le 0$. These relations imply that

$$F(W(t)) \ge F(W(0)) - O(t^2), \quad t \to 0_-,$$

hence $F(W(t))'_{t=0} \le 0$. Thus $F(W(t))'_{t=0} = 0$.

Shifting the parameter of γ we deduce that the derivative of F(W(t)) vanishes everywhere on (a,b). Therefore F(W(t)) is constant.

4. Linear parallel transport and Berwaldness

In this section we prove three characterizations of Berwald metrics in terms of the linear parallel transport. The proof of Theorems 1 and 2 in the next section uses Proposition 4.2, which in its turn refers to Proposition 4.1.

Proposition 4.1. Let (M, F) be a Finsler manifold. Let g be a smooth Riemannian metric on M such that the linear parallel transport (determined by F) along any F-geodesic preserves g. Then g is affinely equivalent to F, in particular F is Berwald.

Proof. Let ∇ denote the Levi-Civita connection induced by g. For $p \in M$ and $v \in T_pM \setminus \{0\}$ define $\kappa(v) \in T_pM$ by

$$\kappa(v) = \frac{\nabla}{dt} \Big|_{t=0} \dot{\gamma}_v(t) \,,$$

where γ_v is, as usual, the F-geodesic with $\dot{\gamma}_v(0) = v$. That is, $\kappa(v)$ is the Riemannian second derivative of γ_v . Note that $\kappa(v)$ depends smoothly on v. We need to show that every geodesic of the Finsler manifold (M, F) is a geodesic in (M, g). Thus, it suffices to prove that $\kappa(v) = 0$ for all p and v.

Let W be a linearly parallel vector field along γ_v and w=W(0). We claim that

$$\frac{d}{ds}\Big|_{s=0}\kappa(v+sw) = 2 \cdot \frac{\nabla}{dt}\Big|_{t=0}W(t). \tag{4.1}$$

To prove this, we use the Riemannian exponential coordinates with respect to g at p to identify a small neighborhood U of p with an open subset of $\mathbb{R}^n = T_p M$. By means of coordinates we also identify vector fields along γ_v with \mathbb{R}^n -valued functions of t. Recall that the Levi-Civita connection coefficients in exponential coordinates vanish at the origin. Hence in these coordinates we have

$$\kappa(v) = \frac{d^2}{dt^2} \Big|_{t=0} \gamma_v(t) \tag{4.2}$$

and

$$\frac{\nabla}{dt}\Big|_{t=0}W(t) = \frac{d}{dt}\Big|_{t=0}W(t).$$

Differentiating (4.2) yields

$$\frac{d}{ds}\Big|_{s=0}\kappa(v+sw) = \frac{\partial^3}{\partial s\partial t^2}\Big|_{(s,t)=(0,0)}\gamma_{v+sw}(t) = \frac{d^2}{dt^2}\Big|_{t=0}J(t),$$

where J is the Jacobi field along γ_v defined by (2.3), see Lemma 2.2. By the relation $J(t) = t \cdot W(t) + O(t^3)$ from Lemma 2.2, the right-hand side can be rewritten as

$$\frac{d^2}{dt^2}\Big|_{t=0}J(t) = \frac{d^2}{dt^2}\Big|_{t=0}(t \cdot W(t)) = 2 \cdot \frac{d}{dt}\Big|_{t=0}W(t) = 2 \cdot \frac{\nabla}{dt}\Big|_{t=0}W(t).$$

This finishes the proof of (4.1).

Now we forget about coordinates and introduce some shortcut notation. We write $\langle \cdot, \cdot \rangle$ instead of $g(\cdot, \cdot)$ and denote by X'(t) the Levi-Civita derivative (induced by g) of a vector field X(t) along a path.

Since g is preserved by the linear parallel transport, for any linearly parallel vector fields X and Y along γ_v , the inner product $\langle X(t), Y(t) \rangle$ is constant, hence

$$0 = \frac{d}{dt} \langle X(t), Y(t) \rangle = \langle X', Y \rangle + \langle X, Y' \rangle. \tag{4.3}$$

Note that $\dot{\gamma}_v$ is linearly parallel along γ_v . For $X=Y=\dot{\gamma}_v$ and t=0 the identity (4.3) boils down to

$$\langle \kappa(v), v \rangle = 0. \tag{4.4}$$

Fix $w \in T_p M$ and let W be the linearly parallel vector field along γ_v with W(0) = w. We replace v by v + sw in (4.4) and differentiate with respect to s using (4.1):

$$0 = \frac{d}{ds}\Big|_{s=0} \langle \kappa(v+sw), v+sw \rangle = \langle 2 \cdot W'(0), v \rangle + \langle \kappa(v), w \rangle.$$

On the other hand, plugging X = W and $Y = \dot{\gamma}_v$ into (4.3) we see that

$$0 = \langle W'(0), v \rangle + \langle w, \kappa(v) \rangle.$$

The last two equations together imply that $\langle \kappa(v), w \rangle = 0$. Since w was arbitrary in $T_p M$, we conclude that $\kappa(v) = 0$. This finishes the proof of the proposition. \square

We now deduce the main ingredient of the proofs of Theorems 1 and 2.

Proposition 4.2. Let (M, F) be a Finsler manifold. Suppose that the linear parallel transport along any geodesic preserves the Finsler norm F. Then there exists a Riemannian metric on M affinely equivalent to F. In particular F is Berwald.

Proof. Consider a smooth Riemannian metric g on M whose value g_p at every $p \in M$ is canonically determined by the norm $F_p := F|_{T_pM}$, where "canonically" means that any linear isometry between normed spaces (T_pM, F_p) and (T_qM, F_q) takes g_p to g_q . See [14] for a possible construction of such g.

Since the linear parallel transports are linear and preserve F, they also preserve g. The claim now follows from Proposition 4.1.

Our next result generalizes Proposition 4.2. In order to state it we need the notion of linear holonomy group, defined as follows. Let (M, F) be a Finsler manifold. For a piecewise geodesic path $\gamma \colon [a,b] \to M$ denote by P_{γ} the linear parallel transport along γ . Recall that P_{γ} is a linear isomorphism from $T_{\gamma(a)}M$ to $T_{\gamma(b)}M$. Fix a point $p \in M$ and let $GL(T_pM)$ denote the group of linear self-isomorphisms of T_pM . The *linear holonomy group* $LH_p(M,F)$ is the subgroup of $GL(T_pM)$ generated by maps of the form $P_{\gamma_2}^{-1} \circ P_{\gamma_1}$ where γ_1 and γ_2 are piecewise geodesic paths starting at p and having a common end point. (The definition is so cumbersome because linear parallel transports in opposite directions are in general not inverse to each other.)

Proposition 4.3. Let (M, F) be a connected Finsler manifold and $p \in M$. Then the closure of the linear holonomy group $LH_p(M, F)$ in $GL(T_pM)$ is compact if and only if F is Berwald.

Proof. Let $H = LH_p(M, F)$. If F is Berwald then by Szabó's metrization theorem the linear parallel transport is the Levi-Civita parallel transport of some Riemannian metric. Thus H preserves an inner product on T_pM and is therefore contained in the corresponding orthogonal group, which is compact.

On the other hand, if the closure of H is compact then there exists an H-invariant inner product g_p on T_pM . For every $x \in M$, pick a piecewise geodesic γ connecting p to x and define an inner product g_x on T_xM as the push-forward of g_p by the linear parallel transport P_γ . Since g_p is H-invariant, the resulting Riemannian metric $g = \{g_x\}_{x \in M}$ is well-defined and it is preserved by the linear parallel transport along any F-geodesic. It is easy to see that g is smooth. By Proposition 4.1 it follows that F is a Berwald metric.

5. Proof of the theorems

We can now collect the harvest and prove Theorems 1 and 2.

The "if" part of Theorem 2 follows from the main result of [11]. To prove the "if" part of Theorem 1, let g be a Riemannian metric on M of nonpositive sectional curvature whose Levi-Civita connection preserves the Finsler norm F. Then F is Berwald and affinely equivalent to g (see Section 2.4). Now Lemma 2.4 implies

that F has nonpositive flag curvature. Hence, by the "if" part of Theorem 2, (M, F) is Busemann NPC.

Now we prove the "only if" implications of the theorems. Let (M, F) be a Busemann NPC Finsler manifold. Then by Lemma 3.1 the linear parallel transport preserves F. Therefore, by Proposition 4.2 the Finsler metric F is Berwald and affinely equivalent to a Riemannian metric g.

It remains to prove that g has nonpositive sectional curvature and F has nonpositive flag curvature. This can be seen from results of [9] but for reader's convenience we give a short proof here.

First let us show that F has nonpositive flag curvature. Suppose the contrary. Then, as explained in Section 2.3, for some geodesic γ the Jacobi operator R^{γ} has a positive eigenvalue at $p=\gamma(0)$. Let $w\in T_pM$ be a nonzero vector such that $R^{\gamma}(w)=\lambda w, \, \lambda>0$. Consider the Jacobi field J along γ with the initial conditions J(0)=w and $D_{\gamma}J(0)=0$. Let $\overline{J}(t)\in T_pM$ be the image of J(t) in T_pM under the linear parallel transport along γ_v . Then $\overline{J}'(0)=D_{\gamma}J(0)=0$ and

$$\bar{J}''(0) = D_{\gamma} D_{\gamma} J(0) = -R^{\gamma} (J(0)) = -\lambda w.$$

Since the linear parallel transport preserves F, we have $F(J(t)) = F(\overline{J}(t))$ for all t. Therefore

$$F(J(t))_{t=0}^{"} = F(\overline{J}(t))_{t=0}^{"} = -\lambda \cdot F(w) < 0,$$

contrary to the fact that F(J(t)) is a convex function (see the proof of Lemma 3.1). This contradiction shows that F has nonpositive flag curvature and finishes the proof of Theorem 2.

Now Lemma 2.4 and the affine equivalence of F and g imply that g has nonpositive sectional curvature. This finishes the proof of Theorem 1.

6. Non-smooth examples

In this section we describe two simple examples of Busemann convex spaces not affinely equivalent to any CAT(0) space. Both examples are constructed from a plane V with a strictly convex non-Euclidean norm.

The first example X_1 arises from V by attaching a ray at the origin.

The second example X_2 is the double branched cover of V with branching locus the origin $0 \in V$. We give X_2 the naturally induced length metric. Note that X_2 is biLipschitz to the Euclidean plane.

It is not difficult to see that X_1 and X_2 are Busemann convex. Assume now that there exists a space Z and an affine bijection $F: X_i \to Z$ for i = 1 or i = 2, hence F sends constant speed geodesics to constant speed geodesics.

Every ray in V starting at the origin can be continued in X_1 to an infinite geodesic by the attached ray in X. Thus, in the case of X_1 , any ray in V is stretched by the

same factor as the attached ray. Hence all rays in V through the origin are stretched by the same factor. The metric on the punctured plane $Z_0 = F(V \setminus \{0\})$ is affinely equivalent to the flat Riemannian manifold $\mathbb{R}^2 \setminus \{0\}$. Due to [13, Theorem 1.5], the metric on Z_0 must come from a constant norm on V. In particular, parallel rays in V must be stretched by F by equal factors. We conclude that F must stretch all distances by the same factor. Thus, F is a dilation and Z cannot be CAT(0).

Every pair of rays $\gamma_{1,2}$ in X_2 starting at the origin can be continued to infinite geodesics by the same ray γ_3 . Therefore, also in case of X_2 , the map F must stretch all rays through the origin by the same factor. Using [13, Theorem 1.5] as above, this again implies that F is just a dilation and Z cannot be CAT(0).

However, in both cases there is a natural CAT(0) metric on the spaces X_1 , X_2 which arises in the same way as $X_{1,2}$ from the Euclidean plane V_0 instead of V. These CAT(0) metrics have the same unparametrized geodesics as the original metrics and are affinely equivalent to the original metric outside the branching locus.

References

- [1] W. Ballmann, *Lectures on spaces of nonpositive curvature*. With an appendix by M. Brin, DMV Seminar, 25, Birkhäuser Verlag, Basel, 1995. Zbl 0834.53003 MR 1377265
- [2] D. Bao, S.-S. Chern, and Z. Shen. *An introduction to Riemann-Finsler geometry*, Graduate Texts in Mathematics, 200, Springer-Verlag, New York, 2000. Zbl 0954.53001 MR 1747675
- [3] M. Berger, Sur les groupes d'holonomie homogène des variétés à connexion affine et des variétés riemanniennes, *Bull. Soc. Math. France*, 83 (1955), 279–330. Zbl 0068.36002 MR 79806
- [4] M. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 319, Springer-Verlag, Berlin, 1999. Zbl 0988.53001 MR 1744486
- [5] H. Busemann, Spaces with non-positive curvature, *Acta Math.*, 80 (1948), 259–310.Zbl 0038.10005 MR 29531
- [6] S. Chern and Z. Shen, *Riemann-Finsler geometry*, Nankai Tracts in Mathematics, 6, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005. Zbl 1085.53066 MR 2169595
- [7] M. Kapovich, B. Leeb, and J. Porti, Finsler bordification of symmetric and certain locally symmetric spaces, *Geom. Topol.*, 22 (2018), no. 5, 2533–2646. Zbl 1417.53058 MR 3811766
- [8] P. Kelly and E. G. Straus, Curvature in Hilbert geometries. II, *Pacific J. Math.*, **25** (1968), 549–552. Zbl 0169.24605 MR 232287
- [9] A. Kristály and L. Kozma, Metric characterization of Berwald spaces of non-positive flag curvature, *J. Geom. Phys.*, **56** (2006), no. 8, 1257–1270. Zbl 1103.53046 MR 2234441
- [10] A. Kristály and A. Roth, Testing metric relations on Finsler manifolds via a geodesic detecting algorithm, in *Conference Proceedings of the 9th IEEE International Symposium on Applied Computational Intelligence and Informatics. SACI 2014 (Timisoara, Romania, 15–17 May, 2014)*, Institute of Electrical and Electronics Engineers (IEEE), 2014.

- [11] A. Kristály, C. Varga, and L. Kozma. The dispersing of geodesics in Berwald spaces of non-positive flag curvature, *Houston J. Math.*, 30 (2004), no. 2, 413–420. Zbl 1075.53077 MR 2084910
- [12] U. Lang, Injective hulls of certain discrete metric spaces and groups, *J. Topol. Anal.*, **5** (2013), no. 3, 297–331. Zbl 1292.20046 MR 3096307
- [13] A. Lytchak, Affine images of Riemannian manifolds, *Math. Z.*, **270** (2012), no. 3-4, 809–817. Zbl 1250.53035 MR 2892925
- [14] V. S. Matveev and M. Troyanov, The Binet-Legendre metric in Finsler geometry, *Geom. Topol.*, **16** (2012), no. 4, 2135–2170. Zbl 1258.53080 MR 3033515
- [15] Z. Muzsnay and P. Nagy, Holonomy Theory of Finsler Manifolds, in *Lie Groups, Differential Equations, and Geometry: Advances and Surveys*, 265–320, UNIPA Springer Ser., Springer, Cham, 2017. Zbl 1392.53079 MR 3726538
- [16] A. Papadopoulos, *Metric spaces, convexity and non-positive curvature*. Second edition, IRMA Lectures in Mathematics and Theoretical Physics, 6, European Mathematical Society (EMS), Zürich, 2014. Zbl 1296.53007 MR 3156529
- [17] H. Rademacher, Nonreversible Finsler metrics of positive flag curvature, in *A sampler of Riemann-Finsler geometry*, 261–302, Math. Sci. Res. Inst. Publ., 50, Cambridge Univ. Press, Cambridge, 2004. Zbl 1074.53062 MR 2132661
- [18] Z. Shen, *Differential geometry of spray and Finsler spaces*, Kluwer Academic Publishers, Dordrecht, 2001. Zbl 1009.53004 MR 1967666
- [19] Z. Shen, Some open problems in Finsler geometry, *preprint*, 2009. Available at: https://www.math.iupui.edu/~zshen/Research/papers/Problem.pdf
- [20] Z. I. Szabó. Positive definite Berwald spaces. Structure theorems on Berwald spaces. *Tensor* (*N.S.*), **35** (1981), no. 1, 25–39. Zbl 0464.53025 MR 614132

Received January 4, 2018

S. Ivanov, St. Petersburg Department of Steklov Mathematical Institute, Russian Academy of Sciences, Fontanka 27, St. Petersburg 191023, Russia E-mail: svivanov@pdmi.ras.ru

A. Lytchak, Mathematisches Institut, Universität Köln, Weyertal 86–90, 50931 Köln, Germany

E-mail: alytchak@math.uni-koeln.de