

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 95 (2020)
Heft: 1

Artikel: On the universality of the Epstein zeta function
Autor: Andersson, Johan / Södergren, Anders
DOI: <https://doi.org/10.5169/seals-882409>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On the universality of the Epstein zeta function

Johan Andersson and Anders Södergren*

Abstract. We study universality properties of the Epstein zeta function $E_n(L, s)$ for lattices L of large dimension n and suitable regions of complex numbers s . Our main result is that, as $n \rightarrow \infty$, $E_n(L, s)$ is universal in the right half of the critical strip as L varies over all n -dimensional lattices L . The proof uses a novel combination of an approximation result for Dirichlet polynomials, a recent result on the distribution of lengths of lattice vectors in a random lattice of large dimension and a strong uniform estimate for the error term in the generalized circle problem. Using the same approach we also prove that, as $n \rightarrow \infty$, $E_n(L_1, s) - E_n(L_2, s)$ is universal in the full half-plane to the right of the critical line as (L_1, L_2) varies over all pairs of n -dimensional lattices. Finally, we prove a more classical universality result for $E_n(L, s)$ in the s -variable valid for almost all lattices L of dimension n . As part of the proof we obtain a strong bound of $E_n(L, s)$ on the critical line that is subconvex for $n \geq 5$ and almost all n -dimensional lattices L .

Mathematics Subject Classification (2010). 11E45, 30K10, 41A30; 11H06, 60G55.

Keywords. Epstein zeta function, universality, random lattice, Poisson process, subconvexity.

1. Introduction

In 1975 Voronin [27, 28] proved the following remarkable approximation theorem for the Riemann zeta function:

Theorem 1.1 (Voronin). *Let $K = \{s \in \mathbb{C} : |s - \frac{3}{4}| \leq r\}$ for some $r < \frac{1}{4}$, and suppose that f is any nonvanishing continuous function on K that is analytic in the interior of K . Then, for any $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \operatorname{meas} \left\{ t \in [0, T] : \max_{s \in K} |\zeta(s + it) - f(s)| < \varepsilon \right\} > 0.$$

This theorem, known as *Voronin's Universality Theorem*, shows that any nonvanishing analytic function in a small disc may be approximated by a vertical

*The second author was partially supported by a postdoctoral fellowship from the Swedish Research Council, by the National Science Foundation under agreement No. DMS-1128155, as well as by a grant from the Danish Council for Independent Research and FP7 Marie Curie Actions-COFUND (grant id: DFF-1325-00058).

shift of the Riemann zeta function. It has been improved and generalized in various directions (see Steuding's monograph [20] and Matsumoto's survey paper [14] for detailed discussions); for example it is known that the set K may be chosen as any compact set with connected complement that lies in the vertical strip

$$D := \{s \in \mathbb{C} : \frac{1}{2} < \operatorname{Re}(s) < 1\}. \quad (1.1)$$

Let us also note that, since $\zeta(s)$ has an Euler product, it is necessary to assume that the function f in Theorem 1.1 is nonvanishing on K . However, for zeta functions without an Euler product, such as the Hurwitz zeta function $\zeta(s, \alpha)$ with rational or transcendental¹ parameter $0 < \alpha < 1$, $\alpha \neq \frac{1}{2}$, this condition can be removed.

Similar universality theorems have been proved for large classes of zeta functions and L -functions. Here we will content ourselves with a short review of the situation for Dirichlet L -functions. In his thesis, Voronin [30] proved² the *joint universality* of Dirichlet L -functions, i.e. that vertical shifts of Dirichlet L -functions attached to nonequivalent Dirichlet characters can be used to simultaneously approximate any finite number of nonvanishing analytic functions on K . In a different direction, Bagchi [1] has proved a universality theorem for Dirichlet L -functions $L(s, \chi)$ in which the imaginary shifts in the complex argument s from Theorem 1.1 has been replaced by a variation of the character χ over the set of characters of prime modulus. To be precise, let K be a compact subset of D with connected complement and let f be a nonvanishing continuous function on K that is analytic in the interior of K . Then, for any $\varepsilon > 0$,

$$\liminf_{p \rightarrow \infty} \frac{1}{\varphi(p)} \# \{ \chi \bmod p : \max_{s \in K} |L(s, \chi) - f(s)| < \varepsilon \} > 0. \quad (1.2)$$

Recently Mishou and Nagoshi [15, 16] proved a related result for Dirichlet L -functions associated with real characters. Let \mathcal{D}^+ (resp. \mathcal{D}^-) denote the set of positive fundamental discriminants (resp. negative fundamental discriminants) and define

$$\mathcal{D}^\pm(X) := \{d \in \mathcal{D}^\pm : |d| \leq X\}.$$

For a discriminant d , we let χ_d denote the quadratic Dirichlet character modulo $|d|$ defined by the Kronecker symbol $\chi_d(n) = (\frac{d}{n})$.

Theorem 1.2 (Mishou–Nagoshi). *Let Ω be a simply connected region in D that is symmetric with respect to the real axis. Then, for any $\varepsilon > 0$, any compact set $K \subset \Omega$ and any nonvanishing holomorphic function f on Ω which takes positive real values on $\Omega \cap \mathbb{R}$, we have*

$$\liminf_{X \rightarrow \infty} \frac{1}{\#\mathcal{D}^\pm(X)} \# \{ d \in \mathcal{D}^\pm(X) : \max_{s \in K} |L(s, \chi_d) - f(s)| < \varepsilon \} > 0.$$

¹For the case of algebraic irrational α , the Hurwitz zeta function does not have an Euler product and it is likely to be universal without assuming the nonvanishing condition. However, proving this seems quite difficult and constitutes a major open problem in the theory of universality.

²Similar results were established (independently) by Gonek [9] and Bagchi [1].

The purpose of the present paper is to prove universality results for the Epstein zeta function that turn out to have similarities with both Theorem 1.2 and Bagchi's result described in (1.2) above. In order to state and describe our results, we first need to properly introduce the setting.

Let X_n denote the space of all n -dimensional lattices $L \subset \mathbb{R}^n$ of covolume one and let μ_n denote Siegel's measure [18] on X_n , normalized to be a probability measure. For $L \in X_n$ and $\text{Re}(s) > \frac{n}{2}$, the Epstein zeta function is defined by

$$E_n(L, s) := \sum_{\mathbf{m} \in L \setminus \{0\}} |\mathbf{m}|^{-2s}.$$

$E_n(L, s)$ has an analytic continuation to \mathbb{C} except for a simple pole at $s = \frac{n}{2}$ with residue $\pi^{\frac{n}{2}} \Gamma(\frac{n}{2})^{-1}$. Furthermore, $E_n(L, s)$ satisfies the functional equation

$$F_n(L, s) = F_n(L^*, \frac{n}{2} - s), \quad (1.3)$$

where $F_n(L, s) := \pi^{-s} \Gamma(s) E_n(L, s)$ and L^* is the dual lattice of L . The Epstein zeta function has many properties in common with the Riemann zeta function. In fact, the functions $E_n(L, s)$ (actually a slightly more general family of functions) were introduced by Epstein [5, 6] in an attempt to find the most general form of a function satisfying a functional equation of the same type as the Riemann zeta function. Note in particular the relation

$$E_1(\mathbb{Z}, s) = 2\zeta(2s).$$

However, we stress that there are also important differences between $E_n(L, s)$ and $\zeta(s)$. Typically $E_n(L, s)$ has no Euler product and it is well known that the Riemann hypothesis for $E_n(L, s)$ generally fails (cf., e.g., [21] and the references therein).³

Let V_n denote the volume of the n -dimensional unit ball. We recall the explicit formula

$$V_n = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)},$$

and stress that V_n decays extremely fast as $n \rightarrow \infty$. In most of our results V_n will appear naturally as part of a factor normalizing $E_n(L, s)$.

Our first main result is a universality theorem for $E_n(L, s)$ in the lattice aspect, i.e. a universality result where the lattice L varies over the space X_n but no vertical shifts are applied to the complex variable s . The situation is related to the one in Bagchi's theorem (1.2), and similarly, in order to obtain still finer approximations it is natural to consider the limit $n \rightarrow \infty$. Let us also point out that the relation with Theorem 1.2 lies in the fact that both $L(s, \chi_d)$ and $E_n(L, s)$ are real-valued for real values of s , resulting in the same sort of conditions on the involved functions and

³Here the words *typically* and *generally* are to be interpreted in terms of the measure μ_n .

regions. However, since $E_n(L, s)$ typically has no Euler product, the nonvanishing condition on the function f can be removed.

Theorem 1.3. *Let Ω be a simply connected region in D that is symmetric with respect to the real axis. Then, for any $\varepsilon > 0$, any compact set $K \subset \Omega$ and any holomorphic function f on Ω that is real-valued on $\Omega \cap \mathbb{R}$, we have*

$$\liminf_{n \rightarrow \infty} \text{Prob}_{\mu_n} \left\{ L \in X_n : \max_{s \in K} \left| 2^{s-1} V_n^{-s} E_n \left(L, \frac{ns}{2} \right) - f(s) \right| < \varepsilon \right\} > 0.$$

Remark 1.4. In particular it follows from Theorem 1.3 that, given $\varepsilon > 0$, $K \subset \Omega$ and f as above, there exists some $n \in \mathbb{Z}^+$ and a lattice $L \in X_n$ such that

$$\max_{s \in K} \left| 2^{s-1} V_n^{-s} E_n \left(L, \frac{ns}{2} \right) - f(s) \right| < \varepsilon.$$

As an immediate consequence of Theorem 1.3, we have the following denseness result.

Corollary 1.5. *For any fixed $s \in D \setminus (D \cap \mathbb{R})$ the set*

$$\left\{ 2^{s-1} V_n^{-s} E_n \left(L, \frac{ns}{2} \right) : n \in \mathbb{Z}^+, L \in X_n \right\}$$

is dense in \mathbb{C} . Moreover, for any fixed $x \in D \cap \mathbb{R}$, we have

$$\left\{ 2^{x-1} V_n^{-x} E_n \left(L, \frac{nx}{2} \right) : n \in \mathbb{Z}^+, L \in X_n \right\} = \mathbb{R}. \quad (1.4)$$

To prove the first part of the corollary, it is sufficient to note that for any $c \in \mathbb{C}$ we may use Theorem 1.3 to approximate $f(s) = c$ on the set $K = \{s\}$ to arbitrary precision. Next, to prove (1.4) we note that for any $N \in \mathbb{Z}^+$ we may use Theorem 1.3, with $\varepsilon = 1$ and $K = \{x\}$, to approximate the two functions

$$f_1(s) = -N - 1 \quad \text{and} \quad f_2(s) = N + 1.$$

It follows that for all sufficiently large n (depending on N) there exist lattices $L_1, L_2 \in X_n$ such that

$$2^{x-1} V_n^{-x} E_n \left(L_1, \frac{nx}{2} \right) < -N \quad \text{and} \quad 2^{x-1} V_n^{-x} E_n \left(L_2, \frac{nx}{2} \right) > N.$$

We note that in contrast to the situation in [15, Corollary 1.2], the lattice variable L varies continuously in X_n and, moreover, the Epstein zeta function is continuous in the lattice variable.⁴ Hence, since X_n is connected and the Epstein zeta function is real-valued on the real line, the intermediate value theorem implies that

$$[-N, N] \subseteq \left\{ 2^{x-1} V_n^{-x} E_n \left(L, \frac{nx}{2} \right) : L \in X_n \right\}$$

⁴In fact, $E_n(L, s)$ is an Eisenstein series associated to a maximal parabolic subgroup of $\text{SL}(n, \mathbb{Z})$ (see, e.g., [8, Section 10.7]). See also [26, Theorem 2].

for all sufficiently large n . This confirms that the left-hand side of (1.4) is not only dense in \mathbb{R} , but in fact equal to \mathbb{R} .

Let us also point out that Theorem 1.3, together with the Weierstrass approximation theorem, readily gives a universality result for approximation of continuous functions on closed subintervals of $(\frac{1}{2}, 1)$.

Corollary 1.6. *Let $\frac{1}{2} < a < b < 1$. Then, for any $\varepsilon > 0$ and any continuous real-valued function f on $[a, b]$, we have*

$$\liminf_{n \rightarrow \infty} \text{Prob}_{\mu_n} \left\{ L \in X_n : \max_{x \in [a, b]} |2^{x-1} V_n^{-x} E_n \left(L, \frac{nx}{2} \right) - f(x)| < \varepsilon \right\} > 0.$$

We will now use Corollary 1.6 to give a short and more elementary proof of (one part of) [25, Corollary 1.5]. Let $\varepsilon < 1$ and let, for any $\delta \in (0, \frac{1}{4})$, K_δ denote the closed interval $[\frac{1}{2} + \delta, 1 - \delta] \subset (\frac{1}{2}, 1)$. Now, by applying Corollary 1.6 with the continuous function $f(x) = -1$ on K_δ , we obtain the following result.

Corollary 1.7. *For any $\delta \in (0, \frac{1}{4})$, we have*

$$\liminf_{n \rightarrow \infty} \text{Prob}_{\mu_n} \left\{ L \in X_n : E_n \left(L, \frac{nx}{2} \right) < 0 \text{ for all } x \in [\frac{1}{2} + \delta, 1 - \delta] \right\} > 0.$$

Remark 1.8. It follows from [25, Corollary 1.5] that we can replace the \liminf in Corollary 1.7 by a proper limit. Let us also point out that it follows from [25, Corollary 1.7] that this limit probability tends to zero as δ tends to zero.

Our next corollary discusses zeros of $E_n(L, \frac{nx}{2})$ in the interval $\frac{1}{2} < x < 1$. Its proof is an application of Theorem 1.3 with the function $f(s) = \sin(\pi(\frac{m(s-a)}{b-a} + \frac{1}{2}))$, together with Rouche's theorem and the observation that since $E_n(L, \frac{nx}{2})$ is real-valued the zeros of $E_n(L, \frac{ns}{2})$ approximating the zeros of $f(s)$ must remain on the real line.⁵

Corollary 1.9. *Let $\frac{1}{2} < a < b < 1$ and let $m \in \mathbb{Z}^+$. Then we have*

$$\liminf_{n \rightarrow \infty} \text{Prob}_{\mu_n} \left\{ L \in X_n : E_n \left(L, \frac{nx}{2} \right) \text{ has exactly } m \text{ zeros in } [a, b] \right\} > 0.$$

Let us give a brief sketch of the proof of Theorem 1.3, which is given in full detail in Section 3. To begin, we decompose the expression under consideration into two parts as

$$\left(2^{s-1} V_n^{-s} E_n \left(L, \frac{ns}{2} \right) - \zeta(s) \right) + (\zeta(s) - f(s)), \quad (1.5)$$

and approximate the second part, i.e. $\zeta(s) - f(s)$ (which is holomorphic on the compact set K), to any desired accuracy by a Dirichlet polynomial. We continue by dividing the first part of (1.5) into a main term and a tail term. Using a recent

⁵If we allow the number of zeros to also be greater than m , then this result is an immediate consequence of Corollary 1.6.

result of the second author [22] on the distribution of lengths of lattice vectors in a random lattice of large dimension, we show that the main term approximate the Dirichlet polynomial found above as well as desired for a positive proportion of lattices $L \in X_n$. To conclude the proof we use a strong uniform estimate of the error term in the generalized circle problem (cf. [17, 25]; see also Section 2.1) to show that the tail term is sufficiently small.

Using the same general idea of proof, we arrive at our second main result. As far as we are aware, this is the first example of a universality theorem that is valid in the half-plane of absolute convergence (of the Dirichlet series under consideration).

Theorem 1.10. *Let Ω be a simply connected region in the half-plane $\operatorname{Re}(s) > \frac{1}{2}$ that is symmetric with respect to the real axis. Then, for any $\varepsilon > 0$, any compact set $K \subset \Omega$ and any holomorphic function f on Ω that is real-valued on $\Omega \cap \mathbb{R}$, we have*

$$\liminf_{n \rightarrow \infty} \operatorname{Prob}_{\mu_n \times \mu_n} \left\{ (L_1, L_2) \in X_n^2 : \max_{s \in K} \left| 2^{s-1} V_n^{-s} E_n \left(L_1, \frac{ns}{2} \right) - 2^{s-1} V_n^{-s} E_n \left(L_2, \frac{ns}{2} \right) - f(s) \right| < \varepsilon \right\} > 0.$$

Classical Voronin universality (variants of Theorem 1.1) for the Epstein zeta function follows in some special cases, such as when the Epstein zeta function is (a constant multiple of) a Dedekind zeta function of an imaginary quadratic number field, since then it is a product of $\zeta(s)$ and a Dirichlet L -function, and universality follows from the joint universality of Dirichlet L -functions. Universality has also been proved in the case when the Epstein zeta function can be written as a linear combination of Hecke L -functions by joint universality for such functions, see [13, pp. 279–284] and [29]. The general case seems to be more difficult. While we cannot prove classical universality for $E_n(L, s)$ with any single given lattice, we will prove universality for almost every lattice $L \in X_n$. At present time our methods produce a result in the strip $\frac{3}{4} < \operatorname{Re}(s) < 1$.⁶

Theorem 1.11. *Let $n \geq 2$. Then, for almost all $L \in X_n$ and for any $\varepsilon > 0$, any compact set $K \subset \{s \in \mathbb{C} : \frac{3}{4} < \operatorname{Re}(s) < 1\}$ with connected complement, and any function f that is continuous on K and analytic in the interior of K , we have*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \operatorname{meas} \left\{ t \in [0, T] : \max_{s \in K} \left| E_n \left(L, \frac{n(s+it)}{2} \right) - f(s) \right| < \varepsilon \right\} > 0.$$

There are a number of standard corollaries of this type of universality (see, e.g., [20]). As one such example, we mention an immediate consequence of Theorem 1.11 that follows by approximating $f(s) = s - \frac{a+b}{2}$ on

$$K = \left\{ s \in \mathbb{C} : \left| s - \frac{a+b}{2} \right| \leq \frac{b-a}{2} \right\}$$

and using Rouché's theorem.

⁶We note that a result recently stated by Blomer [3, p. 2] implies that Theorem 1.11 holds for all compact sets $K \subset D = \{s \in \mathbb{C} : \frac{1}{2} < \operatorname{Re}(s) < 1\}$ with connected complement.

Corollary 1.12. *Let $n \geq 2$ and let $\frac{3}{4} < a < b < 1$. Then, for almost all $L \in X_n$, we have*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \# \left\{ \rho \in \mathbb{C} : |\operatorname{Im}(\rho)| < T, a < \operatorname{Re}(\rho) < b, E_n \left(L, \frac{n\rho}{2} \right) = 0 \right\} > 0.$$

This corollary complements the recent results in [21], that treats zeros in the half-plane $\operatorname{Re}(s) > 1$, by showing that also the zeros in the strip $\{s : \frac{3}{4} < \operatorname{Re}(s) < 1\}$ violates the Riemann hypothesis in a strong way. We also note that the corresponding result for c -values (values s such that $E_n(L, \frac{ns}{2}) = c$) follows by a similar argument.

Finally, we mention that the proof of Theorem 1.11 uses a result of Drungilas–Garunkštis–Kačėnas [4] on universality of general Dirichlet series. The main ingredient in the proof is a new mean square estimate that is valid for $\operatorname{Re}(s) > \frac{3}{4}$ and almost all lattices $L \in X_n$ (cf. Theorem 5.1). In the process of proving this estimate, we discover a strong bound on the growth of the Epstein zeta function on the critical line, which we find interesting in its own right.

Theorem 1.13. *Let $n \geq 2$ and let $\varepsilon > 0$. Then, for almost all $L \in X_n$, we have*

$$E_n \left(L, \frac{n}{4} + it \right) = O_{L,\varepsilon} \left((1 + |t|)^{1+\varepsilon} \right). \quad (1.6)$$

Recalling that the corresponding convexity bound states that

$$E_n \left(L, \frac{n}{4} + it \right) = O_{L,\varepsilon} \left((1 + |t|)^{\frac{n}{4} + \varepsilon} \right)$$

for all $L \in X_n$, it follows that (1.6) provides a subconvex estimate for almost all $L \in X_n$ as soon as $n \geq 5$. In fact, the bound (1.6) is majorized by any positive power δ of the convexity bound, in all sufficiently large dimensions n (depending on δ).

Theorem 1.13 has recently been improved by Blomer [3]; using the spectral theory of automorphic forms, Blomer arrives at the estimate

$$E_n \left(L, \frac{n}{4} + it \right) = O_{L,\varepsilon} \left((1 + |t|)^{\frac{1}{2} + \varepsilon} \right)$$

for almost all $L \in X_n$. (A detailed proof of this result is provided only for almost all orthogonal lattices). See also [19] for the even stronger bound

$$E_2 \left(L, \frac{1}{2} + it \right) = O_{L,\varepsilon} \left((1 + |t|)^{\frac{1}{4} + \varepsilon} \right)$$

valid for almost all $L \in X_2$.

2. Preliminaries

2.1. Poisson distribution of vector lengths. Recall that we use V_n to denote the volume of the unit ball in \mathbb{R}^n . Given a lattice $L \in X_n$, we order its nonzero vectors

by increasing lengths as $\pm \mathbf{v}_1, \pm \mathbf{v}_2, \pm \mathbf{v}_3, \dots$, set $\ell_j = |\mathbf{v}_j|$ (thus $0 < \ell_1 \leq \ell_2 \leq \ell_3 \leq \dots$), and define

$$\mathcal{V}_j(L) := \frac{1}{2} V_n \ell_j^n, \quad (2.1)$$

so that $\mathcal{V}_j(L)$ equals one half of the volume of an n -dimensional ball of radius ℓ_j . The main result in [22] states that the volumes $\{\mathcal{V}_j(L)\}_{j=1}^\infty$ determined by a random lattice $L \in X_n$ converges in distribution, as $n \rightarrow \infty$, to the points of a Poisson process on the positive real line with constant intensity 1. In other words, if, for $t \geq 0$, we let

$$N_{n,L}(t) := \#\{j : \mathcal{V}_j(L) \leq t\},$$

then we have the following theorem.

Theorem 2.1. *Let $\mathcal{P} = \{\mathcal{N}(t), t \geq 0\}$ be a Poisson process on the positive real line with intensity 1. Then the stochastic process $\{N_{n,L}(t), t \geq 0\}$ converges weakly to \mathcal{P} as $n \rightarrow \infty$.*

Given $L \in X_n$ and $t \geq 0$, we define

$$R_{n,L}(t) := N_{n,L}(t) - t. \quad (2.2)$$

Note that $1 + 2R_{n,L}(t/2)$ equals the error term in the circle problem generalized to an n -dimensional ball of volume t and a general lattice $L \in X_n$. We recall the following bound on $R_{n,L}(t)$ and refer to [25, Theorem 1.3] (see also [17]) for a proof.

Theorem 2.2. *For all $\varepsilon > 0$ there exists $C_\varepsilon > 0$ such that for all $n \geq 3$ and $C \geq 1$ we have*

$$\text{Prob}_{\mu_n} \{L \in X_n : |R_{n,L}(t)| \leq C_\varepsilon (Ct)^{\frac{1}{2}} (\log t)^{\frac{3}{2} + \varepsilon}, \forall t \geq 5\} \geq 1 - C^{-1}. \quad (2.3)$$

In our discussion, we will often find it convenient to work with a close relative of $R_{n,L}(t)$, namely

$$S_{n,L}(t) := N_{n,L}(t) - \#\{j \in \mathbb{Z}^+ : j \leq t\} \quad (t \geq 0). \quad (2.4)$$

The following estimate is an immediate corollary of Theorem 2.2.

Corollary 2.3. *For all $\varepsilon > 0$ there exists $C_\varepsilon > 0$ such that for all $n \geq 3$ and $C \geq 1$ we have*

$$\text{Prob}_{\mu_n} \{L \in X_n : |S_{n,L}(t)| \leq C_\varepsilon (Ct)^{\frac{1}{2}} (\log t)^{\frac{3}{2} + \varepsilon}, \forall t \geq 5\} \geq 1 - C^{-1}.$$

Remark 2.4. Let us note that in the case $n = 2$, Schmidt [17, Theorem 2] has proved that for almost all $L \in X_2$ and for all sufficiently large t (depending on L and ε), we have

$$|R_{2,L}(t)| \ll_{L,\varepsilon} t^{\frac{1}{2}} (\log t)^{\frac{5}{2} + \varepsilon}.$$

Hence, for the same L and t , we also have

$$|S_{2,L}(t)| \ll_{L,\varepsilon} t^{\frac{1}{2}} (\log t)^{\frac{5}{2} + \varepsilon}.$$

2.2. Normalization of $E_n(L, s)$. For any complex number s with $\operatorname{Re}(s) > 1$, we can use (2.1) to write $E_n(L, \frac{ns}{2})$ in the form

$$E_n(L, \frac{ns}{2}) = 2^{1-s} V_n^s \sum_{j=1}^{\infty} \mathcal{V}_j(L)^{-s}.$$

We find it natural to consider the normalized function

$$\mathcal{E}_n(L, s) := 2^{s-1} V_n^{-s} E_n(L, \frac{ns}{2}),$$

so that

$$\mathcal{E}_n(L, s) = \sum_{j=1}^{\infty} \mathcal{V}_j(L)^{-s} \quad (\operatorname{Re}(s) > 1).$$

Note that $\mathcal{E}_n(L, s)$ has a simple pole at $s = 1$ with residue 1. Note also that if $\{T_j\}_{j=1}^{\infty}$ are the points of a Poisson process on the positive real line with intensity 1, then [24, Theorem 1 and Remark 4] state that $\mathcal{E}_n(L, s)$ converges in distribution to $\sum_{j=1}^{\infty} T_j^{-s}$, for any fixed s with $\operatorname{Re}(s) > 1$, as $n \rightarrow \infty$.

3. Proof of Theorem 1.3

To begin with we state the following lemma of Mishou and Nagoshi (cf. [15, Proposition 2.4] for a more general statement). We recall that D denotes the right half of the critical strip (see (1.1)).

Lemma 3.1. *Let Ω be a simply connected region in D that is symmetric with respect to the real axis. Then, for any $\varepsilon > 0$, any compact set $K \subset \Omega$ and any holomorphic function f on Ω that is real-valued on $\Omega \cap \mathbb{R}$, there exist $N_0 \in \mathbb{Z}^+$ and numbers $a_j \in \{-1, 0, 1\}$, $1 \leq j \leq N_0$, such that*

$$\max_{s \in K} \left| \sum_{j=1}^{N_0} a_j j^{-s} - f(s) \right| < \varepsilon.$$

Given $N \in \mathbb{Z}^+$, $L \in X_n$ and $s \in \mathbb{C}$, we define

$$P_N(L, s) := \sum_{\mathcal{V}_j(L) \leq N} \mathcal{V}_j(L)^{-s} - \sum_{j=1}^N j^{-s} + \zeta(s).$$

Clearly P_N is analytic in \mathbb{C} except for a simple pole at $s = 1$ with residue 1.

Remark 3.2. In the proof of Theorem 1.3 we find it convenient to subtract and add $\zeta(s)$ to $\mathcal{E}_n(L, s)$ (see (1.5)). The main reason is that $\mathcal{E}_n(L, s) - \zeta(s)$ is an entire

function and its generalized Dirichlet series representation converges in $\operatorname{Re} s > \frac{1}{2}$ for almost every $L \in X_n$ (cf. the discussion in and just above Lemma 3.4 below). On the other hand, the second copy of $\zeta(s)$ plays no essential role in this investigation. We note that $P_N(L, s)$ is simply the sum of $\zeta(s)$ and a suitable truncation of $\mathcal{E}_n(L, s) - \zeta(s)$ that is easy to work with in connection with the approximation of Dirichlet polynomials (see the proof of Proposition 3.3 below).

The following approximation result is the key technical ingredient in the proof of Theorem 1.3.

Proposition 3.3. *Let Ω be a simply connected region in D that is symmetric with respect to the real axis. Then, for any $\varepsilon > 0$, any compact set $K \subset \Omega$ and any holomorphic function f on Ω that is real-valued on $\Omega \cap \mathbb{R}$, there exist $N \in \mathbb{Z}^+$ and a constant $\delta > 0$ such that*

$$\liminf_{n \rightarrow \infty} \operatorname{Prob}_{\mu_n} \left\{ L \in X_n : \max_{s \in K} |P_M(L, s) - f(s)| < \varepsilon \right\} > \delta \quad (3.1)$$

holds for any fixed $M \geq N$.

Proof. Let $h(s) := \zeta(s) - f(s)$ and note that h is holomorphic on Ω and real-valued on $\Omega \cap \mathbb{R}$. By Lemma 3.1, there exist $N_0 \in \mathbb{Z}^+$ and coefficients $a_j \in \{-1, 0, 1\}$, $1 \leq j \leq N_0$, such that

$$\max_{s \in K} \left| \sum_{j=1}^{N_0} a_j j^{-s} - h(s) \right| < \frac{\varepsilon}{3}. \quad (3.2)$$

We let $N > \max(N_0, 5)$ be an integer and set

$$b_j = \begin{cases} 1 - a_j & \text{if } 1 \leq j \leq N_0, \\ 1 & \text{if } N_0 < j \leq N. \end{cases}$$

Hence

$$P_N(L, s) - f(s) = \sum_{\mathcal{V}_j(L) \leq N} \mathcal{V}_j(L)^{-s} - \sum_{j=1}^N b_j j^{-s} + \left(h(s) - \sum_{j=1}^{N_0} a_j j^{-s} \right). \quad (3.3)$$

In addition it is useful to note that

$$\sum_{j=1}^N b_j j^{-s} = \sum_{j=1}^N n_j^{-s},$$

for some $N - N_0 \leq \mathcal{N} \leq N + N_0$ and a certain nondecreasing sequence $\{n_j\}_{j=1}^{\mathcal{N}}$ of positive integers. Note in particular that $n_{\mathcal{N}} = N$.⁷

⁷It is in principle straightforward, but notationally impractical, to write down explicit formulas for all the integers n_j .

Next, we find that there exist $N_1 \in \mathbb{Z}^+$ and $\delta_0 > 0$ (depending on K, N and ε) such that for $n \geq N_1$, we have

$$\text{Prob}_{\mu_n} \left\{ L \in X_n : \max_{s \in K} \left| \sum_{\mathcal{V}_j(L) \leq N} \mathcal{V}_j(L)^{-s} - \sum_{j=1}^N n_j^{-s} \right| < \frac{\varepsilon}{3} \right\} > \delta_0. \quad (3.4)$$

We recall that $n_N = N$ and define $n_{N+1} := N + 1$. It is clear that (3.4) follows if, for a sufficiently small $0 < \theta < 1$ (depending on K, N and ε),

$$n_j - \theta < \mathcal{V}_j(L) < n_j \quad (j = 1, \dots, N+1)$$

holds with probability $> \delta_0$ whenever $n \geq N_1$. Hence, it is sufficient to prove that

$$\mathcal{V}_1(L) \in I_1 := \left(n_1 - \theta, n_1 - \frac{N\theta}{N+1} \right), \quad (3.5)$$

and that

$$\mathcal{V}_{j+1}(L) - \mathcal{V}_j(L) \in I_{j+1} := \left(n_{j+1} - n_j, n_{j+1} - n_j + \frac{\theta}{N+1} \right) \quad (3.6)$$

holds for $1 \leq j \leq N$, with the given probability for all large enough n . To prove this, we first note that it follows from Theorem 2.1 and [12, Section 4.1] that

$$\{\mathcal{V}_1(L)\} \cup \{\mathcal{V}_{j+1}(L) - \mathcal{V}_j(L)\}_{j=1}^N$$

tend in distribution, as $n \rightarrow \infty$, to a collection $\{Y_j\}_{j=1}^{N+1}$ of independent exponentially distributed random variables of mean 1. The probability that (Y_1, \dots, Y_{N+1}) lies in the open set $\prod_{j=1}^{N+1} I_j$ may be explicitly calculated as

$$\delta_0^* := \left(\exp \left(\frac{\theta}{N+1} \right) - 1 \right)^{N+1} e^{-N-1} > 0.$$

Hence, it follows from [2, Theorem 2.1] that the lower limit as $n \rightarrow \infty$ of the probability that (3.5) and (3.6) holds for $1 \leq j \leq N$ is greater than or equal to δ_0^* . We conclude that (3.4) holds for any $\delta_0^* > \delta_0 > 0$ and all $n \geq N_1$ (where N_1 depends on δ_0).

Note that in the special case where $M = N$ and $\delta = \delta_0$, the inequality (3.1) follows from (3.2), (3.3), (3.4) and an application of the triangle inequality. It is also clear from the argument above that we can increase N as much as we like at the cost of having a possibly smaller constant δ_0 . It remains to show that (3.1) holds with the same right-hand side δ for any fixed $M \geq N$. Hence, for $M \geq N$, we study the finite sums

$$\begin{aligned} Q_{N,M}(L, s) &:= P_M(L, s) - P_N(L, s) \\ &= \sum_{N < \mathcal{V}_j(L) \leq M} \mathcal{V}_j(L)^{-s} - \sum_{j=N+1}^M j^{-s} = \int_N^M t^{-s} dS_{n,L}(t), \end{aligned}$$

where $S_{n,L}(t)$ is defined in (2.4). Let $\sigma_0 = \min\{\text{Re}(s) : s \in K\}$ and $\eta = (\sigma_0 - \frac{1}{2})/2$.

It follows from Corollary 2.3 with $C = 2$ that for each $n \geq 3$ there exists a set $Y_n \subset X_n$ with $\mu_n(Y_n) \geq \frac{1}{2}$ and such that for all lattices $L \in Y_n$ and all $t \geq 5$ we have $|S_{n,L}(t)| \ll_\eta t^{\frac{1}{2}+\eta}$. Here it is important to note that the implied constant is independent of both n and L . Integrating by parts we get, for any $L \in Y_n$ and all $M \geq N$,

$$\begin{aligned} |Q_{N,M}(L, s)| &= \left| \int_N^M t^{-s} dS_{n,L}(t) \right| \\ &= \left| [t^{-s} S_{n,L}(t)]_N^M + s \int_N^M t^{-s-1} S_{n,L}(t) dt \right| \ll_{\eta, K} N^{\frac{1}{2}+\eta-\sigma_0}, \end{aligned}$$

uniformly for all $s \in K$. Since we can make the right-hand side above as small as we desire by increasing N as needed, we obtain, for any $M \geq N$ and $n \geq 3$,

$$\text{Prob}_{\mu_n} \left\{ L \in X_n : \max_{s \in K} |Q_{N,M}(L, s)| < \frac{\varepsilon}{3} \right\} \geq \frac{1}{2}. \quad (3.7)$$

Finally, given $M \geq N$, we would like to use (3.2), (3.3), (3.4) and (3.7) to conclude that (3.1) holds with (say) $\delta = \delta_0/4$. In order to verify that this is possible, we recall that Theorem 2.1 states that the volumes $\{\mathcal{V}_j(L)\}_{j=1}^\infty$ tend in distribution (as $n \rightarrow \infty$) to the points of a Poisson process \mathcal{P} on the positive real line of constant intensity 1. Note that the proof of (3.4) only uses the restriction of these processes to the open interval $(0, N)$ and similarly that the proof of (3.7) only uses the restriction of these processes to (N, ∞) . Now the crucial observation is that the process \mathcal{P} may be realized as a union of a Poisson process on $(0, N)$ and an *independent* Poisson process on (N, ∞) , both of intensity 1 (see, e.g., [12, Section 2.2]). Hence it follows from (3.2), (3.3), (3.4) and (3.7), that when $n \geq \max(N_1, 3)$ (where we might need to increase N_1 depending on M) we have

$$\text{Prob}_{\mu_n} \left\{ L \in X_n : \max_{s \in K} |P_M(L, s) - f(s)| < \varepsilon \right\} > \frac{\delta_0}{4}. \quad (3.8)$$

In fact, as $n \rightarrow \infty$ we have the above inequality with any right-hand side strictly smaller than $\frac{\delta_0}{2}$. Since the lower bound in (3.8) holds for any fixed $M \geq N$, the proof is complete. \square

Next we define, for $N \in \mathbb{Z}^+$ and $\text{Re}(s) > 1$, the function

$$Q_N(L, s) := \sum_{\mathcal{V}_j(L) > N} \mathcal{V}_j(L)^{-s} - \sum_{j=N+1}^{\infty} j^{-s} = \int_N^\infty t^{-s} dS_{n,L}(t),$$

and note that this integral representation is holomorphic in the half-plane $\text{Re}(s) > \frac{1}{2}$ for almost every $L \in X_n$.

Lemma 3.4. *Let $n \geq 3$. Let $\varepsilon, \delta > 0$ and let K be a compact subset of the half-plane $\operatorname{Re}(s) > \frac{1}{2}$. Then there exists $M \in \mathbb{Z}^+$ such that*

$$\operatorname{Prob}_{\mu_n} \left\{ L \in X_n : \max_{s \in K} \left| \int_{M'}^{\infty} t^{-s} dS_{n,L}(t) \right| < \varepsilon \right\} \geq 1 - \delta$$

for all $M' \geq M$.

Proof. Let $\delta > 0$ be given. Recall from the proof of Proposition 3.3 that $\sigma_0 = \min\{\operatorname{Re}(s) : s \in K\}$ and $\eta = (\sigma_0 - \frac{1}{2})/2$. It follows from Corollary 2.3 that for each $n \geq 3$ there exists a set $Z_n \subset X_n$ with $\mu_n(Z_n) \geq 1 - \delta$ and such that for all lattices $L \in Z_n$ and all $t \geq 5$ we have $|S_{n,L}(t)| \ll_{\delta,\eta} t^{\frac{1}{2}+\eta}$, where the implied constant is independent of n and L . Now, for any $L \in Z_n$ and all $M \geq 5$, we get

$$\begin{aligned} \left| \int_M^{\infty} t^{-s} dS_{n,L}(t) \right| &= \left| \left[t^{-s} S_{n,L}(t) \right]_M^{\infty} + s \int_M^{\infty} t^{-s-1} S_{n,L}(t) dt \right| \\ &\ll_{\delta,\eta,K} M^{\frac{1}{2}+\eta-\sigma_0}, \end{aligned}$$

uniformly for all $s \in K$. Since we can make the right-hand side above as small as desired by choosing M large enough, the proof is complete. \square

Proof of Theorem 1.3. Let $\varepsilon > 0$ be given and let N and δ be given by Proposition 3.3 applied with $\varepsilon/2$, K and f . Let furthermore M be given by Lemma 3.4 applied with $\varepsilon/2$, $\delta/2$ and K . (Note that we without loss of generality may assume that $M \geq N$.) Hence, by Proposition 3.3 and Lemma 3.4, we find that

$$\begin{aligned} |\mathcal{E}_n(L, s) - f(s)| &= |\mathcal{E}_n(L, s) - \zeta(s) + \zeta(s) - f(s)| \\ &\leq |P_M(L, s) - f(s)| + |Q_M(L, s)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \end{aligned}$$

holds, in the limit as $n \rightarrow \infty$, with a probability of at least $\delta/2$. This finishes the proof. \square

4. Proof of Theorem 1.10

The general strategy for the proof of Theorem 1.10 is the same as in the proof of Theorem 1.3. The main difference is that Lemma 3.1 is no longer at our disposal. The following approximation lemma will take its place in the present argument.

Lemma 4.1. *Let Ω be a simply connected region in \mathbb{C} that is symmetric with respect to the real axis. Then, for any $\varepsilon > 0$, any compact set $K \subset \Omega$ and any holomorphic function f on Ω that is real-valued on $\Omega \cap \mathbb{R}$, there exists a Dirichlet polynomial*

$$A(s) = \sum_{k=1}^{N_0} a_k \lambda_k^{-s}$$

with integer coefficients such that

$$\max_{s \in K} |A(s) - f(s)| < \varepsilon.$$

Proof. The conditions on f and Ω are sufficient to ensure that

$$f(s) = \overline{f(\bar{s})} \quad (s \in \Omega). \quad (4.1)$$

Let $K^\# = K \cup \bar{K}$, where \bar{K} is the reflection of K in the real axis. By Runge's theorem (applied to a simply connected compact set K' satisfying $K^\# \subset K' \subset \Omega$) there exists a polynomial $P(s)$ such that

$$|P(s) - f(s)| < \frac{\varepsilon}{2} \quad (s \in K^\#).$$

By the symmetry of $K^\#$, we have

$$|\overline{P(\bar{s})} - \overline{f(\bar{s})}| < \frac{\varepsilon}{2} \quad (s \in K^\#),$$

and by (4.1) and the triangle inequality it follows that

$$|p(s) - f(s)| < \frac{\varepsilon}{2} \quad (s \in K), \quad (4.2)$$

where

$$p(s) = \frac{P(s) + \overline{P(\bar{s})}}{2} = \frac{P(s) + \bar{P}(s)}{2}$$

is a polynomial with real coefficients.

We now proceed to approximate the polynomial $p(s)$ by Dirichlet polynomials of the desired type. It is sufficient to prove that

$$p_m(s) = b_m s(s-1) \cdots (s-m+1)$$

may be approximated by a Dirichlet polynomial $A_m(s)$, since $p(s)$ can be written as a sum of such polynomials, that is,

$$p(s) = \sum_{m=0}^M p_m(s)$$

for some nonnegative integer M and suitable real numbers $\{b_m\}_{m=0}^M$. Let

$$g(x) = b_m x^s.$$

It is clear that

$$g^{(m)}(x) = b_m s(s-1) \cdots (s-m+1) x^{s-m}.$$

Note that this derivative can be approximated up to any given accuracy by $\Delta_h^m g(x)$, where

$$\Delta_h g(x) := \frac{g(x + h) - g(x)}{h},$$

by choosing h small enough. Note also that

$$\Delta_h^m g(x) = \frac{b_m}{h^m} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} (x + jh)^s.$$

It follows that any such $\Delta_h^m g(x)$, where we choose $x = 1$ and h such that b_m/h^m is an integer, will be of the desired type and can be chosen as our Dirichlet polynomial $A_m(s)$. Thus, by choosing h sufficiently small, we can ensure that

$$|A_m(s) - p_m(s)| < \frac{\varepsilon}{2(M+1)} \quad (0 \leq m \leq M, s \in K). \quad (4.3)$$

The result follows by choosing

$$A(s) = \sum_{m=0}^M A_m(s)$$

and applying the triangle inequality together with the inequalities (4.2) and (4.3). \square

Given $\Lambda \in \mathbb{R}^+$, two lattices $L_1, L_2 \in X_n$ and a complex number $s \in \mathbb{C}$, we define

$$\tilde{P}_\Lambda(L_1, L_2, s) := \sum_{\mathcal{V}_j(L_1) \leq \Lambda} \mathcal{V}_j(L_1)^{-s} - \sum_{\mathcal{V}_j(L_2) \leq \Lambda} \mathcal{V}_j(L_2)^{-s},$$

which clearly is an entire function of s .

Proposition 4.2. *Let Ω be a simply connected region in the half-plane $\operatorname{Re}(s) > \frac{1}{2}$ that is symmetric with respect to the real axis. Then, for any $\varepsilon > 0$, any compact set $K \subset \Omega$ and any holomorphic function f on Ω that is real-valued on $\Omega \cap \mathbb{R}$, there exist constants $\delta, \Lambda_0 \in \mathbb{R}^+$ such that*

$$\liminf_{n \rightarrow \infty} \operatorname{Prob}_{\mu_n \times \mu_n} \{ (L_1, L_2) \in X_n^2 : \max_{s \in K} |\tilde{P}_\Lambda(L_1, L_2, s) - f(s)| < \varepsilon \} > \delta \quad (4.4)$$

holds for any fixed $\Lambda \geq \Lambda_0$.

Proof. By Lemma 4.1, there exist an integer $N_0 \geq 1$ and sequences $\{\lambda_k\}_{k=1}^{N_0}$ and $\{a_k\}_{k=1}^{N_0}$ of positive real numbers and integers respectively, such that

$$\max_{s \in K} \left| \sum_{k=1}^{N_0} a_k \lambda_k^{-s} - f(s) \right| < \frac{\varepsilon}{3}. \quad (4.5)$$

Let $\lambda_0 = 5$, $C_1 = \sum_{a_k > 0} a_k$ and $C_2 = \sum_{a_k < 0} |a_k|$. By choosing $\Lambda_0 \geq \max\{\lambda_k : 0 \leq k \leq N_0\}$ sufficiently large, we find that there exist $N_1 \in \mathbb{Z}^+$ and $\delta_0 > 0$ (depending on K , Λ_0 and ε) such that for $n \geq N_1$, we have

$$\text{Prob}_{\mu_n \times \mu_n} \left\{ (L_1, L_2) \in X_n^2 : \max_{s \in K} \left| \tilde{P}_{\Lambda_0}(L_1, L_2, s) - \sum_{k=1}^{N_0} a_k \lambda_k^{-s} \right| < \frac{\varepsilon}{3} \right\} > \delta_0. \quad (4.6)$$

The proof of (4.6) is essentially the same as the proof of (3.4) (see the proof of Proposition 3.3). The only difference is that we in this case use the sequence $\{\mathcal{V}_j(L_1)\}_{j=1}^{C_1}$ to approximate the numbers λ_k with positive coefficients a_k and the sequence $\{\mathcal{V}_j(L_2)\}_{j=1}^{C_2}$ to approximate the numbers λ_k with negative coefficients a_k (in both cases with multiplicities according to the sizes of the corresponding $|a_k|$). Now, in the case where $\Lambda = \Lambda_0$ and $\delta = \delta_0$, the inequality (4.4) follows from (4.5), (4.6) and an application of the triangle inequality. Note that we may increase Λ_0 at the cost of having a possibly smaller constant δ_0 .

In order to show that (4.4) holds with the same right-hand side δ for any fixed $\Lambda \geq \Lambda_0$, we study the finite sums

$$\begin{aligned} \tilde{Q}_{\Lambda_0, \Lambda}(L_1, L_2, s) &:= \tilde{P}_{\Lambda}(L_1, L_2, s) - \tilde{P}_{\Lambda_0}(L_1, L_2, s) \\ &= \sum_{\Lambda_0 < \mathcal{V}_j(L_1) \leq \Lambda} \mathcal{V}_j(L_1)^{-s} - \sum_{\Lambda_0 < \mathcal{V}_j(L_2) \leq \Lambda} \mathcal{V}_j(L_2)^{-s} \\ &= \int_{\Lambda_0}^{\Lambda} t^{-s} dR_{n, L_1}(t) - \int_{\Lambda_0}^{\Lambda} t^{-s} dR_{n, L_2}(t), \end{aligned}$$

where $R_{n, L}(t)$ is defined in (2.2). Let $\sigma_0 = \min\{\text{Re}(s) : s \in K\}$ and $\eta = (\sigma_0 - \frac{1}{2})/2$. It follows from Theorem 2.2 with $C = 2$ that for each $n \geq 3$ there exists a set $U_n \subset X_n$ with $\mu_n(U_n) \geq \frac{1}{2}$ and such that for all lattices $L \in U_n$ and all $t \geq 5$ we have $|R_{n, L}(t)| \ll_{\eta} t^{\frac{1}{2} + \eta}$. Integrating by parts we get, for all $L_1, L_2 \in U_n$ and all $\Lambda \geq \Lambda_0$,

$$\begin{aligned} |\tilde{Q}_{\Lambda_0, \Lambda}(L_1, L_2, s)| &\leq \left| \int_{\Lambda_0}^{\Lambda} t^{-s} dR_{n, L_1}(t) \right| + \left| \int_{\Lambda_0}^{\Lambda} t^{-s} dR_{n, L_2}(t) \right| \\ &\ll_{\eta, K} \Lambda_0^{\frac{1}{2} + \eta - \sigma_0}, \end{aligned} \quad (4.7)$$

uniformly for all $s \in K$. Thus, since we can make the right-hand side above as small as we like by a sufficient increase of Λ_0 , we obtain, for any $\Lambda \geq \Lambda_0$ and $n \geq 3$,

$$\text{Prob}_{\mu_n \times \mu_n} \left\{ (L_1, L_2) \in X_n^2 : \max_{s \in K} |\tilde{Q}_{\Lambda_0, \Lambda}(L_1, L_2, s)| < \frac{\varepsilon}{3} \right\} \geq \frac{1}{4}. \quad (4.8)$$

Finally, given $\Lambda \geq \Lambda_0$, we use (4.5), (4.6), (4.8) and Theorem 2.1 to conclude that when $n \geq \max(N_1, 3)$ (where we might need to increase N_1 depending on Λ) we have

$$\text{Prob}_{\mu_n \times \mu_n} \left\{ (L_1, L_2) \in X_n^2 : \max_{s \in K} |\tilde{P}_\Lambda(L_1, L_2, s) - f(s)| < \varepsilon \right\} > \frac{\delta_0}{8}. \quad (4.9)$$

This finishes the proof. \square

We define, for each $\Delta \in \mathbb{R}^+$ and $\text{Re}(s) > 1$, the function

$$\begin{aligned} \tilde{Q}_\Delta(L_1, L_2, s) &:= \sum_{\mathcal{V}_j(L_1) > \Delta} \mathcal{V}_j(L_1)^{-s} - \sum_{\mathcal{V}_j(L_2) > \Delta} \mathcal{V}_j(L_2)^{-s} \\ &= \int_{\Delta}^{\infty} t^{-s} dR_{n, L_1}(t) - \int_{\Delta}^{\infty} t^{-s} dR_{n, L_2}(t), \end{aligned}$$

and note that this integral representation is holomorphic in the half-plane $\text{Re}(s) > \frac{1}{2}$ for almost every $(L_1, L_2) \in X_n^2$.

Lemma 4.3. *Let $n \geq 3$. Let $\varepsilon, \delta > 0$ and let K be a compact subset of the half-plane $\text{Re}(s) > \frac{1}{2}$. Then there exists $\Lambda \in \mathbb{R}^+$ such that*

$$\begin{aligned} \text{Prob}_{\mu_n \times \mu_n} \left\{ (L_1, L_2) \in X_n^2 : \right. \\ \left. \max_{s \in K} \left| \int_{\Lambda'}^{\infty} t^{-s} dR_{n, L_1}(t) - \int_{\Lambda'}^{\infty} t^{-s} dR_{n, L_2}(t) \right| < \varepsilon \right\} \geq 1 - \delta \end{aligned}$$

for all $\Lambda' \geq \Lambda$.

Proof. The result follows from Theorem 2.2 applied with $C \geq 2/\delta$ to an estimate similar to the one in (4.7). \square

Proof of Theorem 1.10. Let $\varepsilon > 0$ be given and let Λ_0 and δ be given by Proposition 4.2 applied with $\varepsilon/2$, K , and f . Let furthermore Λ be given by Lemma 4.3 applied with $\varepsilon/2$, $\delta/2$ and K . (We may assume that $\Lambda \geq \Lambda_0$.) Then, it follows from Proposition 4.2 and Lemma 4.3 that

$$\begin{aligned} |\mathcal{E}_n(L_1, s) - \mathcal{E}_n(L_2, s) - f(s)| &\leq |\tilde{P}_\Lambda(L_1, L_2, s) - f(s)| + |\tilde{Q}_\Lambda(L_1, L_2, s)| \\ &< \varepsilon \end{aligned}$$

holds, in the limit as $n \rightarrow \infty$, with a probability of at least $\delta/2$. \square

5. Proof of the imaginary shift case (Theorem 1.11)

We will use Drungilas–Garunkštis–Kačėnas variant [4, Theorem 2.1] of a result of Gonek [9] for general Dirichlet series. To do this we need a mean square estimate in a half-plane.⁸

Theorem 5.1. *Let $n \geq 2$. Then, for $\sigma > \frac{3}{4}$ and for almost all $L \in X_n$, we have*

$$\int_1^T \left| E_n \left(L, \frac{n(\sigma + it)}{2} \right) \right|^2 dt \ll_{L,\sigma} T.$$

By the definition of the normalized Epstein zeta function, the triangle inequality and the well known mean square estimate for the Riemann zeta function in the half-plane $\sigma > \frac{1}{2}$ (see e.g. Ivić [11]), it is sufficient to prove that

$$\int_1^T |\mathcal{E}_n(L, \sigma + it) - \zeta(\sigma + it)|^2 dt \ll_{L,\sigma} T. \quad (5.1)$$

We will prove this estimate using Gallagher's lemma [7, Lemma 1].

Lemma 5.2 (Gallagher). *Let $\delta = \theta/T$, with $0 < \theta < 1$, and let*

$$S(t) := \sum_k c_k e(v_k t)$$

be an absolutely convergent exponential sum having only real frequencies v_k . Then

$$\int_{-T}^T |S(t)|^2 dt \ll_{\theta} \int_{-\infty}^{\infty} \left| \delta^{-1} \sum_{x \leq v_k \leq x+\delta} c_k \right|^2 dx.$$

We will not be able to apply Gallagher's lemma directly, since it assumes that the exponential sum $S(t)$ is absolutely convergent. Thus, we will need to truncate our Dirichlet series; in fact, we will apply the following weak approximate functional equation.

Lemma 5.3. *Let $\varepsilon > 0$. Let $n \geq 2$ and assume that $\operatorname{Re}(s) > \frac{1}{2}$. Then, for any $X \geq 5$ and almost every $L \in X_n$, we have*

$$\mathcal{E}_n(L, s) - \zeta(s) = \sum_{v_j(L) \leq X} \mathcal{V}_j(L)^{-s} - \sum_{1 \leq j \leq X} j^{-s} + O_{L,\varepsilon}(|s|X^{\frac{1}{2}-\operatorname{Re}(s)+\varepsilon}).$$

Proof. The lemma follows immediately from Lemma 3.4 and its proof (using the estimate in Remark 2.4 in place of Corollary 2.3 in the case $n = 2$). \square

⁸We note here that Blomer [3] recently stated a mean square estimate on the critical line $\frac{n}{4} + it$ for almost all $L \in X_n$. Using this bound instead of our Theorem 5.1 in the proof of Theorem 1.11 would result in a stronger universality result valid in $D = \{s \in \mathbb{C} : \frac{1}{2} < \operatorname{Re}(s) < 1\}$.

Proof of Theorem 1.13. The theorem is a direct consequence of Lemma 5.3, the functional equation (1.3) and the Phragmén–Lindelöf principle. \square

In order to apply Gallagher's lemma, we also need estimates for short partial sums of $\mathcal{E}_n(L, s) - \zeta(s)$. We define

$$\Delta(x) = \Delta_{\sigma, L, X}(x) = \Delta_1(x) - \Delta_2(x), \quad (5.2)$$

where

$$\Delta_1(x) = \Delta_{\sigma, L, X; 1}(x) = \sum_{\mathcal{V}_j(L) \leq \min(e^x, X)} \mathcal{V}_j(L)^{-\sigma}$$

and

$$\Delta_2(x) = \Delta_{\sigma, L, X; 2}(x) = \sum_{1 \leq j \leq \min(e^x, X)} j^{-\sigma}.$$

To be precise, we need the following result.

Lemma 5.4. *Let $\varepsilon > 0$. Let further $n \geq 2$, $\sigma > \frac{1}{2}$, $0 \leq \delta \leq 1$ and $\alpha \in \mathbb{R}$. Then, for almost all $L \in X_n$, we have*

$$\int_{\alpha}^{\alpha+1} |\Delta_{\sigma, L, X}(x + \delta) - \Delta_{\sigma, L, X}(x)|^2 dx \ll_{L, \sigma, \varepsilon} \delta e^{\alpha(\frac{3}{2} - 2\sigma + \varepsilon)}.$$

Proof. To begin, we have

$$\begin{aligned} & \int_{\alpha}^{\alpha+1} |\Delta(x + \delta) - \Delta(x)|^2 dx \\ & \leq \max_{x \in [\alpha, \alpha+1]} |\Delta(x + \delta) - \Delta(x)| \int_{\alpha}^{\alpha+1} |\Delta(x + \delta) - \Delta(x)| dx. \end{aligned}$$

By Corollary 2.3 (or Remark 2.4 in the case $n = 2$) and integration by parts, we obtain

$$|\Delta(x + \delta) - \Delta(x)| = \left| \int_{\min(e^x, X)}^{\min(e^{x+\delta}, X)} t^{-\sigma} dS_{n, L}(t) \right| \ll_{L, \sigma, \varepsilon} e^{x(\frac{1}{2} - \sigma + \varepsilon)}$$

for almost all $L \in X_n$. For the rest of this proof we restrict our attention to the set of lattices satisfying the above estimate, for which we have

$$\int_{\alpha}^{\alpha+1} |\Delta(x + \delta) - \Delta(x)|^2 dx \ll_{L, \sigma, \varepsilon} e^{\alpha(\frac{1}{2} - \sigma + \varepsilon)} \int_{\alpha}^{\alpha+1} |\Delta(x + \delta) - \Delta(x)| dx. \quad (5.3)$$

Using (5.2) and the triangle inequality, we find that the latter integral is bounded by

$$\sum_{i=1}^2 \int_{\alpha}^{\alpha+1} |\Delta_i(x + \delta) - \Delta_i(x)| dx. \quad (5.4)$$

Since $\Delta_i(t)$ is nondecreasing for $i = 1, 2$, the absolute values in (5.4) can be removed and the above sum equals

$$\sum_{i=1}^2 \left(\int_{\alpha}^{\alpha+1} \Delta_i(x + \delta) dx - \int_{\alpha}^{\alpha+1} \Delta_i(x) dx \right).$$

Changing variables $y = x + \delta$ in the first integral, we obtain

$$\sum_{i=1}^2 \left(\int_{\alpha+\delta}^{\alpha+1+\delta} \Delta_i(y) dy - \int_{\alpha}^{\alpha+1} \Delta_i(x) dx \right) = \sum_{i=1}^2 \int_{\alpha}^{\alpha+\delta} (\Delta_i(x+1) - \Delta_i(x)) dx,$$

which by Corollary 2.3 (or Remark 2.4 in the case $n = 2$) is $O_{L,\sigma}(\delta e^{\alpha(1-\sigma)})$. Using this estimate, together with (5.3), we obtain the desired result. \square

Proof of Theorem 5.1. Let $\sigma > \frac{3}{4}$ be given. By the triangle inequality, the mean square estimate for the Riemann zeta function and Lemma 5.3 (applied with $X = T^4$) we have, for almost all $L \in X_n$,

$$\begin{aligned} & \int_1^T \left| E_n \left(L, \frac{n(\sigma + it)}{2} \right) \right|^2 dt \\ & \ll_{L,\sigma} \int_1^T \left| \sum_{\mathcal{V}_j(L) \leq T^4} \mathcal{V}_j(L)^{-s} - \sum_{1 \leq j \leq T^4} j^{-s} \right|^2 dt + T. \end{aligned} \quad (5.5)$$

The integral in the right-hand side of (5.5) can be estimated by an application of Lemma 5.2 with $\theta = \frac{1}{2}$, resulting in the upper bound

$$O(T^2) \int_{-\infty}^{\infty} \left| \Delta \left(x + \frac{1}{2T} \right) - \Delta(x) \right|^2 dx, \quad (5.6)$$

where $\Delta(x) = \Delta_{\sigma,L,X}(x)$ is defined in (5.2). The integrand has compact support contained in $[A, B]$, where $A = \log(\min(\mathcal{V}_1(L), 1)) + O(T^{-1})$ and $B = 4 \log T$. By dividing the integral into parts of length 1 and using Lemma 5.4 to estimate each term, we find that the expression in (5.6) is bounded by

$$O_{L,\sigma,\varepsilon}(T) \sum_{k=\lfloor A \rfloor}^{\lfloor B \rfloor} e^{k(\frac{3}{2} - 2\sigma + \varepsilon)},$$

where the geometric sum is bounded with respect to T whenever we choose $0 < \varepsilon < 2\sigma - \frac{3}{2}$. This finishes the proof. \square

Before we turn to the proof of Theorem 1.11, we introduce some more notation. We let \hat{L} denote a set containing one representative from each pair $\{\mathbf{m}, -\mathbf{m}\}$ of

primitive vectors⁹ in L , and write $\{\widehat{\mathcal{V}}_j(L)\}_{j=1}^\infty$ for the subsequence of $\{\mathcal{V}_j(L)\}_{j=1}^\infty$ corresponding to vectors $\mathbf{v} \in \widehat{L}$. We also let

$$\widehat{N}_{n,L}(t) := \#\{j : \widehat{\mathcal{V}}_j(L) \leq t\} \quad (t \geq 0),$$

and note that

$$N_{n,L}(t) = \sum_{1 \leq m \leq (t/\mathcal{V}_1(L))^{1/n}} \widehat{N}_{n,L}\left(\frac{t}{m^n}\right). \quad (5.7)$$

We will need a bound on the error term in the primitive circle problem, generalized to an n -dimensional ball of volume t and a generic lattice $L \in X_n$, of the same quality as the bounds in Theorem 2.2 and Remark 2.4.

Lemma 5.5. *Let $\varepsilon > 0$ and $n \geq 2$. Then, for almost all $L \in X_n$, we have*

$$\widehat{N}_{n,L}(t) = \frac{t}{\zeta(n)} + O_{L,\varepsilon}\left(t^{\frac{1}{2}+\varepsilon}\right). \quad (5.8)$$

Proof. We begin by defining (for $r \geq 0$)

$$\mathcal{N}_{n,L}(r) := \frac{1}{2} \#\{\mathbf{m} \in L \setminus \{\mathbf{0}\} : |\mathbf{m}| \leq \ell_1 r\}$$

and

$$\widehat{\mathcal{N}}_{n,L}(r) := \#\{\mathbf{m} \in \widehat{L} : |\mathbf{m}| \leq \ell_1 r\},$$

where ℓ_1 denotes the length of a shortest nonzero vector in L . We note that these counting functions are related to the ones in (5.7) by

$$N_{n,L}(t) = \mathcal{N}_{n,L}\left(\left(\frac{t}{\mathcal{V}_1(L)}\right)^{\frac{1}{n}}\right)$$

and

$$\widehat{N}_{n,L}(t) = \widehat{\mathcal{N}}_{n,L}\left(\left(\frac{t}{\mathcal{V}_1(L)}\right)^{\frac{1}{n}}\right).$$

Hence, writing $x = x(t, L) = (t/\mathcal{V}_1(L))^{1/n}$, we can reformulate (5.7) as

$$\mathcal{N}_{n,L}(x) = \sum_{1 \leq m \leq x} \widehat{\mathcal{N}}_{n,L}\left(\frac{x}{m}\right).$$

Now, using the Möbius inversion formula in the form given in [10, Theorem 268] together with Theorem 2.2 and Remark 2.4, we obtain

$$\widehat{N}_{n,L}(x) = \sum_{1 \leq m \leq x} \mu(m) \mathcal{N}_{n,L}\left(\frac{x}{m}\right) = \sum_{1 \leq m \leq x} \mu(m) \left(\frac{t}{m^n} + O_{L,\varepsilon}\left(\frac{t}{m^n}\right)^{\frac{1}{2}+\varepsilon}\right)$$

⁹A primitive vector (in a lattice L) is a non-zero lattice vector which is not a positive integral multiple of another lattice vector.

for almost every $L \in X_n$. Extending the upper bound in the summation of the main terms to infinity and then estimating the error terms trivially, we arrive at the desired result. \square

We further define

$$\hat{\mathcal{E}}_n(L, s) := \hat{\mathcal{V}}_1(L)^s \sum_{\mathbf{m} \in \hat{L}} \left(\frac{1}{2} V_n |\mathbf{m}|^n\right)^{-s} = 1 + \sum_{j=2}^{\infty} \left(\frac{\hat{\mathcal{V}}_j(L)}{\hat{\mathcal{V}}_1(L)}\right)^{-s} \quad (\operatorname{Re}(s) > 1)$$

and note that

$$E_n(L, \frac{ns}{2}) = 2\zeta(ns) \left(\frac{V_n}{2\hat{\mathcal{V}}_1(L)}\right)^s \hat{\mathcal{E}}_n(L, s). \quad (5.9)$$

Proof of Theorem 1.11. To begin, we note that since $\sigma_0 = \min\{\operatorname{Re}(s) : s \in K\} > \frac{3}{4} > \frac{1}{2}$, we can for any $\varepsilon_1 > 0$ choose $N_0 \in \mathbb{Z}^+$ so that for $n \geq 2$

$$\max_{\operatorname{Re}(s) \geq \sigma_0} \left| \prod_{p < N_0} (1 - p^{-ns}) - \frac{1}{\zeta(ns)} \right| < \varepsilon_1. \quad (5.10)$$

Furthermore, it follows from [21, Lemma 2] that the set

$$\{\log \hat{\mathcal{V}}_j(L) - \log \hat{\mathcal{V}}_1(L) : j \geq 2\}$$

is linearly independent over \mathbb{Q} for almost all lattices $L \in X_n$. From now on we restrict our attention to lattices $L \in X_n$ with this property. We let

$$\Gamma = \left\{ \frac{\hat{\mathcal{V}}_j(L)}{\hat{\mathcal{V}}_1(L)} : j \geq 2 \right\} \cup \{p^n : p < N_0\} \cup \left\{ \frac{V_n}{2\hat{\mathcal{V}}_1(L)} \right\}$$

and choose $\Gamma_{\max} \subseteq \Gamma$ to be maximal with respect to the property of having the logarithms of all its elements linearly independent over \mathbb{Q} . Since the differences $\log \hat{\mathcal{V}}_j(L) - \log \hat{\mathcal{V}}_1(L)$ for $j \geq 2$ are linearly independent over \mathbb{Q} by assumption, we will always include these elements in Γ_{\max} . We write

$$\Gamma_{\max} = \{\lambda_k\}_{k=1}^{\infty}.$$

Note that the logarithm of each element λ in the finite set $\Gamma \setminus \{\hat{\mathcal{V}}_j(L)/\hat{\mathcal{V}}_1(L)\}_{j=2}^{\infty}$ can be written as

$$\log \lambda = \sum_{k=1}^{N_2} q_{k,\lambda} \log \lambda_k = \sum_{k=1}^{N_2} \frac{h_{k,\lambda}}{N_1} \log \lambda_k \quad (5.11)$$

for $\lambda_k \in \Gamma_{\max}$, $q_{k,\lambda} \in \mathbb{Q}$, $h_{k,\lambda} \in \mathbb{Z}$ and $N_1, N_2 \in \mathbb{Z}^+$ where N_1 can be chosen as the least common multiple of the denominators of the rational numbers $q_{k,\lambda}$ (for all k

and λ). Note in particular that N_1 and N_2 are chosen independently of λ . This is possible since $\Gamma \setminus \{\widehat{\mathcal{V}}_j(L)/\widehat{\mathcal{V}}_1(L)\}_{j=2}^\infty$ is finite and the integers $h_{k,\lambda}$ are allowed to be zero. We also find it convenient to introduce

$$N_3 := \max_{\lambda} \sum_{k=1}^{N_2} |h_{k,\lambda}|.$$

Next, we define the Dirichlet series

$$A(s) := \sum_{k=1}^{N_2} \lambda_k^{-s/N_1} + \sum_{k=N_2+1}^{\infty} \lambda_k^{-s} \quad (\operatorname{Re}(s) > 1)$$

and the Dirichlet polynomials

$$B(s) := \sum_{k=1}^{N_2} (\lambda_k^{-s/N_1} - \lambda_k^{-s})$$

and

$$C(s) := \sum_{\lambda_k \in \Gamma_{\max} \setminus \{\widehat{\mathcal{V}}_j(L)/\widehat{\mathcal{V}}_1(L)\}_{j=2}^\infty} \lambda_k^{-s}.$$

Note that these functions are related to $\widehat{\mathcal{E}}_n(L, s)$ via

$$\widehat{\mathcal{E}}_n(L, s) = 1 + A(s) - B(s) - C(s). \quad (5.12)$$

We are interested in applying [4, Theorem 2.1] to the Dirichlet series $A(s)$. Hence we need to verify that $A(s)$ satisfies the four conditions in [4, Theorem 2.1] for almost all $L \in X_n$. The first condition, the so-called packing condition, follows from (5.8) which implies that for almost every $L \in X_n$, we have

$$|\widehat{N}_{n,L}(e^{x \pm \frac{c}{x^2}}) - \widehat{N}_{n,L}(e^x)| \gg \frac{e^x}{x^2} \gg e^{(1-\varepsilon')x}$$

for any $c > 0$ and any $\varepsilon' > 0$ (which in turn immediately implies the packing condition for $A(s)$). The second condition, the linear independence over \mathbb{Q} of the logarithms $\log \lambda_k$ corresponding to elements $\lambda_k \in \Gamma_{\max}$, follows from the defining property of Γ_{\max} . The third condition follows immediately from the fact that $\widehat{\mathcal{E}}_n(L, s)$ is absolutely convergent in the half-plane $\operatorname{Re}(s) > 1$. Finally, the fourth condition (on approximation) follows for almost all $L \in X_n$ from (5.9), the mean square estimate in Theorem 5.1 and [4, Proposition 2.2]. Thus, for almost all lattices $L \in X_n$, we may apply [4, Theorem 2.1] to $A(s)$. For the rest of this proof we assume that we are working with such a lattice $L \in X_n$.

Let

$$g(s) := \frac{f(s)}{2\zeta(ns)} \left(\frac{V_n}{2\widehat{\mathcal{V}}_1(L)} \right)^{-s} + B(s) + C(s) - 1,$$

where $f(s)$ is the function given in the statement of the theorem, and let

$$\beta := \sup_{\operatorname{Re}(s) \in [\frac{3}{4}, 1]} \left| 2\zeta(ns) \left(\frac{V_n}{2\hat{V}_1(L)} \right)^s \right| = \max_{x \in [\frac{3}{4}, 1]} \left| 2\zeta(nx) \left(\frac{V_n}{2\hat{V}_1(L)} \right)^x \right|.$$

Then we can apply [4, Theorem 2.1], with suitable δ and μ , to $A(s)$ and $g(s)$ to obtain, for a positive proportion of $t \in [0, T]$ as $T \rightarrow \infty$,

$$\max_{s \in K} |A(s + it) - g(s)| < \frac{\varepsilon}{4\beta} \quad (5.13)$$

and

$$\left\| \frac{t \log \lambda_k}{2\pi N_1} \right\| < \frac{\delta}{N_3} \quad (5.14)$$

for $1 \leq k \leq N_2$, where $\|\cdot\|$ denotes the distance to the closest integer. Note that it follows from (5.11), (5.14) and the triangle inequality that

$$\left\| \frac{t \log(p^n)}{2\pi} \right\| < \delta$$

for all $p < N_0$ and

$$\left\| \frac{t \log(V_n/(2\hat{V}_1(L)))}{2\pi} \right\| < \delta.$$

Hence, using (5.10) and the absolute convergence of $\zeta(s)$ in $\operatorname{Re}(s) > 1$, we obtain, for all sufficiently small choices of ε_1 and δ ,

$$\max_{s \in K} \left| \frac{(V_n/(2\hat{V}_1(L)))^{-s}}{2\zeta(ns)} - \frac{(V_n/(2\hat{V}_1(L)))^{-s-it}}{2\zeta(n(s+it))} \right| < \frac{\varepsilon}{4\beta(\max_{s \in K} |f(s)| + 1)}. \quad (5.15)$$

Similarly, we use (5.14) with a possibly smaller constant δ to conclude that

$$\max_{s \in K} |B(s) - B(s + it)| < \frac{\varepsilon}{4\beta} \quad \text{and} \quad \max_{s \in K} |C(s) - C(s + it)| < \frac{\varepsilon}{4\beta}. \quad (5.16)$$

We stress that (5.15) and (5.16) holds for the same values of t as (5.13) and (5.14). Throughout, we restrict our attention to such values of t .

In order to finish the proof it remains to put the above pieces together. Using (5.12), (5.13) and (5.16), together with the triangle inequality, we obtain

$$\begin{aligned} \max_{s \in K} \left| \hat{\mathcal{E}}_n(L, s + it) - \frac{f(s)}{2\zeta(ns)} \left(\frac{V_n}{2\hat{V}_1(L)} \right)^{-s} \right| &\leq \max_{s \in K} |A(s + it) - g(s)| \\ &+ \max_{s \in K} |B(s) - B(s + it)| + \max_{s \in K} |C(s) - C(s + it)| < \frac{3\varepsilon}{4\beta}. \end{aligned}$$

Next, it follows from the above and (5.15) that

$$\begin{aligned} \max_{s \in K} \left| \widehat{\mathcal{E}}_n(L, s + it) - \frac{f(s)}{2\zeta(n(s + it))} \left(\frac{V_n}{2\widehat{\mathcal{V}}_1(L)} \right)^{-s-it} \right| \\ \leq \max_{s \in K} \left| \widehat{\mathcal{E}}_n(L, s + it) - \frac{f(s)}{2\zeta(ns)} \left(\frac{V_n}{2\widehat{\mathcal{V}}_1(L)} \right)^{-s} \right| \\ + \max_{s \in K} |f(s)| \max_{s \in K} \left| \frac{(V_n/(2\widehat{\mathcal{V}}_1(L)))^{-s}}{2\zeta(ns)} - \frac{(V_n/(2\widehat{\mathcal{V}}_1(L)))^{-s-it}}{2\zeta(n(s + it))} \right| < \frac{\varepsilon}{\beta}. \end{aligned}$$

Finally, we use (5.9) to arrive at

$$\begin{aligned} \max_{s \in K} \left| E_n \left(L, \frac{n(s + it)}{2} \right) - f(s) \right| \\ \leq \beta \max_{s \in K} \left| \widehat{\mathcal{E}}_n(L, s + it) - \frac{f(s)}{2\zeta(n(s + it))} \left(\frac{V_n}{2\widehat{\mathcal{V}}_1(L)} \right)^{-s-it} \right| < \varepsilon, \end{aligned}$$

which is the desired conclusion. \square

References

- [1] B. Bagchi, The statistical behaviour and universality properties of the Riemann zeta function and other allied Dirichlet series, Ph.D. thesis, Indian Statistical Institute, Calcutta, 1981.
- [2] P. Billingsley, *Convergence of probability measures*. Second edition, Wiley Series in Probability and Statistics, John Wiley & Sons Inc., New York, 1999. Zbl 0944.60003 MR 1700749
- [3] V. Blomer, Epstein zeta-functions, subconvexity, and the purity conjecture, *J. Inst. Math. Jussieu*, (2018), 1–16.
- [4] P. Drungilas, R. Garunkštis, and A. Kačėnas, Universality of the Selberg zeta-function for the modular group, *Forum Math.*, **25** (2013), no. 3, 533–564. Zbl 1328.11093 MR 3062863
- [5] P. Epstein, Zur Theorie allgemeiner Zetafunctionen, *Math. Ann.*, **56** (1903), no. 4, 615–644. Zbl 34.0461.02 MR 1511190
- [6] P. Epstein, Zur Theorie allgemeiner Zetafunktionen. II, *Math. Ann.*, **63** (1906), no. 2, 205–216. Zbl 37.0433.02 MR 1511399
- [7] P. X. Gallagher, A large sieve density estimate near $\sigma = 1$, *Invent. Math.*, **11** (1970), 329–339. Zbl 0219.10048 MR 279049
- [8] D. Goldfeld, *Automorphic forms and L-functions for the group $GL(n, \mathbb{R})$* . With an appendix by Kevin A. Broughan, Cambridge Studies in Advanced Mathematics, 99, Cambridge University Press, Cambridge, 2006. Zbl 1108.11003 MR 2254662
- [9] S. M. Gonek, Analytic properties of zeta and L-functions, Ph.D. thesis, University of Michigan, Ann Arbor, 1979. MR 2628587

- [10] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*. Sixth edition. Revised by D. R. Heath-Brown and J. H. Silverman. With a foreword by Andrew Wiles, Oxford University Press, Oxford, 2008. Zbl 1159.11001 MR 2445243
- [11] A. Ivić, *The Riemann zeta-function. Theory and applications*. Reprint of the 1985 original (Wiley, New York), Dover Publications Inc., Mineola, NY, 2003. Zbl 1034.11046 MR 1994094
- [12] J. F. C. Kingman, *Poisson processes*, Oxford Studies in Probability, 3, The Clarendon Press, Oxford University Press, New York, 1993. Zbl 0771.60001 MR 1207584
- [13] A. A. Karatsuba and S. M. Voronin, *The Riemann zeta-function*. Translated from the Russian by Neal Koblitz, de Gruyter Expositions in Mathematics, 5, Walter de Gruyter & Co., Berlin, 1992. Zbl 0756.11022 MR 1183467
- [14] K. Matsumoto, A survey on the theory of universality for zeta and L -functions, in *Number theory*, 95–144, Ser. Number Theory Appl., 11, World Sci. Publ., Hackensack, NJ, 2015. Zbl 1381.11070 MR 3382056
- [15] H. Mishou and H. Nagoshi, Functional distribution of $L(s, \chi_d)$ with real characters and denseness of quadratic class numbers, *Trans. Amer. Math. Soc.*, **358** (2006), no. 10, 4343–4366. Zbl 1160.11041 MR 2231380
- [16] H. Mishou and H. Nagoshi, The universality of quadratic L -series for prime discriminants, *Acta Arith.*, **123** (2006), no. 2, 143–161. Zbl 1154.11024 MR 2232505
- [17] W. M. Schmidt, A metrical theorem in geometry of numbers, *Trans. Amer. Math. Soc.*, **95** (1960), 516–529. Zbl 0101.27904 MR 117222
- [18] C. L. Siegel, A mean value theorem in geometry of numbers, *Ann. of Math.* (2), **46** (1945), 340–347. Zbl 0063.07011 MR 12093
- [19] F. Spinu, The L^4 norm of the Eisenstein series, Ph.D. thesis, Princeton University, Princeton, 2003. MR 2705106
- [20] J. Steuding, *Value-distribution of L -functions*, Lecture Notes in Mathematics, 1877, Springer, Berlin, 2007. Zbl 1130.11044 MR 2330696
- [21] A. Strömgrensson and A. Södergren, On the location of the zero-free half-plane of a random Epstein zeta function, *Math. Ann.*, **371** (2018), no. 3–4, 1191–1227. Zbl 06923800 MR 3831269
- [22] A. Södergren, On the Poisson distribution of lengths of lattice vectors in a random lattice, *Math. Z.*, **269** (2011), no. 3–4, 945–954. Zbl 1257.60002 MR 2860272
- [23] A. Södergren, On the distribution of angles between the N shortest vectors in a random lattice, *J. London Math. Soc.* (2), **84** (2011), no. 3, 749–764. Zbl 1268.60068 MR 2855800
- [24] A. Södergren, On the value distribution and moments of the Epstein zeta function to the right of the critical strip, *J. Number Theory*, **131** (2011), no. 7, 1176–1208. Zbl 1228.11051 MR 2782836
- [25] A. Södergren, On the value distribution of the Epstein zeta function in the critical strip, *Duke Math. J.*, **162** (2013), no. 1, 1–48. Zbl 1285.11065 MR 3011871
- [26] A. Terras, The minima of quadratic forms and the behavior of Epstein and Dedekind zeta functions, *J. Number Theory*, **12** (1980), no. 2, 258–272. Zbl 0432.10010 MR 578821
- [27] S. M. Voronin, A theorem on the “universality” of the Riemann zeta-function, *Izv. Akad. Nauk SSSR Ser. Mat.*, **39** (1975), no. 3, 475–486 (Russian); *Math. USSR Izv.*, **9** (1975), no. 3, 443–453. Zbl 0315.10037 MR 472727

- [28] S. M. Voronin, A theorem on the distribution of values of the Riemann zeta-function, *Dokl. Akad. Nauk SSSR*, **221** (1975), no. 4, 771 (Russian); *Soviet Math. Dokl.*, **16** (1975), no. 2, 410. Zbl 0326.10035 MR 382191
- [29] S. M. Voronin, The zeros of zeta-functions of quadratic forms, Number theory, mathematical analysis and their applications, *Trudy Mat. Inst. Steklov.*, **142** (1976), 135-147, 269 (Russian). Zbl 0432.10009 MR 562285
- [30] S. M. Voronin, Analytic properties of Dirichlet generating functions of arithmetic objects, Ph.D. thesis, Steklov Mathematical Institute, Moscow, 1977 (Russian).

Received July 4, 2018

J. Andersson, Department of Mathematics, School of Science and Technology,
Örebro University, SE-701 82 Örebro, Sweden

E-mail: johan.andersson@oru.se

A. Södergren, Department of Mathematical Sciences, University of Copenhagen,
Universitetsparken 5, 2100 Copenhagen Ø, Denmark;
Current address: Department of Mathematical Sciences,
Chalmers University of Technology and the University of Gothenburg,
SE-412 96 Gothenburg, Sweden

E-mail: andesod@chalmers.se

