

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 95 (2020)
Heft: 3

Artikel: Hyperbolic surfaces with sublinearly many systoles that fill
Autor: Fortier Bourque, Maxime
DOI: <https://doi.org/10.5169/seals-882418>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Hyperbolic surfaces with sublinearly many systoles that fill

Maxime Fortier Bourque

Abstract. For any $\varepsilon > 0$, we construct a closed hyperbolic surface of genus $g = g(\varepsilon)$ with a set of at most εg systoles that fill, meaning that each component of the complement of their union is contractible. This surface is also a critical point of index at most εg for the systole function, disproving the lower bound of $2g - 1$ posited by Schmutz Schaller.

Mathematics Subject Classification (2010). 32G15, 30F60.

Keywords. Systole, moduli space, spine.

1. Introduction

The moduli space $\mathcal{M}_{g,n}$ of Riemann surfaces of genus g with n punctures is an object of great interest to many geometers and topologists. It encodes all the different complex structures, conformal structures, or hyperbolic structures (provided $2g + n > 2$) supported on a surface with given topology. The topology of moduli space is largely encoded in its orbifold fundamental group $\Gamma_{g,n}$, the mapping class group.

All the torsion-free finite index subgroups of $\Gamma_{g,n}$ have the same cohomological dimension, which is called the *virtual cohomological dimension* (vcd) of $\Gamma_{g,n}$. Harer computed the vcd of $\Gamma_{g,n}$ for all g and $n \geq 0$ and found a spine (a deformation retract) for $\mathcal{M}_{g,n}$ with this smallest possible dimension whenever $n > 0$ [9]. When $n = 0$, the vcd of the mapping class group is equal to $4g - 5$, but a spine of this dimension has yet to be found [4, Question 1]. The largest codimension attained so far is equal to 2 [10] (the space $\mathcal{M}_{g,0}$ has dimension $6g - 6$).

In an unpublished preprint [28], Thurston claimed that the set \mathcal{X}_g of closed hyperbolic surfaces of genus $g \geq 2$ whose systoles fill forms a spine for $\mathcal{M}_g = \mathcal{M}_{g,0}$. Recall that a *systole* is a closed geodesic of minimal length, and a set of curves *fills* if each component of the complement of their union is simply connected. Thurston's proof that \mathcal{M}_g deformation retracts onto \mathcal{X}_g appears to be difficult to complete [10]. Furthermore, the dimension of \mathcal{X}_g is still not known, mostly because we do not understand which filling sets of curves can be systoles. Indeed, Thurston writes:

Unfortunately, we do not have a combinatorial characterization of collections of curves which can be the collection of shortest geodesics on a surface. This seems like a challenging problem, and until more is understood about how to answer it, there are probably not many applications of the current result.

The paper [2] provides some partial answers to Thurston's question. On a closed hyperbolic surface, systoles do not self-intersect and distinct systoles can intersect at most once. This obvious necessary condition is, however, far from being sufficient. Indeed, a filling set of systoles must contain at least $\sim \pi g / \log g$ curves [2, Theorem 3], but there exist filling collections of $\sim 2\sqrt{g}$ geodesics pairwise intersecting at most once [2, Corollary 2]. There is a discrepancy in the opposite direction as well: a closed hyperbolic surface can have at most $Cg^2 / \log g$ systoles [16, Corollary 1.4], but there exist filling collections with more than g^2 geodesics pairwise intersecting at most once [14, Theorem 1.1].

Our main result here is a construction of closed hyperbolic surfaces with filling sets of systoles containing sublinearly many curves in terms of the genus. Compare this with [21, 22] where surfaces with superlinearly many systoles are found. Though we are still very far¹ from the lower bound of $\pi g / \log g$, our examples improve upon the previous record of surfaces with filling sets of $2g$ systoles ([2, Section 5], [18]).

Theorem 1.1. *For every $\varepsilon > 0$, there exist an integer $g \geq 2$ and a closed hyperbolic surface of genus g with a filling set of at most εg systoles.*

Near a surface with a filling set of at most εg systoles, Thurston's set \mathcal{X}_g contains the set of solutions to the same number of equations. This should imply that \mathcal{X}_g has codimension at most εg in \mathcal{M}_g . However, the equations requiring the curves to have equal length can be redundant, preventing us from applying the implicit function theorem. We only manage to prove that \mathcal{X}_g has dimension at least $4g - 5$ when g is even, but conjecture the following.

Conjecture 1.2. *For every $\varepsilon > 0$, there exists an integer $g \geq 2$ such that \mathcal{X}_g has dimension at least $(6 - \varepsilon)g$.*

On the other hand, we can prove that a closely related spine, the Morse–Smale complex for the systole function, has dimension much larger than the virtual cohomological dimension of the mapping class group.

In a series of papers [19, 20, 23, 24], Schmutz Schaller initiated the study of the systole function $\text{sys}: \mathcal{M}_{g,n} \rightarrow \mathbb{R}_+$, which records the length of any of the shortest closed geodesics on a surface. He proved that the systole function is a topological Morse function on the Teichmüller space $\mathcal{T}_{g,n}$ whenever $n > 0$ [24] and Akrouf extended this result to $n = 0$ (and to a more general class of functions) in [1].

Schmutz Schaller constructed a critical point of index $2g - 1$ for the systole function in every genus $g \geq 2$ and thought it was “quite possible” that this was

¹The genus g in Theorem 1.1 grows like a tower of exponentials of length roughly $1/\varepsilon$.

the smallest achievable index [24, p. 439]. He verified this hypothesis for $g = 2$ by finding all the critical points in \mathcal{M}_2 . If this were true in general, it would imply that the Morse–Smale complex for the systole function has the smallest possible dimension

$$4g - 5 = (6g - 6) - (2g - 1)$$

for a spine of \mathcal{M}_g . However, our surfaces show that no such inequality holds.

Theorem 1.3. *For every $\varepsilon > 0$, there exist an integer $g \geq 2$ and a critical point of index at most εg for the systole function on \mathcal{T}_g .*

Organization of the paper. The surfaces arising in Theorems 1.1 and 1.3 are built in two steps, in a similar fashion as the local maxima from [7]. First, in Section 2, we define a building block (depending on some parameters) which is a surface whose systoles are the boundary components. This surface is modelled on a flag-transitive surface map (a generalization of Platonic solids) and can be cut into isometric right-angled polygons along a collection of geodesic arcs. We then glue building blocks together according to the combinatorics of certain graphs of large girth with strong transitivity properties in Section 3. We do this in such a way that the boundaries of the blocks remain systoles in the larger surface and that the arcs in the blocks connect up to form systoles as well (see Section 4). In Section 5, we show that $X/\text{Isom}(X)$ is isometric to a triangle or a quadrilateral. This easily implies that X is a critical point of the systole function, which we prove in Section 6. Finally, we discuss our failed attempt to prove Conjecture 1.2 in Section 7.

Acknowledgements. I thank the anonymous referee for their useful comments and corrections.

2. Building blocks

Graphs. A *graph* is a 1-dimensional cell complex, where there can be multiple edges between two vertices and edges from vertices to themselves. The *valence* of a vertex in a graph is the number of half-edges adjacent to it. If every vertex in a graph has the same valence, then this number is called the valence of the graph.

Flag-transitive maps. A *map* M is a graph embedded on a surface S such that the closure of each complementary component is an embedded closed disk (called a *face* of the map). All maps considered in this paper will be *orientable*, meaning that the surface S is required to be orientable. If all the faces of a map M have the same number p of edges and all the vertices have the same valence q , then M is said to have *type* $\{p, q\}$.

A *flag* in a map is a triple consisting of a vertex v , an edge e containing v , and a face f containing e . A *map-automorphism* of M is an automorphism of the underlying graph which can be realized by a homeomorphism of the surface S . A map is *flag-transitive*² if its group of map-automorphisms acts transitively on flags.

Any flag-transitive map has type $\{p, q\}$ for some $p \geq 1$ and $q \geq 2$. The five Platonic solids are the only flag-transitive maps on the sphere with $p, q \geq 3$; their types are $\{3, 3\}$, $\{4, 3\}$, $\{3, 4\}$, $\{5, 3\}$, and $\{3, 5\}$. Beach balls assembled from q spherical bigons are flag-transitive maps of type $\{2, q\}$.

Maps of large girth. A *cycle* in a graph is a sequence of oriented edges (e_1, \dots, e_k) such that the endpoint of e_i coincides with the starting point of e_{i+1} for every $i \in \mathbb{Z}_k$. Cycles are considered up to cyclic permutation of their edges and reversal of orientation. The *length* of a cycle is the number of edges that it uses. A cycle is *non-trivial* if it cannot be homotoped to a point by deleting *backtracks*, that is, consecutive edges (modulo k) with opposite orientations. The *girth* of a graph is the length of any of its shortest non-trivial cycles. These shortest non-trivial cycles will be called *girth cycles*. A graph of girth at most 2 is often called a *multigraph*, and a graph of girth larger than 2 is *simple*.

The girth of a flag-transitive map M of type $\{p, q\}$ is at most p since the faces are non-trivial cycles of length p . If M is finite, then one can actually unwrap all the cycles shorter than p by taking a suitable finite normal cover, thereby obtaining a finite flag-transitive map N of girth p [6, Theorem 11]. That such covers exist follows from Mal'cev's theorem on the residual finiteness of finitely generated linear groups [13].

Theorem 2.1 (Evans). *For any $p, q \geq 2$, there exists a finite flag-transitive map of type $\{p, q\}$ and girth p .*

See also [15] and [27] for constructive proofs of this result.

Regular polygons. Let $q \geq 3$. Up to isometry, there exists a unique polygon P in the hyperbolic plane with $2q$ sides of the same length L and all interior angles equal to $\pi/2$. We will call P the *regular right-angled $2q$ -gon*. By connecting the center of P to the midpoint of a side and one of its vertices, we obtain a triangle with interior angles $\pi/2$, $\pi/4$, and $\pi/2q$ and a side of length $L/2$ from which we obtain the equation

$$\cosh(L/2) = \cos(\pi/2q) / \sin(\pi/4) = \sqrt{2} \cos(\pi/2q) \quad (2.1)$$

(see [5, p. 454]). We color the sides of P red and blue in such a way that adjacent sides have different colors.

²These maps are usually called *regular*, but if we stuck to standard terminology, this word would be used with five different meanings throughout the paper.

Lemma 2.2. *Any arc α between two disjoint sides in the regular right-angled $2q$ -gon P has length at least L , with equality only if α is a side of P .*

Proof. Let α have minimal length among such arcs. Then α must be geodesic and orthogonal to ∂P at its endpoints. These endpoints are separated by m sides of P in one direction and n sides in the other, where $m + n + 2 = 2q$ and $m \leq n$. First suppose that $m > 1$. Let d be a main diagonal of P which is linked with α and has one endpoint at an extremity of one of the two sides of P joined by α . Let z be the intersection point between α and d , and let α_{\pm} be the two components of $\alpha \setminus \{z\}$ labelled in such a way that α_+ and d have endpoints in a common side of P . If R_d denotes the reflection about d , then the arc $\gamma = \alpha_- \cup R_d(\alpha_+)$ has the same length as α and joins two disjoint sides of P (because $m > 1$). By minimality, γ must be geodesic and orthogonal to ∂P , which is absurd. This shows that $m = 1$, in which case α is a side of P (the orthogonal segment between two geodesics in the hyperbolic plane is unique when it exists). \square

One can also prove this using trigonometry (see [2, p. 91]).

Gluing regular polygons along maps. Let M be an oriented map of type $\{p, q\}$ where $q \geq 3$. Let P be the unique right-angled regular hyperbolic $2q$ -gon with sides colored red and blue as above. We now define a hyperbolic surface B modelled on M . For each vertex $v \in M$, take a copy P_v of P . The blue sides of P_v are labelled in counterclockwise order by the edges adjacent to v in M , which come with a cyclic ordering from the orientation. For each edge $e = \{u, v\}$ in M , we glue the polygons P_u and P_v along their sides labelled e by an orientation-reversing isometry. The resulting surface is denoted B and will be called a *block* in the sequel. The polygons $P_v \subset B$ are its *tiles*.

Topologically, B is the same as the surface $S \supset M$ with a hole cut out in each face. Indeed, if we join the center of each polygon P_v to the midpoints of its blue sides, we obtain an embedded copy of M in B . Since each P_v deformation retracts onto the star $M \cap P_v$, the surface B deformation retracts onto M . Each boundary component of B is the concatenation of p red sides of polygons P_v coming from the p vertices v around a face of M . In particular, each boundary component of B has length pL , where L is the positive number implicitly defined by Equation (2.1).

Lemma 2.3. *Let M be a map of type $\{p, q\}$ and girth p , where $p \geq 2$ and $q \geq 3$. Then the systoles in B are the boundary geodesics, of length pL .*

Proof. Let γ be a systole in B . As explained above, the map M embeds in B as the dual graph to the decomposition into the $2q$ -gons P_v . Let $\pi : B \rightarrow M$ be the nearest point projection. The image $\pi(\gamma)$ must be non-trivial in M since γ is non-trivial in B and π is a deformation retraction. It follows that the combinatorial length of $\pi(\gamma)$ in M is at least p . In other words, γ intersects at least p tiles P_v , joining distinct blue

sides each time. By Lemma 2.2, the length of $\gamma \cap P_v$ is at least L for any tile P_v that γ intersects. The total length of γ is therefore greater than or equal to pL . If equality occurs, then γ must be a concatenation of red arcs, that is, a boundary geodesic. \square

Corollary 2.4. *Let M be a map of type $\{p, q\}$ and girth p , where $p \geq 2$ and $q \geq 3$. Then any arc from a boundary component to itself in B which cannot be homotoped into ∂B has length strictly larger than $pL/2$.*

Proof. Suppose that α is a non-trivial arc of length at most $pL/2$ from a boundary geodesic b to itself. The arc α followed by the shorter of the two subarcs of b between its endpoints is a non-trivial closed curve γ of length at most pL in B . The closed geodesic homotopic to γ is strictly shorter, contradicting Lemma 2.3. \square

Lemma 2.5. *Let M be a map of type $\{p, q\}$ and girth p , where $p \geq 2$ and $q \geq 3$. Then any arc α from ∂B to ∂B which cannot be homotoped into ∂B has length at least L , with equality only if α is a blue arc.*

Proof. Let α be a geodesic arc from ∂B to ∂B . By Lemma 2.3, we may assume that α joins consecutive sides of any tile P_v it intersects. Since the starting point of α is on a red side, it has to next intersect a blue side, and then a red. This means that α is homotopic to a blue arc in a union $P_u \cup P_v$ of two adjacent tiles. This blue arc is shortest among all arcs in $P_u \cup P_v$ joining the same two sides, as it is orthogonal to the boundary at both endpoints. \square

The above results do not require the map M to be finite or flag-transitive, but we will impose these conditions in the next sections.

3. Gluing graphs

In this section, we explain how to glue blocks together along certain graphs of large girth with large automorphism groups in order to get closed hyperbolic surfaces with many symmetries and few systoles.

Strict polygonal graphs. A *strict polygonal graph* is a graph G such that any embedded path of length 2 in G is contained in a unique girth cycle (where cycles are considered up to cyclic reordering and reversal). This notion was introduced by Perkel in his thesis [17]. Examples of strict polygonal graphs include polygons, the tetrahedron, the dodecahedron, and the cube of any dimension. See [25] for a short survey on the subject.

Archdeacon and Perkel [3] found a way to double the girth of a strict polygonal graph G (or any graph) by taking an appropriate normal covering space. The girth cycles in this cover \tilde{G} are precisely those that wrap twice around a girth cycle in G under the covering map. Repeated applications of their construction yield strict

polygonal graphs of arbitrarily large girth and constant valence (equal to the valence of G).

Seress and Swartz [26, Theorem 3.2] proved that any automorphism of the base graph G lifts to an automorphism of the girth-doubling cover \tilde{G} . They concluded that if G is vertex transitive, edge transitive, arc transitive or 2-arc transitive, then so is \tilde{G} . We will need an even stronger transitivity property, described in the next paragraph.

Isotropic graphs. The *star* $\text{st}(v)$ of a vertex v in a graph is the set of half-edges adjacent to v . A graph G is *locally symmetric* if for every vertex $v \in V(G)$, any bijection of $\text{st}(v)$ can be extended to an automorphism of G that fixes v . We say that a graph is *isotropic* if it is vertex transitive and locally symmetric. To spell it out, G is isotropic if every injection $\text{st}(u) \hookrightarrow \text{st}(v)$ between stars in G extends to an automorphism of G .

In an isotropic graph, there is a girth cycle passing through any embedded path of length 2, but there can be more than one.

Example 3.1. The Petersen graph P (the quotient of the dodecahedron by the antipodal involution) is an isotropic graph of valence 3 and girth 5 on 10 vertices. However, P is not strict polygonal since every embedded path of length 2 is contained in two distinct girth cycles in P .

Lubotzky [12] constructed infinitely many isotropic Cayley graphs of any valence $d \geq 3$ and any even girth ≥ 6 (the generators are involutions, allowing the valence to be odd). Since we want better control on the girth cycles of our isotropic graphs, we use the girth-doubling construction of Archdeacon and Perkel instead. The proof that the girth-doubling cover \tilde{G} of a graph G is isotropic provided that G is isotropic follows immediately from [26, Theorem 3.2], which states that any automorphism of G lifts to \tilde{G} , and the fact that the covering $\tilde{G} \rightarrow G$ is normal, so that its deck group acts transitively on fibers.

The simplest isotropic strict polygonal graph is a pair of vertices joined by $d \geq 2$ edges. Repeated applications of the girth-doubling construction to this graph Θ yield a sequence of finite, isotropic, strict polygonal graphs of any valence and arbitrarily large girth.

Theorem 3.2 (Archdeacon–Perkel, Seress–Swartz). *For any $d \geq 2$ and $n \geq 1$, there exists a finite, isotropic, strict polygonal graph G of valence d and girth 2^n . In fact, G can be chosen to be a covering space of the bipartite graph Θ of valence d on 2 vertices, in which case the girth cycles in G project to powers of girth cycles in Θ under the covering map.*

Gluing. We now explain how to glue copies of the block B from Section 2 along a finite isotropic strict polygonal graph G to get a closed hyperbolic surface X with a small set of systoles that fill.

Let $q \geq 3$, let $n \geq 1$, and write $p = 2^n$. Let M be a finite flag-transitive map of type $\{p, q\}$ and girth p whose existence is guaranteed by Theorem 2.1.

Let B be the block obtained by gluing regular right-angled $2q$ -gons along the map M as in Section 2. Let d be the number of boundary components of B , which is equal to the number of faces in M .

Let G be a finite, isotropic, strict polygonal graph of valence d and girth $p = 2^n$ covering the bipartite graph Θ on two vertices as in Theorem 3.2, and let $\pi: G \rightarrow \Theta$ be a covering map. Let $\sigma: V(\Theta) \rightarrow \{-1, 1\}$ and $\chi: E(\Theta) \rightarrow \{1, \dots, d\}$ be bijections, where $V(\Theta)$ and $E(\Theta)$ are the sets of vertices and edges of Θ respectively. These induce proper colorings $\sigma \circ \pi$ and $\chi \circ \pi$ of the vertices and edges of G respectively.

For each $v \in V(G)$, let B_v be a copy of the block B , equipped with its standard orientation if $\sigma(v) = 1$ and with the reverse orientation if $\sigma(v) = -1$. Let b_1, \dots, b_d be the boundary components of B and label the boundary components of any copy B_v in the same way so that the isometric identification $B_v \cong B$ preserves the indices of boundary components.

Here is how we define the closed hyperbolic surface X given the above combinatorial data. For any edge $e = \{u, v\}$ in G , glue B_u to B_v by the identity map along their j -th boundary component, where $j = \chi \circ \pi(e)$. The surface X is defined as the quotient of $\sqcup_{v \in V(G)} B_v$ by these gluings. Since the gluing maps are orientation-reversing, X is an oriented surface. It has empty boundary since the coloring $\chi \circ \pi$ takes all values in $\{1, \dots, d\}$ on the edges containing a given vertex v , so that all the boundary components of B_v are glued. Lastly, X is compact because M and G are finite.

The main reason for using strict polygonal graphs in this construction is so that the blue arcs in the blocks B_v all close up to curves of the same length in X .

Lemma 3.3. *Any blue arc in a block $B_v \subset X$ is part of a closed geodesic of length pL in X .*

Proof. Any blue arc α_v in B_v connects two boundary geodesics b_i and b_j . The block B_v is glued to two other blocks B_u and B_w via these boundary components, and there are blue arcs $\alpha_u \subset B_u$ and $\alpha_w \subset B_w$ corresponding to α_v under the isometric identifications $B_u \cong B_v \cong B_w$. The concatenation $\alpha_u \cup \alpha_v \cup \alpha_w$ is geodesic since α_v is orthogonal to ∂B_v .

By our convention, the arc α_u (resp. α_w) connects the boundary components of B_u (resp. B_w) labelled b_i and b_j . By repeating the above reflection process with α_u or α_w instead of α_v (and so on), we obtain a bi-infinite path $\delta = (\dots, u, v, w, \dots)$ in the graph G whose edges alternate between the colors i and j . There is also a bi-infinite geodesic

$$\beta = \dots \cup \alpha_u \cup \alpha_v \cup \alpha_w \cup \dots$$

in X obtained by concatenating the corresponding blue arcs.

Since G is a strict polygonal graph, the path (u, v, w) is contained in a unique non-trivial cycle γ of length p (the girth of G). Furthermore, Theorem 3.2 stipulates that γ covers a closed cycle of length 2 in Θ under the covering map $\pi: G \rightarrow \Theta$. This cycle of length 2 is necessarily formed by the edges $\chi^{-1}(i)$ and $\chi^{-1}(j)$ since π respects the coloring of edges. This means that the edges of γ alternate between the colors i and j , and hence that the path δ wraps around γ periodically in both directions. In other words, δ closes up after p steps. Similarly, the geodesic β is closed and its length is equal to pL since each of its p subarcs has length L . \square

Note that we have not used the hypotheses that M is flag-transitive nor that G is isotropic yet. This will come up in Section 5 where we determine the isometry group of X .

4. Systoles

In this section, we determine and count the systoles in the surface X constructed above.

Proposition 4.1. *Let X be the surface constructed in Section 3. The systoles in X are the red curves and the blue curves. These systoles fill X and there are $\frac{4q}{(q-2)p}(g-1)$ of them, where q is the valence of the map M , p is the girth of M and the gluing graph G , and g is the genus of X .*

Proof. Let γ be a systole in X . If γ is contained in a single block $B_v \subset X$, then γ is a red curve (of length pL) by Lemma 2.3. Now assume that γ is not contained in any block. Then the blocks B_{v_1}, \dots, B_{v_k} ($k \geq 2$) that it visits define a closed cycle $s = (v_1, \dots, v_k, v_1)$ in the graph G . First suppose that s is trivial in G . Then s contains at least two backtracks, that is, vertices v_j in the sequence such that $v_{j-1} = v_{j+1}$. If s backtracks at a vertex $u \in G$, this means that a subarc ω of γ enters and leaves the block B_u via the same boundary component. By Corollary 2.4, ω has length strictly larger than $pL/2$. Since there are at least two disjoint subarcs like this, γ is longer than pL . We conclude that s is non-trivial in G , so that its length is at least p , the girth of G . But for each vertex u along s , the corresponding subarc of γ in B_v has length at least L by Lemma 2.5. Thus the total length of γ is at least pL . If equality occurs, then γ is a concatenation of blue arcs. Conversely, any concatenation of blue arcs has length pL by Lemma 3.3.

The complementary components of the set of systoles in X are precisely the interiors of the tiles from which the blocks are assembled. In particular, the systoles fill.

The number of systoles in X is equal to the total number of red arcs and blue arcs divided by p . This is because the red arcs are joined in groups of p to form systoles, and similarly for the blue arcs. Each such arc α (either red or blue) belongs to exactly

two tiles. The rhombus with one vertex in the center of each of these two tiles and diagonal α has area $\pi(q-2)/q$ by the Gauss–Bonnet formula (it has two right angles and two angles π/q). These rhombi tile X , which has area $4\pi(g-1)$. Therefore, the number of systoles is $4\pi(g-1)$ divided by $\pi(q-2)/q$, divided by p . \square

Recall that in the construction of X we could take any $q \geq 3$ and $p = 2^n$ for any $n \geq 1$. Given any $\varepsilon > 0$, taking n sufficiently large and any $q \geq 3$ gives a surface with a filling set of at most εg systoles. This proves Theorem 1.1. At the other extreme, the largest number of systoles is obtained when $q = 3$ and $p = 2$, which gives $6g - 6$ systoles. By [19, Theorem 2.8], such a surface has too few systoles to be a local maximum of the systole function, but we will see later that it is nevertheless a critical point of lower index.

Example 4.2. For any $g \geq 2$, if we take the map M to be the bipartite graph of valence $g + 1$ on two vertices (as a map on the sphere), then the resulting block B has $g + 1$ boundary components. Taking the gluing graph G to be equal to M , we obtain a surface X which is the double of B across its boundary. The genus of X is then equal to g . Since $q = g + 1$ and $p = 2$, the number of systoles is $2g + 2$ according to the formula in Proposition 4.1. Removing any two intersecting systoles leaves a filling set of $2g$ systoles. This example was previously described in [24, Theorem 36] and [2, Section 5] and was the starting point of this paper.

Remark 4.3. We could allow the graphs G and M to have different girths p and r by replacing the polygons P in the blocks to be semi-regular with side lengths L_{blue} and L_{red} satisfying $pL_{\text{blue}} = rL_{\text{red}}$. A version of Proposition 4.1 still holds for this generalization, with the count of systoles coming to

$$\frac{2q}{(q-2)} \left(\frac{1}{p} + \frac{1}{r} \right) (g-1).$$

All one has to do is change Lemma 2.2 to say that the distance between any two blue sides is at least L_{red} and the distance between any two red sides is at least L_{blue} , and modify the other lemmata accordingly.

5. Isometries

In this section, we determine the isometry group of the surface X up to index 2. Recall that the blocks $B_v \subset X$ (where $v \in V(G)$) are tiled by regular right-angled $2q$ -gons P_u (where $u \in V(M)$). By connecting the center of each polygon P_u to the midpoints of its edges with geodesics, we obtain a tiling \mathcal{Q} of X by $(2, 2, 2, q)$ -quadrilaterals (i.e., quadrilaterals with three right angles and one angle equal to π/q). Since any isometry of X preserves the set of systoles, it permutes the complementary polygons P_u and therefore the quadrilaterals in \mathcal{Q} . In fact, any quadrilateral can be sent to any other by an isometry.

Proposition 5.1. *The isometry group of X acts transitively on the quadrilaterals in the tiling \mathcal{Q} .*

Proof. The hypothesis that M is flag-transitive implies that the isometry group of B acts transitively on its $(2, 2, 2, q)$ -quadrilaterals. This is because there is a one-to-one correspondence between the flags in M and the quadrilaterals in B . The correspondence works as follows. Recall that M naturally embeds in B , connecting the centers of polygons P to their blue sides. A flag in M is the same as a half-edge e together with a choice of a face f containing e , either on the left or the right. In the tiling of B by quadrilaterals, there are exactly two quadrilaterals that have e as an edge. The side of e on which f lies determines which quadrilateral to pick. Since any map-automorphism of M can be realized as an isometry of B and M is flag-transitive, the claim follows.

Let $v \in V(G)$ and let $\phi: B_v \rightarrow B_v$ be an isometry. We claim that ϕ extends to an isometry Φ of X . First, the isometry ϕ induces a permutation τ on $\{1, \dots, d\}$ such that ϕ sends the boundary component b_i of B_v to the component $b_{\tau(i)}$ for every i . Now the edges adjacent to v in G are colored with the numbers $\{1, \dots, d\}$ according to the coloring $\chi \circ \pi$. Thus the permutation τ induces a bijection on the star of v . Since G is locally symmetric, this bijection can be extended to an automorphism ψ of G . If $x \in B_u \subset X$, then define $\Phi(x)$ to be the point $\phi(x)$ in $B_{\psi(u)}$, where we use the canonical identifications $B_w \cong B_v$ to transport the action of ϕ onto any block. This map is well-defined, for if $x \in B_u \cap B_v$ then x belongs to the boundary component labelled $i = \chi \circ \pi(\{u, v\})$ of B_u and B_v . By definition, $\phi(x)$ belongs to the $\tau(i)$ -th boundary component of B_v . Now $B_{\psi(u)}$ and $B_{\psi(v)}$ are glued along their boundary component labelled $\chi \circ \pi(\psi(\{u, v\}))$. This number equals $\tau(i)$ provided that the automorphism ψ is chosen to be a lift of the automorphism of Θ induced by τ , and this is possible according to [26, Theorem 3.2]. The map Φ is an isometry since it is a locally isometry as well as a bijection.

Similarly, any automorphism ψ of G which preserves the coloring $\chi \circ \pi$ defines an isometry Ψ of X by sending $x \in B_v$ to the corresponding x in $B_{\psi(v)}$. This simply shuffles the blocks around, acting by the identity map on the blocks. Note that the group of such automorphisms ψ acts transitively on the vertices of G .

Combining these two types of isometries gives the desired result. In order to send a quadrilateral $Q \subset B_u$ to another quadrilateral $Q' \subset B_v$, first apply an isometry Ψ as in the previous paragraph to send B_u to B_v . Then move $\Psi(Q)$ to Q' via an isometry Φ of the first type, preserving the block B_v . \square

Since there are at most two isometries of X sending one quadrilateral to another, this determines the isometry group of X up to index 2. We can reformulate this as follows. Subdivide \mathcal{Q} further into a tiling \mathcal{T} by $(2, 4, 2q)$ -triangles by bisecting the quadrilaterals at their smallest angle. Then the isometry group of X may or may

not act transitively on the tiles of \mathcal{T} depending on the graphs M and G used to construct X .

In Example 4.2, the isometry group of X acts transitively on these triangles, but that is not the case in general. That is, there can be an asymmetry between the red and blue curves in X . For example, let M be the flag-transitive map of type $\{4, 4\}$ obtained by subdividing the square torus into a 5×5 grid (so that B is a torus with 25 holes) and let G be the 1-skeleton of the 25-dimensional cube. Then each component of $X \setminus \{\text{blue curves}\}$ is a torus with 16 boundary components corresponding to a 4-dimensional subcube of G , while the complementary components of the red curves are the blocks with 25 boundary curves each. In this case, no isometry of X can interchange the two families of systoles.

6. Critical point and index

A real-valued function f on an n -dimensional manifold M is a *topological Morse function* if for every $p \in M$, there is an open neighborhood U of p and an injective continuous map $\phi: U \rightarrow \mathbb{R}^n$ with $\phi(p) = 0$ such that $f \circ \phi^{-1} - f(p)$ takes either the form

$$(x_1, \dots, x_n) \mapsto x_1$$

or

$$(x_1, \dots, x_n) \mapsto -\sum_{i=1}^j x_i^2 + \sum_{i=j+1}^n x_i^2$$

for some $j \in \{0, \dots, n\}$. In the first case, p is an *ordinary point* and in the second case p is a *critical point of index j* . Critical points of index 0 and n are local minima and maxima respectively.

Let $g \geq 2$ and let \mathcal{T}_g be the Teichmüller space of marked, connected, oriented, closed, hyperbolic surfaces of genus g . This space is a smooth manifold diffeomorphic to \mathbb{R}^{6g-6} . The systole $\text{sys}(Y)$ of a surface $Y \in \mathcal{T}_g$ is the length of any of its shortest closed geodesics. As mentioned in the introduction, Akrouf [1] proved that $\text{sys}: \mathcal{T}_g \rightarrow \mathbb{R}_+$ is a topological Morse function.

Let $Y \in \mathcal{T}_g$ and let \mathcal{S} be the set of (homotopy classes of) systoles in Y . For each $\alpha \in \mathcal{S}$ and $Z \in \mathcal{T}_g$, we let $\ell_\alpha(Z)$ be the length of the unique closed geodesic homotopic to α in Z . These functions are differentiable on \mathcal{T}_g and we denote their differentials by $d\ell_\alpha$.

Definition 6.1. The point $Y \in \mathcal{T}_g$ is *eutactic* if for every $v \in T_Y \mathcal{T}_g$, the following implication holds: if $d\ell_\alpha(v) \geq 0$ for every $\alpha \in \mathcal{S}$, then $d\ell_\alpha(v) = 0$ for every $\alpha \in \mathcal{S}$. The *rank* of a eutactic point Y is the dimension of the image of the linear map

$$(d\ell_\alpha)_{\alpha \in \mathcal{S}}: T_Y \mathcal{T}_g \rightarrow \mathbb{R}^{\mathcal{S}}.$$

With these definitions, we have the following characterization of the critical points of sys [1, Theorem 1].

Theorem 6.2 (Akrouit). *The critical points of index j of the systole function are the eutactic points of rank j .*

We can now show that the surface X constructed in Section 3 is a critical point of sys and give an upper bound for its index.

Proposition 6.3. *Let X be as in Section 3. Then X is a critical point of index at most $\frac{4q}{(q-2)p}(g-1)$ for the systole function.*

Proof. Let \mathcal{S} be the set of systoles of X (the red curves and the blue curves). Suppose that $v \in T_X \mathcal{T}_g$ is such that $d\ell_\alpha(v) \geq 0$ for every $\alpha \in \mathcal{S}$ and let

$$w = \sum_{f \in \text{Isom}(X)} f_* v.$$

Then

$$d\ell_\alpha(w) = \sum_{f \in \text{Isom}(X)} d\ell_\alpha(f_* v) = \sum_{f \in \text{Isom}(X)} d\ell_{f(\alpha)}(v) \geq d\ell_\alpha(v) \geq 0 \quad (6.1)$$

for every $\alpha \in \mathcal{S}$. On the other hand, w is the lift to X of a deformation of the quotient orbifold $Q = X/\text{Isom}(X)$. By Proposition 5.1, Q is either a $(2, 4, 2q)$ -triangle or a $(2, 2, 2, q)$ -quadrilateral. If Q is a triangle, then $w = 0$ so that $d\ell_\alpha(w)$ and $d\ell_\alpha(v)$ are both zero by Equation (6.1), for every $\alpha \in \mathcal{S}$. If Q is a quadrilateral, then its deformation space is 1-dimensional. This is because any $(2, 2, 2, q)$ -quadrilateral is determined by the lengths a and b of the two sides disjoint from the angle π/q , which satisfy the relation

$$\sinh a \sinh b = \cos(\pi/q)$$

(see [5, p. 454]). This equation implies that the lengths of the red curves and the blue curves in \mathcal{S} have opposite derivatives in the direction of w . Since the derivatives are non-negative, they must all be zero. We conclude that $d\ell_\alpha(v) = 0$ for every $\alpha \in \mathcal{S}$ from Equation (6.1). This shows that X is eutactic. The number of systoles in X is a trivial upper bound for the rank of X , and this number is equal to $\frac{4q}{(q-2)p}(g-1)$ by Proposition 4.1. \square

Once again, by taking p sufficiently large we obtain critical points of index at most εg for any $\varepsilon > 0$, thereby proving Theorem 1.3. This disproves the possibility envisaged by Schmutz Schaller [24, p. 410] that the minimal index were $2g - 1$.

7. Deformations preserving the systoles

Let \mathcal{X}_g be the subset of \mathcal{T}_g whose systoles fill. We would like to show that \mathcal{X}_g has relatively small codimension in \mathcal{T}_g . By Proposition 4.1, the systoles of any surface X constructed in Section 3 fill (recall that X depends on several parameters). Let \mathcal{S} be the set of systoles in X . If we deform X in such a way that the curves in \mathcal{S} remain of equal length, then these curves will still be the systoles for sufficiently small deformations. This is because the second shortest curve on X is longer by a definite amount and length varies continuously.

In other words, the intersection between the inverse image of the diagonal $\Delta \subset \mathbb{R}^{\mathcal{S}}$ by the map $(\ell_\alpha)_{\alpha \in \mathcal{S}}: \mathcal{T}_g \rightarrow \mathbb{R}^{\mathcal{S}}$ and a small neighborhood of X is contained in \mathcal{X}_g . One might be tempted to conclude directly that \mathcal{X}_g has codimension at most $|\mathcal{S}| - 1$ in \mathcal{T}_g . The subtlety is that the image of $(\ell_\alpha)_{\alpha \in \mathcal{S}}$ is not necessarily transverse to Δ . Indeed, the rank of X can be strictly less than $|\mathcal{S}| - 1$. For instance, the surface in Example 4.2 has rank $2g - 1$ according to [24, Theorem 36], while $|\mathcal{S}| = 2g + 2$.

To remedy this, one could try to get rid of redundant equations, i.e., to find a filling subset of curves $\mathcal{R} \subset \mathcal{S}$ for which the differential $(d_X \ell_\alpha)_{\alpha \in \mathcal{R}}$ is surjective and apply the implicit function theorem. The problem is that even if the curves in \mathcal{R} stay of equal length, the curves in $\mathcal{S} \setminus \mathcal{R}$ might become shorter and so the systoles might not fill anymore.

Another approach would be to find a nearby surface X_θ which has the same set of systoles as X , and hope that the differential $(d_{X_\theta} \ell_\alpha)_{\alpha \in \mathcal{S}}$ has full rank there. Below we will describe a 1-dimensional family of deformations of X with the same systoles. This fixes the issue of rank in some (but not all) cases. A similar idea was used in [18] to find a path of surfaces in \mathcal{X}_g with $2g$ systoles.

A 1-dimensional deformation. Recall that X is assembled from right-angled regular $2q$ -gons P whose sides are colored alternately red and blue, where $q \geq 3$. Given any $\theta \in (0, \pi)$, there exists a unique polygon P_θ with $2q$ equal sides and interior angles alternating between θ and $\pi - \theta$ (start with a triangle with angles π/q , $\theta/2$, and $(\pi - \theta)/2$ and reflect repeatedly across the two sides at angle π/q). To fix ideas, let us say that θ is the counter-clockwise (interior) angle from a red side to a blue side when going clockwise around P_θ and $\pi - \theta$ is the angle from blue to red. Now replace all the polygons P in X by P_θ while keeping the same gluing combinatorics. By construction, the total angle around vertices of the resulting tiling is 2π so the deformed surface X_θ is still a closed hyperbolic surface. Moreover, the red sides still line up to form closed geodesics and similarly for the blue sides. These closed geodesics all have equal length, namely, p times the side length of P_θ . As long as θ is close enough to $\pi/2$, these curves will remain the systoles.

The goal is then to show that the linear map $(d_{X_\theta} \ell_\alpha)_{\alpha \in \mathcal{S}}$ has full rank when $\theta \neq \pi/2$. We can do this for some small examples (see below), but we do not know how to handle surfaces with complicated gluing graphs of large girth. We present examples with full rank for girth 2 and 3 below.

Computing the rank. To prove that the derivative of lengths has full rank, it suffices to find a set of tangent vectors $\{v_\beta\}_{\beta \in \mathcal{S}}$ to Teichmüller space for which the square matrix $(d_{X_\theta} \ell_\alpha(v_\beta))_{\alpha, \beta \in \mathcal{S}}$ has non-zero determinant. For this, we can choose each vector v_β to be the Fenchel-Nielsen twist deformation (i.e., left earthquake) around the curve β . The cosine formula of Wolpert [29] and Kerckhoff [11] then says that

$$d\ell_\alpha(v_\beta) = \sum_{p \in \alpha \cap \beta} \cos \angle_p(\alpha, \beta)$$

whenever α and β are transverse, where $\angle_p(\alpha, \beta)$ is the counter-clockwise angle from α to β at the point p . In our case, two distinct curves $\alpha, \beta \in \mathcal{S}$ intersect at most once, with angle θ from red to blue or $\pi - \theta$ from blue to red. If we split the rows and columns of $D = (d_{X_\theta} \ell_\alpha(v_\beta))_{\alpha, \beta \in \mathcal{S}}$ by color we get a block matrix of the form

$$D = \cos \theta \begin{pmatrix} 0 & A \\ -A^\top & 0 \end{pmatrix}$$

where A is the matrix of zeros and ones recording which red curves intersect which blue curves. If $\theta \neq \pi/2$, then D has full rank if and only if the matrix

$$\tilde{D} = \begin{pmatrix} 0 & A \\ A^\top & 0 \end{pmatrix}$$

does. This matrix is the adjacency matrix of some graph $I_\mathcal{S}$, namely, the graph whose vertices are the systoles of X and where two vertices are joined by an edge if and only if the corresponding systoles intersect.

The determinant of the adjacency matrix of a graph counts something combinatorial on the graph. Indeed, according to [8] we have

$$\det(\tilde{D}) = \sum_{J \subset I_\mathcal{S}} (-1)^{\#\{\text{even components of } J\}} 2^{\#\{\text{cycles in } J\}}$$

where the sum is over all spanning subgraphs of $J \subset I_\mathcal{S}$ (subgraphs containing all vertices) which are *elementary*, meaning that their components are either edges or embedded cycles. The even components are those with an even number of vertices. In our case, $I_\mathcal{S}$ is bipartite so that all its cycles are even.

We are now ready to give some examples where \tilde{D} is invertible.

Examples of girth 2. The first family of examples comes from Example 4.2. In that example, the red and blue curves form a chain, that is, $I_\mathcal{S}$ is a cycle of length $2g + 2$. It follows that $I_\mathcal{S}$ has exactly three elementary spanning subgraphs: $I_\mathcal{S}$ itself, and two subgraphs obtained by deleting every other edge in $I_\mathcal{S}$. If $g = 2m$ is even, then $I_\mathcal{S}$ has $4m + 2$ edges and

$$\det(\tilde{D}) = (-1)^1 2^1 + (-1)^{2m+1} 2^0 + (-1)^{2m+1} 2^0 = -4 \neq 0. \tag{7.1}$$

Alternatively, one could compute the determinant of \tilde{D} by using the fact that A is a circulant matrix in this case.

The fact that \tilde{D} has non-zero determinant implies that the derivative of $\ell = (\ell_\alpha)_{\alpha \in \mathcal{S}}: \mathcal{T}_g \rightarrow \mathbb{R}^{\mathcal{S}}$ has full rank at X_θ whenever $\theta \neq \pi/2$. By the implicit function theorem, near X_θ we have that $\ell^{-1}(\Delta)$ is a smooth submanifold of codimension

$$|\mathcal{S}| - 1 = (2g + 2) - 1 = 2g + 1,$$

hence of dimension $4g - 7$. As explained earlier, $\ell^{-1}(\Delta)$ intersected with a sufficiently small ball around X_θ is contained in \mathcal{X}_g . We have thus proved that \mathcal{X}_g has dimension at least $4g - 7$ when g is even. We can push the proof a little further to obtain the following.

Theorem 7.1. *For every even $g \geq 2$, the set $\mathcal{X}_g \subset \mathcal{T}_g$ of closed hyperbolic surfaces of genus g whose systoles fill contains a cell of dimension $4g - 5$.*

Proof. Let X be the surface of genus g from Example 4.2 and let

$$\mathcal{S} = \{\alpha_1, \dots, \alpha_{2g+2}\}$$

be its set of systoles labelled in such a way that α_j intersects α_{j-1} and α_{j+1} for every j , where the indices are taken modulo $2g + 2$. Let X_θ be the deformation of X described above, where θ is close enough to $\pi/2$ so that its sets of systoles is still equal to \mathcal{S} . By Equation (7.1), the map $\ell = (\ell_\alpha)_{\alpha \in \mathcal{S}}: \mathcal{T}_g \rightarrow \mathbb{R}^{\mathcal{S}}$ is a submersion at the point X_θ . In particular, ℓ is open in a neighborhood of X_θ . This implies that there exist surfaces Y arbitrarily close to X_θ such that

$$\ell_{\alpha_1}(Y) = \dots = \ell_{\alpha_{2g}}(Y)$$

and such that these lengths are strictly less than $\ell_{\alpha_{2g+1}}(Y)$ and $\ell_{\alpha_{2g+2}}(Y)$. If Y is close enough to X_θ , then its set of systoles is a subset of \mathcal{S} by continuity of the length functions. Therefore, there is a sequence Y_n converging to X_θ such that the systoles in Y_n are given by the set $\mathcal{R} = \{\alpha_1, \dots, \alpha_{2g}\}$.

If n is large enough, then the square matrix $(d_{Y_n} \ell_\alpha(v_\beta))_{\alpha, \beta \in \mathcal{R}}$ will have non-zero determinant. Indeed, in the limit the matrix has the form

$$(d_{X_\theta} \ell_\alpha(v_\beta))_{\alpha, \beta \in \mathcal{R}} = \cos \theta \begin{pmatrix} 0 & B \\ -B^\top & 0 \end{pmatrix}$$

which is invertible because $\begin{pmatrix} 0 & B \\ B^\top & 0 \end{pmatrix}$ is the adjacency matrix of a tree $I_{\mathcal{R}}$ with an even number of vertices. Up to sign, its determinant is the number of perfect matchings (spanning subgraphs whose components are edges) in $I_{\mathcal{R}}$, which is equal to one. Furthermore, the entries of $(d_Y \ell_\alpha(v_\beta))_{\alpha, \beta \in \mathcal{R}}$ depend continuously on the surface Y in the same way that the angles of intersection between geodesics do.

Let $Y = Y_n$ for any such large enough n . Then the systoles in Y are given by the set \mathcal{R} and the map $(\ell_\alpha)_{\alpha \in \mathcal{R}}: \mathcal{T}_g \rightarrow \mathbb{R}^{\mathcal{R}}$ is a submersion at Y . By the implicit function theorem, the inverse image of the diagonal by this map is a submanifold of codimension $2g - 1$ near Y . Since the curves in \mathcal{R} fill, a small neighborhood of Y in this submanifold is contained in \mathcal{X}_g . The curves in \mathcal{R} fill because the complement of the curves in \mathcal{S} is a union of four polygons which meet at the intersection of α_{2g+1} and α_{2g+2} . Adding these two curves fuses the four polygons into a single one. \square

When g is odd, the matrix \tilde{D} is singular, but this does not necessarily imply that the image of $(\ell_\alpha)_{\alpha \in \mathcal{S}}$ is not transverse to the diagonal.

An example of girth 3. Next, we present an example of genus $g = 6$ where the underlying graphs M and G for the surface X have girth 3 and the matrix \tilde{D} is non-singular. Let M be the 1-skeleton of a regular tetrahedron (as a map of type $\{3, 3\}$ on the sphere) and let B be the corresponding block. This is a sphere with 4 holes and tetrahedral symmetry. Although this does not fit in the theory of Section 3, it is possible to glue five copies of B along the complete graph K_5 in such a way that the blue arcs connect up in groups of three to form closed geodesics. To see this, it is convenient to draw K_5 in \mathbb{R}^3 with a 3-fold symmetry as in Figure 1.




Figure 1. An embedding of K_5 in \mathbb{R}^3 and the corresponding gluing of tetrahedral blocks. (The red curves are printed in light grey and the blue curves in dark grey.)

The tetrahedral pieces are glued as suggested by the figure, in the simplest possible way (without twist). By inspection, the blue arcs connect in groups of three. The proof of Proposition 4.1 applies without change to show that the systoles in X are the red curves and the blue curves. The genus of X is equal to the number of edges in the complement of any spanning tree in K_5 , which is $10 - 4 = 6$. Let us label the red curves from 1 to 10 and the blue curves from a to j as in Figure 1 (the red curves correspond to the edges in K_5). Then the intersection matrix A is given by

$$A = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

which has determinant $48 \neq 0$. Therefore, the derivative of lengths $d\ell$ has full rank at X_θ whenever $\theta \neq \pi/2$. Note that this only gives us that \mathcal{X}_g has codimension at most 19 in \mathcal{T}_g , hence dimension at least $11 = 4g - 13$. The conclusion is weaker than that of Theorem 7.1, but we wanted to include this example to show that \tilde{D} can have full rank for more complicated graphs.

Questions. We conclude with a few questions related to the strategy we have just outlined.

Question 7.2. *Is there a sequence of graphs M and G as in Section 3 with girth going to infinity such that the corresponding intersection matrices \tilde{D} have non-zero determinants?*

In view of the above reasoning and the counting of Proposition 4.1, a positive answer would imply Conjecture 1.2. A major difficulty is that M and G are given to us in a non-explicit way from Theorem 2.1 and Theorem 3.2.

As the proof of Theorem 7.1 shows, one could bypass the determinant issue by finding a filling subset $\mathcal{R} \subset \mathcal{S}$ of even cardinality such that the corresponding intersection graph $I_{\mathcal{R}}$ is a tree, and a surface Y near X_θ whose systoles are exactly the curves in \mathcal{R} .

Question 7.3. *Given a surface X constructed as in Section 3 with set of systoles \mathcal{S} , is there an induced subtree in $I_{\mathcal{S}}$ with an even number of vertices such that the union of the corresponding curves fill?*

Question 7.4. *Let X be any hyperbolic surface, let \mathcal{S} be its set of systoles and let $\mathcal{R} \subset \mathcal{S}$ be a non-empty subset. Does there exist, in every neighborhood of X , a surface whose set of systoles is equal to \mathcal{R} ?*

Even if these questions have negative answers, they suggest how one should modify the construction of surfaces with sublinearly many systoles that fill in order to show that \mathcal{X}_g has large dimension: the systoles should cut the surface into a single polygon instead of several.

References

- [1] H. Akrouit, Singularités topologiques des systoles généralisées (French), *Topology*, **42** (2003), no. 2, 291–308. Zbl 1054.32006 MR 1941437
- [2] J. Anderson, H. Parlier, and A. Pettet, Small filling sets of curves on a surface, *Topology Appl.*, **158** (2011), no. 1, 84–92. Zbl 1238.30034 MR 2734699
- [3] D. Archdeacon and M. Perkel, Constructing polygonal graphs of large girth and degree. Proceedings of the Twentieth Southeastern Conference on Combinatorics, Graph Theory, and Computing (Boca Raton, FL, 1989), *Congr. Numer.*, **70** (1990), 81–85. Zbl 0699.05049 MR 1041586
- [4] M. Bridson and K. Vogtmann., Automorphism groups of free groups, surface groups and free abelian groups, in *Problems on mapping class groups and related topics*, 301–316, Proc. Sympos. Pure Math., 74, Amer. Math. Soc., Providence, RI, 2006. Zbl 1184.20034 MR 2264548
- [5] P. Buser, *Geometry and spectra of compact Riemann surfaces*. Reprint of the 1992 edition, Modern Birkhäuser Classics, Birkhäuser Boston, Ltd., Boston, MA, 2010. Zbl 1239.32001 MR 2742784
- [6] C. Evans, Net structure and cages, *Discrete Math.*, **27** (1979), no. 2, 193–204. Zbl 0407.05031 MR 537475
- [7] M. Fortier Bourque and K. Rafi, Local maxima of the systole function. arxiv:1807.08367
- [8] F. Harary, The determinant of the adjacency matrix of a graph, *SIAM Rev.*, **4** (1962), 202–210. Zbl 0113.17406 MR 144330
- [9] J. Harer, The virtual cohomological dimension of the mapping class group of an orientable surface, *Invent. Math.*, **84** (1986), no. 1, 157–176. Zbl 0592.57009 MR 830043
- [10] L. Ji, Well-rounded equivariant deformation retracts of Teichmüller spaces, *Enseign. Math.*, **60** (2014), no. 1-2, 109–129. Zbl 1303.32007 MR 3262437
- [11] S. Kerckhoff, The Nielsen realization problem, *Ann. of Math. (2)*, **117** (1983), no. 2, 235–265. Zbl 0528.57008 MR 690845
- [12] A. Lubotzky, Locally symmetric graphs of prescribed girth and Coxeter groups, *SIAM J. Discrete Math.*, **3** (1990), no. 2, 277–280. Zbl 0721.05031 MR 1039298
- [13] A. Mal'cev, On the faithful representation of infinite groups by matrices, *Mat. Sb.*, **8** (50) (1940), 405–422; English translation in *Amer. Math. Soc. Transl. (2)*, **45** (1965), 1–18.
- [14] J. Malestein, I. Rivin, and L. Theran, Topological designs, *Geom. Dedicata*, **168** (2014), 221–233. Zbl 1284.57020 MR 3158040

- [15] R. Nedela and M. Škoviera, Regular maps on surfaces with large planar width, *European J. Combin.*, **22** (2001), no. 2, 243–261. Zbl 0973.05025 MR 1808195
- [16] H. Parlier, Kissing numbers for surfaces, *J. Topol.*, **6** (2013), no. 3, 777–791. Zbl 1285.30023 MR 3100890
- [17] M. Perkel, On finite groups acting on polygonal graphs, Ph. D. thesis, University of Michigan, 1977. MR 2627108
- [18] B. Sanki, Systolic fillings of surfaces, *Bull. Aust. Math. Soc.*, **98** (2018), no. 3, 502–511. Zbl 1411.57005 MR 3877282
- [19] P. Schmutz, Riemann surfaces with shortest geodesic of maximal length, *Geom. Funct. Anal.*, **3** (1993), no. 6, 564–631. Zbl 0810.53034 MR 1250756
- [20] P. Schmutz, Systoles on Riemann surfaces, *Manuscripta Math.*, **85** (1994), no. 3-4, 429–447. Zbl 0819.30027 MR 1305753
- [21] P. Schmutz, Compact Riemann surfaces with many systoles, *Duke Math. J.*, **84** (1996), no. 1, 191–198. Zbl 0867.30029 MR 1394752
- [22] P. Schmutz Schaller, Extremal Riemann surfaces with a large number of systoles, in *Extremal Riemann surfaces (San Francisco, CA, 1995)*, 9–19, Contemp. Math., 201, Amer. Math. Soc., Providence, RI, 1997. Zbl 0867.30028 MR 1429190
- [23] P. Schmutz Schaller, Geometry of Riemann surfaces based on closed geodesics, *Bull. Amer. Math. Soc. (N.S.)*, **35** (1998), no. 3, 193–214. Zbl 1012.30029 MR 1609636
- [24] P. Schmutz Schaller, Systoles and topological Morse functions for Riemann surfaces, *J. Differential Geom.*, **52** (1999), no. 3, 407–452. Zbl 1033.32010 MR 1761080
- [25] A. Seress, Polygonal graphs, in *Horizons of combinatorics*, 179–188, Bolyai Soc. Math. Stud., 17, Springer, Berlin, 2008. Zbl 1170.05317 MR 2432533
- [26] A. Seress and E. Swartz, A note on the girth-doubling construction for polygonal graphs, *J. Graph Theory*, **68** (2011), no. 3, 246–254. Zbl 1229.05185 MR 2838335
- [27] J. Širáň, Triangle group representations and constructions of regular maps, *Proc. London Math. Soc. (3)*, **82** (2001), no. 3, 513–532. MR 1815966 Zbl 1015.05033
- [28] W. Thurston, A spine for Teichmüller space, *preprint*, 1985.
- [29] S. Wolpert, An elementary formula for the Fenchel–Nielsen twist, *Comment. Math. Helv.*, **56** (1981), no. 1, 132–135. Zbl 0467.30036 MR 615620

Received April 9, 2019

M. Fortier Bourque, School of Mathematics and Statistics, University of Glasgow,
University Place, Glasgow G12 8QQ, UK

E-mail: maxime.fortier-bourque@glasgow.ac.uk