

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 95 (2020)
Heft: 4

Artikel: Free rational points on smooth hypersurfaces
Autor: Browning, Tim / Sawin, Will
DOI: <https://doi.org/10.5169/seals-919560>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 03.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Free rational points on smooth hypersurfaces

Tim Browning and Will Sawin

Abstract. Motivated by a recent question of Peyre, we apply the Hardy–Littlewood circle method to count “sufficiently free” rational points of bounded height on arbitrary smooth projective hypersurfaces of low degree that are defined over the rationals.

Mathematics Subject Classification (2010). 11P55; 11D45, 14G05.

Keywords. Circle method, freeness, geometry of numbers, heights, rational points.

1. Introduction

Let $V \subset \mathbb{P}^{n-1}$ be a smooth hypersurface of degree $d \geq 3$, defined over the field of rational numbers. For $B \geq 1$, let

$$N_V(B) = \#\{x \in V(\mathbb{Q}) : H(x) \leq B\},$$

where H is the usual exponential height function on $\mathbb{P}^{n-1}(\mathbb{Q})$. Thanks to the Hardy–Littlewood circle method and work of Birch [2], it follows that there exists a constant $\delta > 0$ such that

$$N_V(B) = cB^{n-d} + O_V(B^{n-d-\delta}), \quad (1.1)$$

as $B \rightarrow \infty$, provided that $n > 2^d(d-1)$. Here $c = \frac{1}{n-d}\omega_H(V(\mathbf{A}_{\mathbb{Q}}))$ and ω_H is the Tamagawa measure on the space of adèles of V . The asymptotic formula (1.1) provided one of the earliest pieces of evidence for the conjecture of Manin [7], and its refinement by Peyre [10], about the distribution of rational points on Fano varieties.

The purpose of this paper is to address a very recent question of Peyre [11] about the distribution of “sufficiently free” rational points of bounded height on V . Peyre associates a measure of “freeness” $\ell(x) \in [0, 1]$ to any $x \in V(\mathbb{Q})$ and advocates the idea of only counting those rational points which satisfy $\ell(x) \geq \varepsilon_B$, where ε_B is a function of B decreasing to zero sufficiently slowly.¹ (See [11, Def. 6.11] for a precise statement for arbitrary Fano varieties over arbitrary number fields.) Peyre’s

¹A similar question was asked by Ellenberg and Venkatesh in a 2015 private communication with the first author.

function $\ell(x)$ is defined in (3.5) using Arakelov geometry and the theory of slopes associated to the tangent bundle \mathcal{T}_V . Let

$$N_V^{\varepsilon\text{-free}}(B) = \#\{x \in V(\mathbb{Q}) : \ell(x) \geq \varepsilon, H(x) \leq B\}. \quad (1.2)$$

In the setting of smooth hypersurfaces $V \subset \mathbb{P}^{n-1}$ of low degree, Peyre predicts that for a suitable range of ε , $N_V^{\varepsilon\text{-free}}(B)$ should have the same asymptotic behaviour as the usual counting function $N_V(B)$, as $B \rightarrow \infty$. The following result confirms this for a range of ε that is independent of B .

Theorem 1.1. *Let $d \geq 3$ and let $n > 3(d-1)2^{d-1}$. Then there exists a constant $c_{d,n} \in (0, 1)$ such that for any*

$$0 \leq \varepsilon < c_{d,n}$$

there exists a further constant $\delta > 0$ such that

$$N_V^{\varepsilon\text{-free}}(B) = cB^{n-d} + O_{V,\varepsilon}(B^{n-d-\delta}),$$

where $c = \frac{1}{n-d}\omega_H(V(\mathbb{A}_{\mathbb{Q}}))$ is the expected leading constant.

Note that in our theorem the parameter ε is a constant, while in Peyre's notion of freeness one takes ε_B tending to zero. Thus our result is stronger than necessary for Peyre's formulation. We shall show in §3 that it suffices to work with a simpler freeness function $\tilde{\ell}(x)$ that is defined in (3.4) in terms of the largest successive minimum of a certain associated lattice. Once this is achieved, the proof of Theorem 1.1 is guided by our investigation [4] of the analogous situation for smooth hypersurfaces over global fields of positive characteristic. We shall find that the role of the Riemann–Roch theorem in [4, §3] is replaced by the Poisson summation formula. After this the argument runs in close parallel to [4], apart from in one essential difference associated to primes of bad reduction for V .

An interesting feature of our method is that it relies on counting integer solutions (\mathbf{x}, \mathbf{y}) to the system of equations $f(\mathbf{x}) = \mathbf{y} \cdot \nabla f(\mathbf{x}) = 0$, where f is the defining polynomial of V . This is equivalent to counting integer points on the tangent bundle of the affine cone over V . This suggests that it may be possible to bound the number of rational points of small freeness on a Fano variety X by using asymptotics for the number of rational points on X together with asymptotics for the number of integral points on the tangent bundle of X .

Acknowledgements. The authors are very grateful to the anonymous referee for numerous pertinent remarks. While working on this paper the first author was supported by EPSRC grant EP/P026710/1. The research was partially conducted during the period the second author served as a Clay Research Fellow, and partially conducted during the period he was supported by Dr. Max Rössler, the Walter Haefner Foundation and the ETH Zurich Foundation.

2. The geometry of numbers and the shape of lattices

Most of the facts that we record in this section are taken from the book by Cassels [5]. Recall that a *lattice* Λ is a discrete additive subgroup of \mathbb{R}^n . Equivalently

$$\Lambda = \{x_1 \mathbf{b}_1 + \cdots + x_r \mathbf{b}_r : x_i \in \mathbb{Z}\},$$

for a set of linearly independent vectors $\mathbf{b}_1, \dots, \mathbf{b}_r \in \mathbb{R}^n$. The *rank* of Λ is then $\text{rank}(\Lambda) = r$ and the *determinant* is $\det(\Lambda) = \sqrt{\det(B^t B)}$, where B is the $n \times r$ matrix formed from the column vectors $\mathbf{b}_1, \dots, \mathbf{b}_r$. For each $1 \leq k \leq r$ let $s_k(\Lambda)$ be the least $\sigma > 0$ such that Λ contains at least k linearly independent vectors of Euclidean length bounded by σ . The $s_k(\Lambda)$ are the *successive minima* of Λ and they satisfy $0 < s_1(\Lambda) \leq s_2(\Lambda) \leq \cdots \leq s_r(\Lambda)$. Furthermore, it follows from Minkowski's second convex body theorem [5, §VIII.3.2] that

$$\det(\Lambda) \leq \prod_{i=1}^r s_i(\Lambda) \ll_n \det(\Lambda), \quad (2.1)$$

where the implied constant depends only on n . The *dual lattice* is defined to be

$$\Lambda^* = \{\mathbf{x} \in \text{span}_{\mathbb{R}}(\Lambda) : \mathbf{x} \cdot \mathbf{y} \in \mathbb{Z} \text{ for all } \mathbf{y} \in \Lambda\}.$$

This lattice has basis matrix $B(B^t B)^{-1}$ and so

$$\text{rank}(\Lambda^*) = r \quad \text{and} \quad \det(\Lambda^*) = \det(\Lambda)^{-1}.$$

Appealing to work of Banaszczyk [1, Thm. 2.1], it follows that

$$1 \leq s_k(\Lambda) s_{r-k+1}(\Lambda^*) \leq r, \quad (2.2)$$

for $1 \leq k \leq r$.

The following result is well-known and will prove instrumental in our work. A proof is given as a special case of work by Heath-Brown [8, Lemma 1].

Lemma 2.1. *For any vector $\mathbf{c} \in \mathbb{Z}_{\text{prim}}^n$ the set $\Lambda = \{\mathbf{x} \in \mathbb{Z}^n : \mathbf{c} \cdot \mathbf{x} = 0\}$ is a lattice of dimension $n - 1$ and determinant $\det(\Lambda) = \|\mathbf{c}\|$, where $\|\cdot\|$ is the Euclidean norm on \mathbb{R}^n .*

Given a lattice $\Lambda \subset \mathbb{R}^n$ of rank r it will be important to detect when the lattice is unusually skew, in the sense that the largest successive minimum is excessively large. To be precise, we seek a useful majorant for the indicator function

$$\mathbf{1}_R(\Lambda) = \begin{cases} 1 & \text{if } s_r(\Lambda) > R, \\ 0 & \text{otherwise.} \end{cases}$$

This is achieved in the following simple result.

Lemma 2.2. *Let $\Lambda \subset \mathbb{R}^n$ be a lattice of rank $r \leq n$ and let $\omega: \mathbb{R}^n \rightarrow \mathbb{R}$ be the Gaussian function $\omega(\mathbf{t}) = \exp(-\pi \|\mathbf{t}\|^2)$. Then*

$$\mathbf{1}_R(\Lambda) \leq \exp(\pi r^2) \frac{\det(\Lambda)}{R^r} \left(\sum_{\mathbf{y} \in \Lambda} \omega(\mathbf{y}/R) - \frac{R^r}{\det(\Lambda)} \right).$$

Proof. Note that $\omega(\mathbf{t}) \geq 0$ for all $\mathbf{t} \in \mathbb{R}^n$ and $\omega(\mathbf{0}) = 1$. It follows from the Poisson summation that

$$\sum_{\mathbf{y} \in \Lambda} \omega(\mathbf{y}/R) = \frac{R^r}{\det(\Lambda)} \sum_{\mathbf{y} \in \Lambda^*} \omega(R\mathbf{y}) = \frac{R^r}{\det(\Lambda)} \left(1 + \sum_{\substack{\mathbf{y} \in \Lambda^*, \\ \mathbf{y} \neq \mathbf{0}}} \omega(R\mathbf{y}) \right),$$

since $\hat{\omega} = \omega$. Thus

$$\sum_{\mathbf{y} \in \Lambda} \omega(\mathbf{y}/R) - \frac{R^r}{\det(\Lambda)} \geq 0$$

for any lattice Λ . Moreover, according to (2.2), we have $s_1(\Lambda^*) < r/R$ if $s_r(\Lambda) > R$. This means that there exists a non-zero vector $\mathbf{y}_0 \in \Lambda^*$ such that $\|\mathbf{y}_0\| < r/R$. But then

$$\sum_{\substack{\mathbf{y} \in \Lambda^* \\ \mathbf{y} \neq \mathbf{0}}} \omega(R\mathbf{y}) \geq \omega(R\mathbf{y}_0) = \exp(-\pi R^2 \|\mathbf{y}_0\|^2) \geq \exp(-\pi r^2).$$

This implies that

$$\sum_{\mathbf{y} \in \Lambda} \omega(\mathbf{y}/R) - \frac{R^r}{\det(\Lambda)} \geq \frac{\exp(-\pi r^2) R^r}{\det(\Lambda)}$$

if $s_r(\Lambda) > R$, which thereby completes the proof of the lemma. \square

3. Free rational points on hypersurfaces

Suppose that $f \in \mathbb{Z}[x_1, \dots, x_n]$ is a non-singular form of degree d that defines the hypersurface $V \subset \mathbb{P}^{n-1}$. Any rational point $x \in V(\mathbb{Q})$ has a representative vector $\mathbf{x} \in \mathbb{Z}_{\text{prim}}^n$ such that $f(\mathbf{x}) = 0$ and $x = (x_1 : \dots : x_n)$. The measure of freeness of x that we shall use in our paper is phrased in terms of the “well-shapedness” of the associated lattice

$$\Lambda_x = \{\mathbf{y} \in \mathbb{Z}^n : \mathbf{y} \cdot \nabla f(\mathbf{x}) = 0\}.$$

It follows from Lemma 2.1 that $\Lambda_x \subset \mathbb{Z}^n$ is a lattice of rank $n - 1$ and determinant

$$\det(\Lambda_x) = \frac{\|\nabla f(\mathbf{x})\|}{\gcd(\nabla f(\mathbf{x}))},$$

where $\|\cdot\|$ is the Euclidean norm. Let $\Delta_f \neq 0$ be the absolute value of the discriminant of the non-singular polynomial f . From the definition of the discriminant as the resultant of the forms $\partial f/\partial x_1, \dots, \partial f/\partial x_n$, it follows that there exists $e \in \mathbb{N}$ and algebraic identities

$$\Delta_f x_i^e = \sum_{1 \leq j \leq n} g_{i,j}(\mathbf{x}) \frac{\partial f}{\partial x_i}(\mathbf{x}), \quad (3.1)$$

for $1 \leq i \leq n$, where each $g_{i,j}$ has integer coefficients. In particular

$$\gcd(\nabla f(\mathbf{x})) | \Delta_f \quad \text{for all } \mathbf{x} \in \mathbb{Z}_{\text{prim}}^n. \quad (3.2)$$

Next we claim that

$$\|\mathbf{x}\|^{d-1} \ll \|\nabla f(\mathbf{x})\| \ll \|\mathbf{x}\|^{d-1} \quad \text{for all } \mathbf{x} \in \mathbb{R}^n, \quad (3.3)$$

for appropriate implied constants that depend only on f . Since f has degree d and so its partial derivatives have degree $d-1$, the upper bound is clear. To see the lower bound we note that $\nabla f(\mathbf{x}) \neq \mathbf{0}$ for all $\mathbf{x} \in \mathbb{R}^n$, since f is non-singular. Thus $\|\nabla f(\mathbf{x})\|$ is nowhere vanishing on the unit sphere and so attains some minimum value C , say, there. Thus we have $\|\nabla f(\mathbf{x})\| \geq C \|\mathbf{x}\|^{d-1}$ in general because f is homogeneous of degree d . This establishes (3.3).

As we shall see shortly, Peyre defines a freeness function relative to the smallest slope on the tangent bundle \mathcal{T}_V . The measure of freeness that we shall work with is related to this, but it is phrased in terms of the relative size of the largest successive minimum of the lattice Λ_x . To be precise, we set

$$\tilde{\ell}(x) = \frac{\log \|\mathbf{x}\| - \log s_{n-1}(\Lambda_x)}{\log \|\mathbf{x}\|}. \quad (3.4)$$

Then $\tilde{\ell}(x) \geq \varepsilon$ if and only if $s_{n-1}(\Lambda_x) \leq \|\mathbf{x}\|^{1-\varepsilon}$.

We gain some feeling for the behaviour of $\tilde{\ell}(x)$ by recalling (2.1). Thus for “typical” x one might expect the successive minima $s_k(\Lambda_x)$ to have the same order of magnitude, for $1 \leq k \leq n-1$. If this were true it would follow from (2.1) that

$$s_{n-1}(\Lambda_x) \asymp (s_1(\Lambda_x) \dots s_{n-1}(\Lambda_x))^{1/(n-1)} \asymp \det(\Lambda_x)^{1/(n-1)} \asymp \|\mathbf{x}\|^{1-\frac{n-d}{n-1}},$$

since $\|\nabla f(\mathbf{x})\| \asymp \|\mathbf{x}\|^{d-1}$ by (3.3). Such x satisfy $\tilde{\ell}(x) = (n-d)/(n-1) + o(1)$, as $H(x) \rightarrow \infty$. The following example shows a familiar situation in which the freeness function is unusually small.

Example 3.1. Consider the case $d = 3$ and $n = 4$ of a smooth cubic surface $V \subset \mathbb{P}^3$. Let $L \subset V$ be a \mathbb{Q} -line and define the associated rank 2 lattice

$$\mathbf{L} = \{\mathbf{0}\} \cup \{\mathbf{z} \in \mathbb{Z}^4 : (z_1 : \dots : z_4) \in L\} \subset \mathbb{Z}^4.$$

We claim that, for any $\varepsilon > 0$, we have $\tilde{\ell}(x) < \varepsilon$ for all but finitely many $x \in L$. To see this we note that $f(\mathbf{x} + t\mathbf{z})$ vanishes identically in t for all $z \in L$. But then it follows that $L \subset \Lambda_x$, in which case we have $1 \leq s_1(\Lambda_x) \leq s_2(\Lambda_x) \ll_L 1$. It now follows from (2.1) and (3.3) that

$$s_3(\Lambda_x) \gg_V \frac{\|\nabla f(\mathbf{x})\|}{s_1(\Lambda_x)s_2(\Lambda_x)} \gg_{L,V} \|\mathbf{x}\|^2.$$

This therefore yields $\tilde{\ell}(x) \leq -1 + o(1)$ and the claim.

We now explain how our freeness function (3.4) relates to that defined by Peyre [11, Déf. 4.11]. To begin with we can extend V to a closed subscheme $V \subset \mathbb{P}_{\mathbb{Z}}^{n-1}$. A rational point $x \in V(\mathbb{Q})$ gives a section $x \in V(\mathbb{Z})$ of this scheme. Because this scheme is smooth of dimension $n - 2$, the pullback $(\mathcal{T}_V)_x$ of its tangent bundle along x is a rank $n - 2$ free \mathbb{Z} -module; i.e. a free lattice of rank $n - 2$. Fixing a Riemannian metric on $V(\mathbb{R})$ gives a metric on this lattice. Peyre defines the freeness of x as

$$\ell(x) = \frac{\max\{(n-2)\mu_{n-2}((\mathcal{T}_V)_x), 0\}}{h(x)}, \quad (3.5)$$

where $h(x) = (n - d) \log \|\mathbf{x}\| + O(1)$ is the logarithmic anticanonical height of x and $\mu_1 \geq \dots \geq \mu_{n-2}$ are the slopes defined by Bost. There are four main differences between Peyre's definition and ours:

- (1) Peyre includes a factor of $n - 2$ in the numerator and the anticanonical height in the denominator instead of $\log \|\mathbf{x}\|$.
- (2) Peyre uses the notion of slopes instead of successive minima. The slopes of a lattice differ from minus the logarithms of its successive minima by $O(1)$.
- (3) Peyre works in a slightly different lattice, namely the tangent lattice instead of the perpendicular lattice to $\nabla f(\mathbf{x})$. These lattices are closely related, but not identical, and this discrepancy means that we only produce an inequality (instead of an identity) between the two notions of freeness.
- (4) Peyre defines the freeness to always be non-negative.

The relationship between the two notions of freeness is articulated in the following result.

Lemma 3.2. *For any $x \in V(\mathbb{Q})$ we have*

$$\ell(x) \geq \frac{n-2}{n-d} \tilde{\ell}(x) + O\left(\frac{1}{h(x)}\right).$$

Proof. We first explain how to relate the tangent lattice to Λ_x , and then why this leads to the stated inequality. We have an Euler exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^{n-1}} \rightarrow \mathcal{O}_{\mathbb{P}^{n-1}}(1)^n \rightarrow \mathcal{T}_{\mathbb{P}^{n-1}} \rightarrow 0$$

on $\mathbb{P}_{\mathbb{Z}}^{n-1}$. This induces an exact sequence

$$0 \rightarrow (\mathcal{O}_{\mathbb{P}^{n-1}})_x \rightarrow (\mathcal{O}_{\mathbb{P}^{n-1}}(1)^n)_x \rightarrow (\mathcal{T}_{\mathbb{P}^{n-1}})_x \rightarrow 0.$$

We have $(\mathcal{O}_{\mathbb{P}^{n-1}})_x = \mathbb{Z}$ and $(\mathcal{O}_{\mathbb{P}^{n-1}}(1)^n)_x = \mathbb{Z}^n$ because $\mathcal{O}_{\mathbb{P}^{n-1}}$ and $\mathcal{O}_{\mathbb{P}^{n-1}}(1)$ are rank one locally free sheaves, so their pullback along x are rank one locally free sheaves on $\text{Spec } \mathbb{Z}$, which are all isomorphic to \mathbb{Z} . The map between them is multiplication by \mathbf{x} , so the tangent lattice of \mathbb{P}^{n-1} is the quotient lattice $\mathbb{Z}^n/\mathbb{Z}\mathbf{x}$. We claim that the induced metric on this is the renormalized metric

$$\|\mathbf{y} + \mathbb{Z}\mathbf{x}\| = \frac{\min_{t \in \mathbb{R}} (\|\mathbf{y} - t\mathbf{x}\|)}{\|\mathbf{x}\|}.$$

Formally this arises from the $\mathcal{O}_{\mathbb{P}^{n-1}}(1)$ twist, but we can see this explicitly since the natural isomorphism between $\mathbb{R}^n/\mathbb{R}\mathbf{x}$ and the tangent space to $\mathbb{P}_{\mathbb{R}}^{n-1}$ at \mathbf{x} depends on the scaling of the vector \mathbf{x} and not just on its equivalence class in \mathbb{P}^{n-1} . The Arakelov metric on the tangent bundle of projective space must depend continuously on a point in projective space. To make it do so, we divide by $\|\mathbf{x}\|$.

Calculating $(\mathcal{T}_V)_x$ is now relatively easy. Consider the exact sequence

$$0 \rightarrow \mathcal{T}_V \rightarrow \mathcal{T}_{\mathbb{P}^{n-1}} \rightarrow \mathcal{O}_V(d) \rightarrow 0,$$

where the second map represents dot product with $\nabla f(\mathbf{x})$. We can realise $(\mathcal{T}_V)_x$ as the kernel of dotting with $\nabla f(\mathbf{x})$ in $\mathbb{Z}^n/\mathbb{Z}\mathbf{x}$, with no further renormalization necessary. Invoking some basic properties of slopes, we deduce that

$$\begin{aligned} \mu_{n-2}((\mathcal{T}_V)_x) &= \mu_{n-2}(\Lambda_x/\mathbb{Z}\mathbf{x}) + \log \|\mathbf{x}\| \\ &\geq \mu_{n-1}(\Lambda_x) + \log \|\mathbf{x}\| \\ &\geq -s_{n-1}(\Lambda_x) + \log \|\mathbf{x}\|. \end{aligned}$$

Indeed, the first step uses the fact that, when we divide the metric of a lattice by $\|\mathbf{x}\|$, we add $\log \|\mathbf{x}\|$ to each slope of the lattice, which is clear from the definition [11, Déf. 4.4] and is a special case of [3, Lemma 4.2]. The second step uses the fact that the minimum slope of a quotient lattice is at least the minimum slope of the original lattice, which is immediate from the definition of the minimum slope as a minimum over quotients of the lattice in [3, p. 195] and the equivalence of Bost's minimum slope and the last slope in Peyre's ordering. The last step uses [11, Remarque 4.7(b)].

The inequality

$$\frac{(n-2)\mu_{n-2}((\mathcal{T}_V)_x)}{h(x)} \geq \frac{n-2}{n-d} \tilde{\ell}(x) + O\left(\frac{1}{h(x)}\right)$$

immediately follows, since $h(x) = (n-d) \log \|\mathbf{x}\| + O(1)$. We therefore have

$$\begin{aligned} l(x) &= \frac{\max\{(n-2)\mu_{n-2}((\mathcal{T}_V)_x), 0\}}{h(x)} \geq \frac{(n-2)\mu_{n-2}((\mathcal{T}_V)_x)}{h(x)} \\ &\geq \frac{n-2}{n-d} \tilde{\ell}(x) + O\left(\frac{1}{h(x)}\right) \end{aligned}$$

except if $h(x) < 0$ where the middle inequality fails, but this happens for only finitely many x and we can handle it by assuming that the constant in the $O(1/h(x))$ term is sufficiently large. \square

Returning to (1.2), we can now make sense of the counting function

$$\begin{aligned} N_V^{\varepsilon\text{-free}}(B) &= \#\{x \in V(\mathbb{Q}) : \ell(x) \geq \varepsilon, H(x) \leq B\} \\ &= \#\{x \in V(\mathbb{Q}) : H(x) \leq B\} - E_{V,\varepsilon}(B), \end{aligned}$$

for any $\varepsilon > 0$, where

$$E_{V,\varepsilon}(B) = \#\{x \in V(\mathbb{Q}) : \ell(x) < \varepsilon, H(x) \leq B\}.$$

The first term is handled by (1.1), since $3(d-1)2^{d-1} \geq 2^d(d-1)$. Moreover, in view of Lemma 3.2 and the fact that $d \geq 3$, we have

$$\begin{aligned} E_{V,\varepsilon}(B) &\leq \#\left\{x \in V(\mathbb{Q}) : \tilde{\ell}(x) < \frac{n-d}{n-2}\varepsilon + o(1), H(x) \leq B\right\} + O(1) \\ &\leq \#\{x \in V(\mathbb{Q}) : \tilde{\ell}(x) < \varepsilon, H(x) \leq B\} + O(1), \end{aligned}$$

where the presence of the $O(1)$ term is needed to account for the low height points. Hence

$$E_{V,\varepsilon}(B) \leq \frac{1}{2} \#\{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^n : f(\mathbf{x}) = 0, \|\mathbf{x}\| \leq B, s_{n-1}(\Lambda_x) > \|\mathbf{x}\|^{1-\varepsilon}\} + O(1),$$

on taking into account the action of the units $\{\pm 1\}$ on $\mathbb{P}^{n-1}(\mathbb{Q})$. We require an upper bound for $E_{V,\varepsilon}(B)$ which is $O_{V,\varepsilon}(B^{n-d-\delta})$ for an appropriate $\delta > 0$, and which is valid for as wide a range of ε as possible.

To handle $E_{V,\varepsilon}(B)$ it will be convenient to break the range for $\|\mathbf{x}\|$ into dyadic intervals. Thus

$$\begin{aligned} E_{V,\varepsilon}(B) &\leq \sum_{\substack{R=2^j, \\ 1 \leq R \leq 2B}} \#\{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^n : f(\mathbf{x}) = 0, R/2 < \|\mathbf{x}\| \leq R, s_{n-1}(\Lambda_x) > R^{1-\varepsilon}\} \\ &\quad + O(1) \\ &= \sum_{\substack{R=2^j, \\ 1 \leq R \leq 2B}} E_{V,\varepsilon}^*(R) + O(1), \end{aligned} \tag{3.6}$$

say. Appealing to Lemma 2.2, we deduce that

$$E_{V,\varepsilon}^*(R) \ll 1 + \sum_{\substack{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^n, f(\mathbf{x})=0, \\ R/2 < \|\mathbf{x}\| \leq R}} \frac{\det(\Lambda_x)}{R^{(1-\varepsilon)(n-1)}} \left(\sum_{\mathbf{y} \in \Lambda_x} \omega(\mathbf{y}/R^{1-\varepsilon}) - \frac{R^{(1-\varepsilon)(n-1)}}{\det(\Lambda_x)} \right), \tag{3.7}$$

where $\omega(\mathbf{t}) = \exp(-\pi \|\mathbf{t}\|^2)$. In what follows it will be convenient to write $d(\mathbf{x}) = \det(\Lambda_x)$ when x is represented by a vector $\mathbf{x} \in \mathbb{Z}_{\text{prim}}^n$. In view of (3.2) we have

$$\gcd(\nabla f(\mathbf{x})) = \gcd(\nabla f(\mathbf{x}), \Delta_f),$$

so that

$$d(\mathbf{x}) = \frac{\|\nabla f(\mathbf{x})\|}{\gcd(\nabla f(\mathbf{x}), \Delta_f)}. \quad (3.8)$$

We shall use this formula to extend the definition of $d(\mathbf{x})$ to all (not necessarily primitive) vectors $\mathbf{x} \in \mathbb{Z}^n$.

Let us write $\|\mathbf{x}\| \sim R$ to denote the inequalities $R/2 < \|\mathbf{x}\| \leq R$. In order to treat $E_{V,\varepsilon}^*(R)$ we begin by analysing the term

$$M_\varepsilon(R) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim R}} d(\mathbf{x}) \sum_{\mathbf{y} \in \Lambda_x} \omega(\mathbf{y}/R^{1-\varepsilon}).$$

It is clear that $\mathbf{y} \cdot \nabla f(k\mathbf{x}) = 0$ if and only if $\mathbf{y} \cdot \nabla f(\mathbf{x}) = 0$, for any $k \in \mathbb{N}$. Hence, an application of Möbius inversion yields

$$\begin{aligned} M_\varepsilon(R) &= \sum_{k \leq R} \mu(k) \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim R/k}} d(k\mathbf{x}) \sum_{\substack{\mathbf{y} \in \mathbb{Z}^n, \\ \mathbf{y} \cdot \nabla f(\mathbf{x})=0}} \omega(\mathbf{y}/R^{1-\varepsilon}) \\ &= \sum_{k \leq R} \mu(k) \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim R/k}} d(k\mathbf{x}) \int_0^1 S(\beta) d\beta, \end{aligned} \quad (3.9)$$

where

$$S(\beta) = \sum_{\mathbf{y} \in \mathbb{Z}^n} \omega(\mathbf{y}/R^{1-\varepsilon}) e(\beta \mathbf{y} \cdot \nabla f(\mathbf{x})).$$

Our plan is to define a set of “major arcs” for the interval $[0, 1]$ whose integral matches the expected main term $R^{(1-\varepsilon)(n-1)} / \det(\Lambda_x)$ from (3.7).

4. Identification of the major arcs

Our identification of the major arcs follows the path that was paved in [4, §4]. Henceforth all implied constants will be allowed to depend on f . It will be convenient to set

$$X = X_k = R/k \quad \text{and} \quad Y = R^{1-\varepsilon},$$

where k is the parameter occurring in (3.9). Since $\hat{\omega} = \omega$, it follows from Poisson summation that

$$\begin{aligned} S(\beta) &= \sum_{\mathbf{y} \in \mathbb{Z}^n} \omega(\mathbf{y}/Y) e(\beta \mathbf{y} \cdot \nabla f(\mathbf{x})) \\ &= Y^n \sum_{\mathbf{y} \in \mathbb{Z}^n} \omega(Y\beta \nabla f(\mathbf{x}) - Y\mathbf{y}), \end{aligned}$$

for any $\beta \in [0, 1]$ and any $\mathbf{x} \in \mathbb{Z}^n$. Let us use $\langle \alpha \rangle = \inf_{m \in \mathbb{Z}} |m - \alpha|$ to denote the distance to the nearest integer. We observe that

$$\omega(\mathbf{t}) \ll_N (1 + \|\mathbf{t}\|)^{-N},$$

for any $\mathbf{t} \in \mathbb{R}^n$ and any $N \geq 0$. Hence it is not hard to see that

$$S(\beta) = Y^n \omega\left(Y\left\langle \beta \frac{\partial f(\mathbf{x})}{\partial x_1} \right\rangle, \dots, Y\left\langle \beta \frac{\partial f(\mathbf{x})}{\partial x_n} \right\rangle\right) + O_N(Y^{-N}), \quad (4.1)$$

for any $N \geq 0$. Led by this we make the following definition.

Definition 4.1 (Major arcs). For any $\eta > 0$ we set

$$\mathfrak{M}_\eta(X, Y) = \bigcup_{q \leq Y^{1-\eta}} \bigcup_{\substack{0 \leq a < q, \\ \gcd(a, q) = 1}} \left\{ \beta \in [0, 1) : |q\beta - a| \leq \frac{1}{C_f X^{d-1}} \right\},$$

where $C_f > 0$ is a sufficiently large constant that only depends on f .

The following result is concerned with the size of the exponential sum $S(\beta)$ when β belongs to this set of major arcs.

Lemma 4.2. *Let $N \geq 0$, let $\mathbf{x} \in \mathbb{Z}^n$ with $\|\mathbf{x}\| \leq X$, and let $\beta = a/q + \theta \in \mathfrak{M}_\eta(X, Y)$ for coprime integers a, q such that $0 \leq a < q$ and $|q\theta| \leq 1/(C_f X^{d-1})$. Then*

$$S(\beta) = Y^n \omega(Y\theta \nabla f(\mathbf{x})) + O_N(Y^{-N})$$

if $q \mid \nabla f(\mathbf{x})$ and $|\theta| \leq Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|$, with $S(\beta) = O_N(Y^{-N})$ otherwise.

Proof. Let $\beta = a/q + \theta \in \mathfrak{M}_\eta(X, Y)$. Then

$$\left| \theta \frac{\partial f(\mathbf{x})}{\partial x_i} \right| \leq \frac{1}{4q}$$

for any $1 \leq i \leq n$, provided that C_f is large enough. Next, we see that

$$\left\langle \beta \frac{\partial f(\mathbf{x})}{\partial x_i} \right\rangle = \left| \theta \frac{\partial f(\mathbf{x})}{\partial x_i} \right|$$

if $q \mid \partial f(\mathbf{x})/\partial x_i$. Thus it follows from (4.1) that

$$S(\beta) = O_N(Y^{-N})$$

if $q \mid \nabla f(\mathbf{x})$ and $|\theta| > Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|$. Alternatively, if we have $q \mid \nabla f(\mathbf{x})$ and $|\theta| \leq Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|$ then clearly

$$Y^n \omega \left(Y \left\langle \beta \frac{\partial f(\mathbf{x})}{\partial x_1} \right\rangle, \dots, Y \left\langle \beta \frac{\partial f(\mathbf{x})}{\partial x_n} \right\rangle \right) = Y^n \omega(Y\theta \nabla f(\mathbf{x})).$$

Finally, if $q \nmid \partial f(\mathbf{x})/\partial x_i$ for some $i \in \{1, \dots, n\}$ then there exists a non-zero integer $u \in [-q/2, q/2]$ such that $a\partial f(\mathbf{x})/\partial x_i \equiv u \pmod{q}$, whence

$$\left\langle \beta \frac{\partial f(\mathbf{x})}{\partial x_i} \right\rangle \geq \frac{3}{4q} \geq \frac{3}{4Y^{1-\eta}},$$

for $\beta \in \mathfrak{M}_\eta(X, Y)$. This shows that $S(\beta) = O_N(Y^{-N})$ in this case, as required to complete the proof of the lemma. \square

The following result is concerned with the evaluation of the integral of $S(\beta)$ over the major arcs.

Lemma 4.3. *Let $N \geq 0$ and assume that $\|\mathbf{x}\| \sim X$. Then*

$$\int_{\mathfrak{M}_\eta(X, Y)} S(\beta) d\beta = \frac{Y^{n-1} \gcd(\nabla f(\mathbf{x}))}{\|\nabla f(\mathbf{x})\|} (1 + O(\mathbf{1}(\mathbf{x}))) + O_N(Y^{-N}),$$

where

$$\mathbf{1}(\mathbf{x}) = \begin{cases} 1 & \text{if } \gcd(\nabla f(\mathbf{x}))C_f^2 > Y^{1-\eta}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Let us set $h = \gcd(\nabla f(\mathbf{x}))$ throughout the proof. We define the modified major arcs $\tilde{\mathfrak{M}}_\eta(X, Y)$ to be the set of $\beta = a/q + \theta \in \mathfrak{M}_\eta(X, Y)$ for which $q \mid h$ and $|\theta| \leq Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|$. We claim that these modified major arcs are non-overlapping. To see this we suppose that $a_1/q_1 + \theta_1 = a_2/q_2 + \theta_2$. Then we may assume without loss of generality that $q_1 = q_2 = h$. But then it follows that

$$|a_1 - a_2| = h|\theta_2 - \theta_1| \ll X^{d-1} \cdot \frac{1}{C_f X^{d-1}}.$$

Assuming that C_f is sufficiently large, this implies that $a_1 = a_2$, which thereby establishes the claim. (In fact it is not hard to check that the major arcs $\mathfrak{M}_\eta(X, Y)$ are also disjoint provided that $X^{d-1} \gg Y^{1-\eta}$.)

An application of Lemma 4.2 yields

$$\begin{aligned} \int_{\mathfrak{M}_\eta(X,Y)} S(\beta) d\beta &= \int_{\widetilde{\mathfrak{M}}_\eta(X,Y)} S(\beta) d\beta + O_N(Y^{-N}) \\ &= Y^n \sum_{\substack{q \leq Y^{1-\eta}, \\ q|h}} \varphi(q) \int_{|\theta| \leq \Theta} \omega(Y\theta \nabla f(\mathbf{x})) d\theta + O_N(Y^{-N}), \end{aligned}$$

for any $N \geq 0$, where

$$\Theta = \min \left\{ \frac{Y^{-1+\eta}}{\|\nabla f(\mathbf{x})\|}, \frac{1}{qC_f X^{d-1}} \right\}.$$

Since $(1 * \varphi)(h) = h$, it is clear that

$$\sum_{\substack{q \leq Y^{1-\eta}, \\ q|h}} \varphi(q) = h + O(h\mathbf{1}(\mathbf{x})),$$

where $\mathbf{1}(\mathbf{x})$ is as in the statement of the lemma.

Next, we observe that

$$\int_{|\theta| \leq Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|} \omega(Y\theta \nabla f(\mathbf{x})) d\theta = \frac{1}{Y\|\nabla f(\mathbf{x})\|} + O_N(Y^{-N}).$$

Moreover,

$$\int_{\Theta < |\theta| \leq Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|} \omega(Y\theta \nabla f(\mathbf{x})) d\theta$$

vanishes unless $\Theta < \|\nabla f(\mathbf{x})\|$, which implies that

$$\frac{1}{qC_f X^{d-1}} < \frac{Y^{-1+\eta}}{\|\nabla f(\mathbf{x})\|}.$$

Appealing to (3.3) and using the fact that $\|\mathbf{x}\| \sim X$, the right hand side is at most $C_f Y^{-1+\eta}/X^{d-1}$, if the constant C_f is taken to be sufficiently large in Definition 4.1. Hence we conclude that $qC_f^2 > Y^{1-\eta}$, which in turn implies that $hC_f^2 > Y^{1-\eta}$ and thus that $\mathbf{1}(\mathbf{x}) = 1$. Because the integrand is nonnegative, the integral over this restricted interval is at most $\frac{1}{Y\|\nabla f(\mathbf{x})\|}$, so that

$$\begin{aligned} \int_{|\theta| \leq \Theta} \omega(Y\theta \nabla f(\mathbf{x})) d\theta &= \int_{|\theta| \leq Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|} \omega(Y\theta \nabla f(\mathbf{x})) d\theta \\ &\quad - \int_{\Theta < |\theta| \leq Y^{-1+\eta}/\|\nabla f(\mathbf{x})\|} \omega(Y\theta \nabla f(\mathbf{x})) d\theta \\ &= \frac{1}{Y\|\nabla f(\mathbf{x})\|} + O\left(\frac{\mathbf{1}(\mathbf{x})}{Y\|\nabla f(\mathbf{x})\|}\right) + O_N(Y^{-N}). \end{aligned}$$

Putting everything together yields the statement of the lemma. \square

It is now time to return to our expression (3.9) for $M_\varepsilon(R)$. First, sticking with the notation $X = R/k$ and $Y = R^{1-\varepsilon}$, we deduce from Lemma 4.3 that

$$\sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim X}} d(k\mathbf{x}) \int_{\mathfrak{M}_\eta(X,Y)} S(\beta) d\beta = Y^{n-1} \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim X}} d(k\mathbf{x}) \cdot \frac{\gcd(\nabla f(\mathbf{x}))}{\|\nabla f(\mathbf{x})\|} \{1 + E(\mathbf{x})\},$$

where

$$E(\mathbf{x}) = O(\mathbf{1}(\mathbf{x})) + O_N(X^{d-1}Y^{-N})$$

for any $N \geq 0$. If $\mathbf{1}(\mathbf{x}) = 1$ then $\gcd(\nabla f(\mathbf{x})) > C_f^{-2}Y^{1-\eta}$. Furthermore, if $\gcd(x_1, \dots, x_n) = \ell$ then (3.1) yields

$$\gcd(\nabla f(\mathbf{x})) = \gcd(\ell^{d-1}\nabla f(\mathbf{x}/\ell)) = \ell^{d-1} \gcd(\nabla f(\mathbf{x}/\ell)) \ll \ell^{d-1},$$

whence in fact $\ell \gg Y^{(1-\eta)/(d-1)}$. Moreover, we have

$$d(k\mathbf{x}) \cdot \frac{\gcd(\nabla f(\mathbf{x}))}{\|\nabla f(\mathbf{x})\|} \leq k^{d-1} \gcd(\nabla f(\mathbf{x})) \ll k^{d-1} \ell^{d-1}.$$

Assume now that $n > 2^d(d-1)$. Then it follows from (1.1) that

$$\#\{\mathbf{x} \in \mathbb{Z}^n : f(\mathbf{x}) = 0, \|\mathbf{x}\| \leq X, \gcd(x_1, \dots, x_n) = \ell\} \ll N_V(X/\ell) \ll \frac{X^{n-d}}{\ell^{n-d}}.$$

Thus

$$\sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim X}} d(k\mathbf{x}) \cdot \frac{\gcd(\nabla f(\mathbf{x}))}{\|\nabla f(\mathbf{x})\|} E(\mathbf{x}) \ll_N k^{d-1} (X^{n-d}Y^{-\delta} + X^{n-1}Y^{-N})$$

for any $N \geq 0$, where

$$\delta = \frac{(1-\eta)(n-2d)}{d-1}.$$

Here the exponent δ arises from summing the $\ell^{-(n-d)+(d-1)} = \ell^{-(n-2d+1)}$ savings over $\ell \gg Y^{(1-\eta)/(d-1)}$. Reintroducing the sum over k , we now see that the overall contribution to $M_\varepsilon(R)$ from the set of major arcs $\mathfrak{M}_{\eta,k} = \mathfrak{M}_\eta(R/k, R^{1-\varepsilon})$ is

$$\begin{aligned} \sum_{k \leq R} \mu(k) \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim R/k}} d(k\mathbf{x}) \int_{\mathfrak{M}_{\eta,k}} S(\beta) d\beta \\ = R^{(1-\varepsilon)(n-1)} \sum_{\substack{\mathbf{x} \in \mathbb{Z}_{\text{prim}}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim R}} 1 + O(R^{n-d+(1-\varepsilon)(n-1-\delta)}), \end{aligned}$$

on taking N sufficiently large.

Putting $\mathfrak{m}_{\eta,k} = [0, 1) \setminus \mathfrak{M}_{\eta,k}$ and bringing everything together in (3.7), it now follows that

$$E_{V,\varepsilon}^*(R) \ll R^{n-d-(1-\varepsilon)\delta} + \left| \sum_{k \leq R} \mu(k) \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ f(\mathbf{x})=0, \|\mathbf{x}\| \sim R/k}} \frac{d(k\mathbf{x})}{R^{(1-\varepsilon)(n-1)}} \int_{\mathfrak{m}_{\eta,k}} S(\beta) d\beta \right|.$$

We may detect the equation $f(\mathbf{x}) = 0$ in the way most familiar to practitioners of the Hardy–Littlewood circle method. On doing so, we are led to the following result, which summarises our discussion of the major arcs.

Lemma 4.4. *Let $n > 2^d(d-1)$. For any $k \in \mathbb{N}$ let $\mathfrak{m}_{\eta,k} = [0, 1) \setminus \mathfrak{M}_{\eta,k}$, where $\mathfrak{M}_{\eta,k} = \mathfrak{M}_{\eta}(R/k, R^{1-\varepsilon})$ is given by Definition 4.1. Then there exists $\delta > 0$ such that*

$$E_{V,\varepsilon}^*(R) \ll R^{n-d-\delta} + \frac{1}{R^{(1-\varepsilon)(n-1)}} \sum_{k \leq \sqrt{R}} \mu^2(k) \int_0^1 \int_{\mathfrak{m}_{\eta,k}} |S(\alpha, \beta)| d\alpha d\beta,$$

where if $d(\mathbf{x})$ is given by (3.8) then

$$S(\alpha, \beta) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n \\ \|\mathbf{x}\| \sim R/k}} \sum_{\mathbf{y} \in \mathbb{Z}^n} d(k\mathbf{x}) \omega(\mathbf{y}/R^{1-\varepsilon}) e(\alpha f(\mathbf{x}) + \beta \mathbf{y} \cdot \nabla f(\mathbf{x})).$$

Proof. The only thing that requires comment is the truncation from $k \leq R$ to $k \leq \sqrt{R}$. But since $d(k\mathbf{x}) \ll R^{d-1}$ the trivial bound yields

$$\frac{|S(\alpha, \beta)|}{R^{(1-\varepsilon)(n-1)}} \ll \frac{R^{d-1} \cdot (R/k)^n \cdot R^{(1-\varepsilon)n}}{R^{(1-\varepsilon)(n-1)}} \ll \frac{R^{n+d-\varepsilon}}{k^n}.$$

Hence the tail of the k -summation makes a satisfactory contribution. \square

5. Treatment of the minor arcs

We begin with a technical result from the geometry of numbers, which generalises the “shrinking lemma” that is due to Davenport [6, Lemma 12.6], and which one recovers by taking $P = Q$ in the following result.

Lemma 5.1. *Let γ be a symmetric $n \times n$ matrix with entries in \mathbb{R} . Let $P > 0$, let $Q > 2$ and let $\theta \in (0, 1]$. Let $N_{\gamma,P,Q}$ be the number of $\mathbf{x} \in \mathbb{Z}^n$ such that $\|\mathbf{x}\| < P$ and $\max_{1 \leq i \leq n} \langle \gamma x_i \rangle < Q^{-1}$. Then*

$$\frac{N_{\gamma,P,Q}}{N_{\gamma,\theta P,\theta^{-1}Q}} \ll \theta^{-n} \max \left\{ \sqrt{\frac{P}{Q}}, 1 \right\}^n,$$

where the implied constant depends only on n .

Proof. We may assume that $P \geq 1$, since the left hand side is 1 when $P < 1$. Define the matrix

$$\Lambda_{P,Q} = \begin{pmatrix} P^{-1}I_n & 0 \\ Q\gamma & QI_n \end{pmatrix},$$

so that

$$\Lambda_{P,Q}^{-t} = \begin{pmatrix} PI_n & -P\gamma \\ 0 & Q^{-1}I_n \end{pmatrix} \quad \text{and} \quad \frac{Q}{P}\Lambda_{P,Q}^{-t} = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix} \Lambda_{P,Q} \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}^{-1}.$$

Let $R_1 \leq \dots \leq R_{2n}$ denote the successive minima of the lattice corresponding to $\Lambda_{P,Q}$ and let $R_1^* \leq \dots \leq R_{2n}^*$ be the successive minima of the dual lattice corresponding to $(Q/P)\Lambda_{P,Q}^{-t}$. Then (2.2) implies that $R_i^* \asymp (Q/P)/R_{2n-i+1}$, for $1 \leq i \leq 2n$. Since the lattices are equal up to left and right multiplication by a matrix in $\text{GL}_{2n}(\mathbb{Z})$, we must have

$$R_i \asymp \frac{Q/P}{R_{2n+1-i}}$$

for all $1 \leq i \leq 2n$. Taking $i = n+1$ we deduce that $\sqrt{Q/P} \ll R_{n+1}$.

Since $Q > 2$, the quantity $N_{\gamma,P,Q}$ is equal to the number of vectors in the lattice corresponding to $\Lambda_{P,Q}$ whose first n entries form a vector of Euclidean norm < 1 and whose last n entries are individually < 1 . Thus it is bounded below by the number of vectors with Euclidean norm < 1 , and bounded above by the corresponding number with Euclidean norm $< \sqrt{n+1}$. On the other hand, $N_{\gamma,\theta P,\theta^{-1}Q}$ is bounded below by the number of vectors in the lattice corresponding to $\Lambda_{P,Q}$ with norm $< \theta$ and above by the corresponding number with norm $< \theta\sqrt{2n+1}$. It therefore follows from Davenport [6, Lemma 12.4] that

$$N_{\gamma,P,Q} \asymp \prod_{i=1}^{2n} \max\{1, R_i^{-1}\} \quad \text{and} \quad N_{\gamma,\theta P,\theta^{-1}Q} \asymp \prod_{i=1}^{2n} \max\{1, \theta R_i^{-1}\},$$

where the implied constants depend only on n . Dividing term by term, we see that each i contributes at most θ^{-1} and each $i \geq n+1$ contributes at most $\max\{\sqrt{P/Q}, 1\}$. Thus the total contribution is at most $\theta^{-n} \max\{\sqrt{P/Q}, 1\}^n$, as claimed in the statement of the lemma. \square

The second technical result required is a simple Diophantine approximation result due to Heath-Brown [9, Lemma 2.3].

Lemma 5.2. *Let $M, R > 0$. Let $m \in \mathbb{Z}$ such that $|m| \leq M$ and let $\alpha = a/q + z$, with coprime integers a, q and $z \in \mathbb{R}$, such that $\langle \alpha m \rangle < R^{-1}$. Assume that*

$$|z| \leq (2qM)^{-1}, \quad q \leq R/2 \quad \text{and} \quad q > \min\{M, (|z|R)^{-1}\}.$$

Then $m = 0$.

The statement of this result requires the assumption that a and q are coprime, which isn't formally stated in [9, Lemma 2.3] but is implicit in the proof.

We now have the tools in place to study our exponential sum on the minor arcs. Let us set

$$X = R/k \quad \text{and} \quad Y = R^{1-\varepsilon}, \quad (5.1)$$

as previously. We want to study

$$S(\alpha, \beta) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n \\ \|\mathbf{x}\| \sim X}} \sum_{\mathbf{y} \in \mathbb{Z}^n} d(k\mathbf{x}) \omega(\mathbf{y}/Y) e(g(\mathbf{x}, \mathbf{y})),$$

for $(\alpha, \beta) \in [0, 1) \times \mathfrak{m}_{\eta, k}$, where $d(\mathbf{x})$ is given by (3.8) and

$$g(\mathbf{x}, \mathbf{y}) = \alpha f(\mathbf{x}) + \beta \mathbf{y} \cdot \nabla f(\mathbf{x}).$$

Let us write $\tilde{\Delta}_f = \Delta_f / \gcd(k^{d-1}, \Delta_f)$. Then we have

$$d(k\mathbf{x}) = \frac{\|\nabla f(k\mathbf{x})\|}{\gcd(\nabla f(k\mathbf{x}), \Delta_f)} = \frac{k^{d-1} \|\nabla f(\mathbf{x})\|}{\gcd(k^{d-1}, \Delta_f) \gcd(\nabla f(\mathbf{x}), \tilde{\Delta}_f)},$$

whence

$$S(\alpha, \beta) = \frac{k^{d-1}}{\gcd(k^{d-1}, \Delta_f)} \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n \\ \|\mathbf{x}\| \sim X}} \sum_{\mathbf{y} \in \mathbb{Z}^n} \frac{\|\nabla f(\mathbf{x})\|}{\gcd(\nabla f(\mathbf{x}), \tilde{\Delta}_f)} \omega(\mathbf{y}/Y) e(g(\mathbf{x}, \mathbf{y})). \quad (5.2)$$

In this section an important role will be played by the multilinear forms

$$m_j(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(d-1)}) = d! \sum_{j_1, \dots, j_{d-1}=1}^n c_{j_1, \dots, j_{d-1}, j} x_{j_1}^{(1)} \cdots x_{j_{d-1}}^{(d-1)},$$

for $1 \leq j \leq n$, where $c_{j_1, \dots, j_d} \in \mathbb{Z}$ are the symmetric coefficients such that

$$f(\mathbf{x}) = \sum_{j_1, \dots, j_d=1}^n c_{j_1, \dots, j_d} x_{j_1} \cdots x_{j_d}.$$

In what follows we shall write $\underline{\mathbf{u}}$ to denote the vector $(\mathbf{u}_1, \dots, \mathbf{u}_{d-1})$. Since f is non-singular, it follows from [2, Lemmas 3.1 and 3.3] that

$$\#\{\underline{\mathbf{u}} \in \mathbb{Z}^{(d-1)n} : \|\mathbf{u}_1\|, \dots, \|\mathbf{u}_{d-1}\| < U, m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) = 0 \forall j \leq n\} \ll 1 + U^{(d-2)n} \quad (5.3)$$

for any $U > 0$. Next, for given $P > 0$, $Q > 2$ and $\tau \in \mathbb{R}$, let

$$\mathcal{M}(\tau; P, Q) = \#\{\underline{\mathbf{u}} \in \mathbb{Z}^{(d-1)n} : \|\mathbf{u}_1\|, \dots, \|\mathbf{u}_{d-1}\| < P, \langle \tau m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) \rangle < Q^{-1} \forall j \leq n\}. \quad (5.4)$$

It follows from $d - 1$ applications of Lemma 5.1 that

$$\mathcal{M}(\tau; P, Q) \ll \frac{\max\{\sqrt{P/Q}, 1\}^{(d-1)n}}{\theta^{(d-1)n}} \mathcal{M}(\tau; \theta P, \theta^{1-d} Q), \quad (5.5)$$

for any $\theta \in (0, 1]$.

Returning to the expression for $S(\alpha, \beta)$ in (5.2), we start by removing the factor $\gcd(\nabla f(\mathbf{x}), \tilde{\Delta}_f)$ via the observation that $\gcd(\nabla f(\mathbf{x}), \tilde{\Delta}_f)$ depends only on $\mathbf{x} \bmod \tilde{\Delta}_f$. Letting $h = \tilde{\Delta}_f$ for compactness of notation, we break the sum into residue classes mod h , getting

$$S(\alpha, \beta) = \frac{k^{d-1}}{\gcd(k^{d-1}, \Delta_f)} \sum_{\mathbf{y} \in \mathbb{Z}^n} \omega(\mathbf{y}/Y) \sum_{\xi \in (\mathbb{Z}/h\mathbb{Z})^n} \frac{1}{\gcd(\nabla f(\xi), h)} T(\mathbf{y}), \quad (5.6)$$

where

$$T(\mathbf{y}) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ \|\mathbf{x}\| \sim X, \mathbf{x} \equiv \xi \bmod h}} \|\nabla f(\mathbf{x})\| e(g(\mathbf{x}, \mathbf{y})).$$

We may write

$$T(\mathbf{y}) = \sum_{\mathbf{x} \in \mathbb{Z}^n} F(\mathbf{x}) e(G(\mathbf{x})),$$

where

$$F(\mathbf{x}) = \begin{cases} \|(\nabla f)(\xi + h\mathbf{x})\| & \text{if } \|\xi + h\mathbf{x}\| \sim X, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$G(\mathbf{x}) = \alpha f(\xi + h\mathbf{x}) + \beta \mathbf{y} \cdot (\nabla f)(\xi + h\mathbf{x}).$$

Note that $G(\mathbf{x})$ has degree d .

We shall estimate $T(\mathbf{y})$ via Weyl differencing, as in Birch [2]. Let

$$\begin{aligned} F_{\mathbf{u}_1}(\mathbf{x}) &= F(\mathbf{x} + \mathbf{u}_1) F(\mathbf{x}), \\ F_{\mathbf{u}_1, \mathbf{u}_2}(\mathbf{x}) &= F(\mathbf{x} + \mathbf{u}_1 + \mathbf{u}_2) F(\mathbf{x} + \mathbf{u}_1) F(\mathbf{x} + \mathbf{u}_2) F(\mathbf{x}), \\ &\vdots \end{aligned}$$

and

$$\begin{aligned} G_{\mathbf{u}_1}(\mathbf{x}) &= G(\mathbf{x} + \mathbf{u}_1) - G(\mathbf{x}), \\ G_{\mathbf{u}_1, \mathbf{u}_2}(\mathbf{x}) &= G(\mathbf{x} + \mathbf{u}_1 + \mathbf{u}_2) - G(\mathbf{x} + \mathbf{u}_1) - G(\mathbf{x} + \mathbf{u}_2) + G(\mathbf{x}), \\ &\vdots \end{aligned}$$

Then, for any $r \in \{1, \dots, d-1\}$, we have

$$\left| \frac{T(\mathbf{y})}{X^n} \right|^{2^r} \ll \frac{1}{X^{rn}} \sum_{\|\mathbf{u}_1\| < X} \cdots \sum_{\|\mathbf{u}_r\| < X} \left| \frac{T_{\mathbf{u}_1, \dots, \mathbf{u}_r}(\mathbf{y})}{X^n} \right|, \quad (5.7)$$

where

$$T_{\mathbf{u}_1, \dots, \mathbf{u}_r}(\mathbf{y}) = \sum_{\mathbf{x} \in \mathbb{Z}^n} F_{\mathbf{u}_1, \dots, \mathbf{u}_r}(\mathbf{x}) e(G_{\mathbf{u}_1, \dots, \mathbf{u}_r}(\mathbf{x})).$$

We shall produce two estimates for $T(\mathbf{y})$. In the first we take $r = d - 1$, which eliminates the effect of the lower degree term $\beta \mathbf{y} \cdot (\nabla f)(\xi + h\mathbf{x})$ and leads to a family of linear exponential sums that depend on the Diophantine approximation properties of α alone. Alternatively, we take $r = d - 2$. After a further application of Cauchy–Schwarz, one brings the \mathbf{y} -sum inside, thereby bringing the Diophantine properties of β into play.

By Dirichlet’s approximation theorem there exist $a, q \in \mathbb{Z}$ and $\psi \in \mathbb{R}$ such that

$$\alpha = \frac{a}{q} + \psi,$$

with

$$\gcd(a, q) = 1, \quad 0 \leq a < q \leq X^{d/2} \quad \text{and} \quad |\psi| \leq \frac{1}{qX^{d/2}}. \quad (5.8)$$

The following is our first bound for $S(\alpha, \beta)$ and only involves the Diophantine approximation properties of α .

Lemma 5.3. *Assume that $\alpha = a/q + \psi$ is such that (5.8) holds and put*

$$D = \frac{n}{2^{d-1}(d-1)}.$$

Then

$$S(\alpha, \beta) \ll \frac{k^{d-1} X^{n+d-1} Y^n (\log X)^n}{q^D} \min \left\{ 1, \frac{1}{|\psi| X^d} \right\}^D.$$

Proof. Taking $r = d - 1$ in (5.7), we first note that

$$G_{\mathbf{u}_1, \dots, \mathbf{u}_{d-1}}(\mathbf{x}) = \alpha h^d \sum_{j=1}^n x_j m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) + H(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}),$$

for some polynomial $H(\mathbf{u}_1, \dots, \mathbf{u}_{d-1})$ that doesn’t depend on \mathbf{x} . It follows that

$$\sum_{\substack{\mathbf{x} \in \mathbb{Z}^n, \\ -cX \leq x_j \leq t_j}} e(G_{\mathbf{u}_1, \dots, \mathbf{u}_r}(\mathbf{x})) \ll \prod_{j=1}^n \min \{ X, \langle \alpha h^d m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) \rangle^{-1} \},$$

for any $t_1, \dots, t_n \ll X$ and any absolute constant $c > 0$. Exploiting (3.3), it readily follows from multi-dimensional partial summation that

$$T_{\mathbf{u}_1, \dots, \mathbf{u}_r}(\mathbf{y}) \ll X^{2^{d-1}(d-1)} \prod_{j=1}^n \min \{ X, \langle \alpha h^d m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) \rangle^{-1} \}.$$

In the standard way (cf. the proof of [6, Lemma 13.2]) one finds that

$$\left| \frac{T(\mathbf{y})}{X^n} \right|^{2^{d-1}} \ll X^{(d-1)(2^{d-1}-n)} (\log X)^n \mathcal{M}(\alpha h^d; X, X),$$

in the notation of (5.4). Applying (5.5) we obtain

$$\frac{\mathcal{M}(\alpha h^d; X, X)}{X^{(d-1)n}} \ll \frac{\mathcal{M}(\alpha h^d; \theta X, \theta^{1-d} X)}{(\theta X)^{(d-1)n}},$$

for any $\theta \in (0, 1]$. By choosing θ to satisfy

$$\theta^{d-1} \asymp \min \left\{ 1, \frac{1}{|q\psi|X^{d-1}}, \frac{X}{q}, \max \left\{ \frac{q}{X^{d-1}}, |q\psi|X \right\} \right\},$$

for appropriate implied constants depending on f , we can make Lemma 5.2 applicable. We then deduce from (5.3) that

$$\begin{aligned} \frac{\mathcal{M}(\alpha h^d; X, X)}{X^{(d-1)n}} &\ll \max \left\{ \frac{1}{X^{d-1}}, q|\psi|, \frac{q}{X^d}, \min \left\{ \frac{1}{q}, \frac{1}{q|\psi|X^d} \right\} \right\}^{n/(d-1)} \\ &\ll \frac{1}{q^{n/(d-1)}} \min \left\{ 1, \frac{1}{|\psi|X^d} \right\}^{n/(d-1)}, \end{aligned}$$

since (5.8) holds. It follows that

$$T(\mathbf{y}) \ll \frac{X^{n+d-1} (\log X)^n}{q^D} \min \left\{ 1, \frac{1}{|\psi|X^d} \right\}^D,$$

with D as in the statement of the lemma. Substituting this into (5.6), we conclude the proof of the lemma by summing trivially over \mathbf{y} and the finitely many possible values of ξ . \square

We now turn to our alternative estimate for $S(\alpha, \beta)$, which is obtained by exploiting the Diophantine approximation properties of β . By Dirichlet's approximation theorem there exist $b, r \in \mathbb{Z}$ and $\varrho \in \mathbb{R}$ such that

$$\beta = \frac{b}{r} + \varrho,$$

with

$$\gcd(b, r) = 1, \quad 0 \leq b < r \leq \sqrt{X^{d-1}Y} \quad \text{and} \quad |\varrho| \leq \frac{1}{r\sqrt{X^{d-1}Y}}. \quad (5.9)$$

We shall prove the following result, which operates under the assumption that X and Y are not too lopsided.

Lemma 5.4. Assume that $\beta = b/r + \varrho$ is such that (5.9) holds and put

$$E = \frac{n}{2^{d-2}(d-1)}.$$

Assume that $Y \leq X^{d-1}$. Then

$$S(\alpha, \beta) \ll \frac{k^{d-1} X^{n+d-1} Y^n \max\{X/Y, 1\}^{(d-1)n/2^{d-1}} (\log X)^n}{r^E} \min\left\{1, \frac{1}{|\varrho| X^{d-1} Y}\right\}^E$$

Proof. This time we begin through an application of Hölder's inequality in (5.6). Recalling that $h = O(1)$, we deduce that

$$\begin{aligned} \left| \frac{S(\alpha, \beta)}{k^{d-1}} \right|^{2^{d-2}} &\ll Y^{(2^{d-2}-1)n} \sum_{\|y\| \ll Y} \left(\sum_{\xi \in (\mathbb{Z}/h\mathbb{Z})^n} |T(y)| \right)^{2^{d-2}} \\ &\ll Y^{(2^{d-2}-1)n} \sum_{\xi \in (\mathbb{Z}/h\mathbb{Z})^n} U(\xi), \end{aligned}$$

where

$$U(\xi) = \sum_{\|y\| \ll Y} |T(y)|^{2^{d-2}}.$$

Taking $r = d - 3$ in (5.7), it follows that

$$\begin{aligned} U(\xi) &\ll \sum_{\|y\| \ll Y} \left| \frac{X^{2^{d-3}n}}{X^{(d-3)n}} \sum_{\|u_1\| < X} \cdots \sum_{\|u_{d-3}\| < X} \left| \frac{T_{u_1, \dots, u_{d-3}}(y)}{X^n} \right| \right|^2 \\ &\ll \frac{X^{2^{d-2}n}}{X^{(d-1)n}} \sum_{\|y\| \ll Y} \sum_{\|u_1\| < X} \cdots \sum_{\|u_{d-3}\| < X} |T_{u_1, \dots, u_{d-3}}(y)|^2. \end{aligned}$$

At this point we carry out a further differencing operation to conclude that

$$|T_{u_1, \dots, u_{d-3}}(y)|^2 = \sum_{u_{d-2} \in \mathbb{Z}^n} \sum_{u_{d-1} \in \mathbb{Z}^n} F_{u_1, \dots, u_{d-2}}(u_{d-1}) e(G_{u_1, \dots, u_{d-2}}(u_{d-1})).$$

There exists a polynomial $H \in \mathbb{Z}[u_1, \dots, u_{d-1}]$ that doesn't depend on y and polynomials $r_1, \dots, r_n \in \mathbb{Z}[u_1, \dots, u_{d-2}]$ that don't depend on u_{d-1} such that

$$\begin{aligned} G_{u_1, \dots, u_{d-2}}(u_{d-1}) &= \beta h^{d-1} \sum_{j=1}^n y_j (m_j(u_1, \dots, u_{d-1}) + r_j(u_1, \dots, u_{d-2})) \\ &\quad + H(u_1, \dots, u_{d-1}), \end{aligned}$$

where r_j may depend on ξ . We may now interchange the order of summation and execute the sum over y , noting that $F_{u_1, \dots, u_{d-2}}(u_{d-1}) \ll X^{2^{d-2}(d-1)}$ for $\|u_i\| < X$.

Hence, in the usual way, we conclude that

$$\begin{aligned} U(\xi) &\ll \frac{X^{2^{d-2}(n+d-1)}}{X^{(d-1)n}} \sum_{\|\mathbf{u}_1\| < X} \cdots \sum_{\|\mathbf{u}_{d-1}\| < X} \\ &\quad \times \prod_{j=1}^n \min \left\{ Y, \left\langle \beta h^{d-1} (m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) + r_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-2})) \right\rangle^{-1} \right\} \\ &\ll \frac{X^{2^{d-2}(n+d-1)} Y^n (\log Y)^n}{X^{(d-1)n}} \tilde{\mathcal{M}} \end{aligned}$$

where $\tilde{\mathcal{M}}$ denotes the number of $\underline{\mathbf{u}} \in \mathbb{Z}^{(d-1)n}$ for which $\|\mathbf{u}_1\|, \dots, \|\mathbf{u}_{d-1}\| < X$ and

$$\left\langle \beta h^{d-1} (m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) + r_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-2})) \right\rangle < Y^{-1},$$

for $1 \leq j \leq n$. Note that m_j is linear when viewed as a polynomial in \mathbf{u}_{d-1} . For fixed $\mathbf{u}_1, \dots, \mathbf{u}_{d-2}$, given a single \mathbf{u}'_{d-1} satisfying the inequality, for all other solutions \mathbf{u}_{d-1} we will have

$$\begin{aligned} &\left\langle \beta h^{d-1} (m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1} - \mathbf{u}'_{d-1})) \right\rangle \\ &= \left\langle \beta h^{d-1} (m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) + r_j - m_j(\mathbf{u}_1, \dots, \mathbf{u}'_{d-1}) - r_j) \right\rangle \\ &< 2Y^{-1}, \end{aligned}$$

where $r_j = r_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-2})$. Thus we may replace \mathbf{u}_{d-1} by $\mathbf{u}_{d-1} - \mathbf{u}'_{d-1}$ to remove the constant term $r_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-2})$. Doing so leads to the conclusion that $\tilde{\mathcal{M}}$ is at most the number of vectors $(\mathbf{u}_1, \dots, \mathbf{u}_{d-2}, \mathbf{u}_{d-1}) \in \mathbb{Z}^{(d-1)n}$ for which $\|\mathbf{u}_1\|, \dots, \|\mathbf{u}_{d-2}\| < X$ and $\|\mathbf{u}_{d-1}\| < 2X$, with

$$\left\langle \beta h^{d-1} m_j(\mathbf{u}_1, \dots, \mathbf{u}_{d-1}) \right\rangle < 2Y^{-1},$$

for $1 \leq j \leq n$. We conclude that

$$U(\xi) \ll \frac{X^{2^{d-2}(n+d-1)} Y^n (\log Y)^n}{X^{(d-1)n}} \mathcal{M}(\beta h^{d-1}; 2X, Y/2),$$

in the notation of (5.4).

Next, on appealing to (5.5), we deduce that

$$\mathcal{M}(\beta h^{d-1}; 2X, Y/2) \ll \frac{\max\{\sqrt{X/Y}, 1\}^{(d-1)n}}{\theta^{(d-1)n}} \mathcal{M}(\beta h^{d-1}; 2\theta X, \theta^{1-d} Y/2),$$

for any $\theta \in (0, 1]$. Assume that $\beta = b/r + \varrho$. If we choose θ so that

$$\theta^{d-1} \asymp \max \left\{ \frac{1}{X^{d-1}}, \min \left\{ 1, \frac{1}{|r\varrho|X^{d-1}}, \frac{Y}{r}, \max \left\{ \frac{r}{X^{d-1}}, |r\varrho|Y \right\} \right\} \right\},$$

for appropriate implied constants that depend only on f , then we can make Lemma 5.2 applicable. In the light of (5.3), this leads to the conclusion that

$$\mathcal{M}(\beta h^{d-1}; 2X, Y/2) \ll \frac{\max\{\sqrt{X/Y}, 1\}^{(d-1)n} (\theta X)^{(d-2)n}}{\theta^{(d-1)n}},$$

since $\theta X \gg 1$. Thus

$$\begin{aligned} U(\xi) &\ll \frac{X^{2^{d-2}(n+d-1)} Y^n \max\{\sqrt{X/Y}, 1\}^{(d-1)n} (\log Y)^n}{(\theta X)^n} \\ &\ll X^{2^{d-2}(n+d-1)} Y^n \max\{\sqrt{X/Y}, 1\}^{(d-1)n} (\log Y)^n M^{n/(d-1)}, \end{aligned}$$

where

$$M \ll \max \left\{ \frac{1}{X^{d-1}}, r|\varrho|, \frac{r}{YX^{d-1}}, \min \left\{ \frac{1}{r}, \frac{1}{r|\varrho|YX^{d-1}} \right\} \right\}.$$

Assuming that (5.9) holds and $Y \leq X^{d-1}$, it follows that

$$M \ll \frac{1}{r} \min \left\{ 1, \frac{1}{|\varrho|X^{d-1}Y} \right\},$$

whence finally

$$\begin{aligned} U(\xi) &\ll X^{2^{d-2}(n+d-1)} Y^n \max\{\sqrt{X/Y}, 1\}^{(d-1)n} (\log Y)^n \\ &\quad \times \frac{1}{r^{n/(d-1)}} \min \left\{ 1, \frac{1}{|\varrho|X^{d-1}Y} \right\}^{n/(d-1)}. \end{aligned}$$

We deduce by summing over the finitely many possible values of ξ that

$$\begin{aligned} \left| \frac{S(\alpha, \beta)}{k^{d-1}} \right|^{2^{d-2}} &\ll \frac{X^{2^{d-2}(n+d-1)} Y^{2^{d-2}n} \max\{\sqrt{X/Y}, 1\}^{(d-1)n} (\log Y)^n}{r^{n/(d-1)}} \\ &\quad \times \min \left\{ 1, \frac{1}{|\varrho|X^{d-1}Y} \right\}^{n/(d-1)}. \end{aligned}$$

The lemma follows since $\log Y \leq (d-1) \log X$. \square

We now have everything in place to complete the estimation of $E_{V,\varepsilon}^*(R)$ via Lemma 4.4. For the moment we continue to adopt the notation (5.1) for X and Y . Since $k \leq \sqrt{R}$ in Lemma 4.4 we may assume that $Y \leq X^{d-1}$ in Lemma 5.4. Given $Q_i, t_i > 0$, let $\mathcal{I}(Q_1, Q_2; t_1, t_2)$ denote the overall contribution to the integral

$$\int_0^1 \int_{\mathfrak{m}_{\eta,k}} |S(\alpha, \beta)| \, d\alpha \, d\beta$$

from $\alpha = a/q + \psi$ and $\beta = b/r + \varrho$ such that

$$q \sim Q_1, \quad r \sim Q_2 \quad \text{and} \quad |\psi| \sim t_1, \quad |\varrho| \sim t_2.$$

Then it follows that from Lemmas 5.3 and 5.4 that

$$\begin{aligned} \mathcal{I}(Q_1, Q_2; t_1, t_2) &\ll k^{d-1} X^{n+d-1} Y^n \max\{X/Y, 1\}^{(d-1)n/2^{d-1}} Q_1^2 Q_2^2 t_1 t_2 \\ &\times (\log X)^n \min\left\{\frac{1}{Q_1^D}, \frac{1}{(Q_1 t_1 X^d)^D}, \frac{1}{Q_2^E}, \frac{1}{(Q_2 t_2 X^{d-1} Y)^E}\right\}. \end{aligned} \quad (5.10)$$

By invoking Dirichlet's approximation theorem twice, as in (5.8) and (5.9), we see that we are only interested in $Q_i, t_i > 0$ such that

$$Q_1 \ll X^{d/2}, \quad Q_1 t_1 \ll \frac{1}{X^{d/2}} \quad \text{and} \quad Q_2 \ll \sqrt{X^{d-1} Y}, \quad Q_2 t_2 \ll \frac{1}{\sqrt{X^{d-1} Y}}.$$

Furthermore, since β belongs to the minor arcs $\mathfrak{m}_{\eta, k}$ it follows from Definition 4.1 that $\mathcal{I}(Q_1, Q_2; t_1, t_2) = 0$ unless

$$\max\{Q_2, Q_2 t_2 X^{d-1} Y\} \gg Y^{1-\eta}.$$

Since there are $O((\log XY)^4)$ possible dyadic values for Q_i, t_i that can contribute, we get an estimate for the minor arc integral by taking a maximum of (5.10) over all Q_i, t_i satisfying these inequalities.

Taking $\min\{A, B\} \leq A^{2/D} B^{1-2/D}$, with

$$A = \frac{1}{\max\{Q_1, Q_1 t_1 X^d\}^D} \quad \text{and} \quad B = \frac{1}{\max\{Q_2, Q_2 t_2 X^{d-1} Y\}^E},$$

and then taking $\max\{1, t_1 X^d\}^2 \geq t_1 X^d$, we deduce from (5.10) that

$$\begin{aligned} \mathcal{I}(Q_1, Q_2; t_1, t_2) &\ll k^{d-1} X^{n-1} Y^n \max\{X/Y, 1\}^{(d-1)n/2^{d-1}} (\log X)^n \\ &\times \frac{Q_2^2 t_2}{\max\{Q_2, Q_2 t_2 X^{d-1} Y\}^{E(1-2/D)}}. \end{aligned}$$

But $2E/D = 4$ and $Q_2^2 t_2 X^{d-1} Y \leq \max\{Q_2, Q_2 t_2 X^{d-1} Y\}^2$. Hence

$$\begin{aligned} \mathcal{I}(Q_1, Q_2; t_1, t_2) &\ll \frac{k^{d-1} X^{n-d} Y^{n-1} \max\{X/Y, 1\}^{(d-1)n/2^{d-1}} (\log X)^n}{\max\{Q_2, Q_2 t_2 X^{d-1} Y\}^{E-6}} \\ &\ll \frac{k^{d-1} X^{n-d} Y^{n-1} \max\{X/Y, 1\}^{(d-1)n/2^{d-1}} (\log X)^n}{Y^{(1-\eta)(E-6)}}. \end{aligned}$$

Note that the exponent of Y in the denominator is strictly positive precisely when $n > 3(d-1)2^{d-1}$. Recalling that X and Y are given by (5.1) we insert this argument into Lemma 4.4 to deduce that

$$E_{V, \varepsilon}^*(R) \ll R^{n-d-\delta}$$

for some $\delta > 0$, provided that ε is sufficiently small in terms of d and n . This completes the proof of Theorem 1.1, on summing over dyadic intervals in (3.6).

We can get an explicit value of the constant $c_{d,n}$ as follows. Since $Y \geq X^{1-\varepsilon}$ in (5.1), we see that

$$\frac{X^{n-d} Y^{n-1} \max\{X/Y, 1\}^{(d-1)n/2^{d-1}} (\log X)^n}{Y^{(1-\eta)(E-6)}} \leq \frac{X^{n-d} Y^{n-1} X^{\varepsilon(d-1)n/2^{d-1}} (\log X)^n}{X^{(1-\varepsilon)(1-\eta)(E-6)}}$$

gives a power saving as soon as $(1-\varepsilon)(1-\eta)(E-6) > \varepsilon(d-1)n/2^{d-1}$. Recalling that $E = n/(2^{d-2}(d-1))$ and multiplying both sides by $2^{d-1}(d-1)$, this condition becomes

$$(1-\varepsilon)(1-\eta)(2n-3(d-1)2^d) > \varepsilon n(d-1)^2$$

or

$$\varepsilon < \frac{2n-3(d-1)2^d}{n(d-1)^2/(1-\eta) + 2n-3(d-1)2^d}.$$

Thus we may take $c_{d,n} = \frac{2n-3(d-1)2^d}{n(d^2-2d+3)-3(d-1)2^d}$ in Theorem 1.1 by letting η converge to 0. Note that for fixed d we have $c_{d,n} \rightarrow \frac{2}{d^2-2d+3}$ as $n \rightarrow \infty$.

References

- [1] W. Banaszczyk, New bounds in some transference theorems in the geometry of numbers, *Math. Ann.*, **296** (1993), no. 4, 625–635. Zbl 0786.11035 MR 1233487
- [2] B. J. Birch, Forms in many variables, *Proc. Roy. Soc. London Ser. A*, **265** (1961/62), 245–263. Zbl 0103.03102 MR 150129
- [3] J.-B. Bost, Algebraic leaves of algebraic foliations over number fields, *Publ. Math. Inst. Hautes Études Sci.*, (2001), no. 93, 161–221. Zbl 1034.14010 MR 1863738
- [4] T. D. Browning and W. Sawin, Free rational curves on low degree hypersurfaces and the circle method, 2018. arXiv:1810.06882
- [5] J. W. S. Cassels, *An introduction to the geometry of numbers*. Second printing, corrected, Die Grundlehren der mathematischen Wissenschaften, Band 99, Springer-Verlag, Berlin-New York, 1971. Zbl 0209.34401 MR 306130
- [6] H. Davenport, *Analytic methods for Diophantine equations and Diophantine inequalities*. Second edition. With a foreword by R. C. Vaughan, D. R. Heath-Brown and D. E. Freeman. Edited and prepared for publication by T. D. Browning, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2005. Zbl 1125.11018 MR 2152164
- [7] J. Franke, Y. I. Manin, and Y. Tschinkel, Rational points of bounded height on Fano varieties, *Invent. Math.*, **95** (1989), no. 2, 421–435. Zbl 0674.14012 MR 974910
- [8] D. R. Heath-Brown, Diophantine approximation with square-free numbers, *Math. Z.*, **187** (1984), no. 3, 335–344. Zbl 0539.10026 MR 757475
- [9] D. R. Heath-Brown, Cubic forms in 14 variables, *Invent. Math.*, **170** (2007), no. 1, 199–230. Zbl 1135.11031 MR 2336082

- [10] E. Peyre, Hauteurs et mesures de Tamagawa sur les variétés de Fano (French), *Duke Math. J.*, **79** (1995), no. 1, 101–218. Zbl 0901.14025 MR 1340296
- [11] E. Peyre, Liberté et accumulation, *Doc. Math.*, **22** (2017), 1615–1659. Zbl 1434.11084 MR 3741845

Received June 24, 2019

T. Browning, IST Austria, Am Campus 1,
3400 Klosterneuburg, Austria

E-mail: tdb@ist.ac.at

W. Sawin, Columbia University, Department of Mathematics,
2990 Broadway, New York, NY 10027, USA

E-mail: sawin@math.columbia.edu

