

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 42 (1967)

Artikel: On Finitely Generated Fuchsian Groups.
Autor: Marden, Albert
DOI: <https://doi.org/10.5169/seals-32132>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On Finitely Generated Fuchsian Groups¹⁾

by ALBERT MARDEN

We will prove the following two theorems.

THEOREM 1. *Let G be a finitely generated Fuchsian group in the unit disk Δ . Then a) $\Omega = \Delta/G$ is a Riemann surface of finite topological type, and b) Δ is ramified over at most a finite number of points $\{p_i\}$ of Ω . Conversely, if G satisfies a) and b), then G is finitely generated.*

THEOREM 2. *G is finitely generated if and only if every fundamental region P has a finite number of sides.*

DEFINITION. A fundamental region P for G is a connected open set in Δ which satisfies the following conditions.

1. Every point in Δ is equivalent under G to a point in \bar{P} (= relative closure in Δ).
2. No two points in P are equivalent.
3. Each component of the relative boundary of P in Δ is an open Jordan arc or a Jordan curve and is the union of possibly infinitely many closed Jordan arcs, called sides (two sides can intersect only at a common end point).
4. The sides of P are arranged in pairs (s_i, s'_i) where (i) $S_i(s'_i) = s_i$ for some $S_i \in G$, (ii) $S_i \neq S_j^{\pm 1}$ for each $j \neq i$ with at most a finite number of exceptions, (iii) each side of P appears once and only once in the set $\{s_i, s'_i\}$.

Theorem 1 is well known and is fundamental in the theory of Fuchsian groups. By the use of variational methods it has been proven by AHLFORS [1] (in a more general form), BERS [2], and EARLE [3]. From a more general point of view it is a consequence of a theorem of SELBERG [8]. Theorem 2 is also known provided P satisfies the additional hypotheses that (a) *its sides are non-Euclidean line segments*, and (b) *only a finite number of images of P under G meet any given compact set in Δ* . In this form, a proof has recently been given by L. GREENBERG [5]. M. HEINS' proof [6] requires that P also be convex (in this paper HEINS also proves Theorem 1). The first proof of Theorem 1 and of Theorem 2 in the case of Poincaré normal polygons was given by FENCHEL and NIELSEN [4].

The purpose of this paper is to give direct, elementary proofs of Theorems 1 and 2 which are much shorter than those referred to above, and in the case of Theorem 2, more general as well. In fact our definition of fundamental region is, in a sense, the weakest one for which Theorem 2 is true: If P does not satisfy the principal condition 4

¹⁾ Supported in part by the National Science Foundation under grants GP 2280 and GP 3904.

(ii) then Theorem 2 is false in general. Our proofs are purely topological in character and make use of some elementary properties of surfaces of finite topological type. We will show that Theorem 1b is a simple consequence of the fact that a cycle in the exterior of a region which is homologous to a cycle in the interior is homologous to a cycle on the boundary. And Theorem 2 is a consequence of the fact that if there are infinitely many mutually disjoint simple closed curves not ~ 1 on a surface of finite type, then two of them bound an annular region.

Proof of a). Suppose A_1, \dots, A_m generate G and let π denote the projection $\Delta \rightarrow \Omega$. Assuming 0 is not a fixed point of G , denote by $[0, A_i(0)]$ the non-Euclidean line segment from 0 to $A_i(0)$ (actually any arc will do) and set $\alpha_i = \pi([0, A_i(0)])$, $1 \leq i \leq m$. We claim that the curves $\{\alpha_i\}$ generate the fundamental group of Ω with origin at $\pi(0)$ and consequently that Ω is of finite topological type.

If τ is a closed curve in Ω with initial point $\pi(0)$ there is a lift τ^* of τ in Δ with initial point 0 and end point $\tau^*(1)$. We may write $\tau^*(1) = B_m B_{m-1} \dots B_1(0)$ where each B_j is some $A_i^{\pm 1}$. Consider the arc τ'^* in Δ from 0 to $\tau^*(1)$ obtained by joining the non-Euclidean line segments $[0, B_m(0)]$, $[B_m(0), B_m B_{m-1}(0)]$, \dots , $[B_m \dots B_2(0), B_m \dots B_1(0)]$; each of these segments projects onto some curve $\alpha_i^{\pm 1}$. Since τ^* is homotopic to τ'^* , τ is homotopic to $\pi(\tau'^*)$, that is τ is homotopic to a product of the α_i .

Proof of b). Let Ω_0 be a relatively compact subregion of Ω containing all the curves α_i such that each component of $\partial\Omega_0$ is a dividing cycle and no component of $\Omega - \Omega_0$ is compact. We claim that Δ is not ramified over $\Omega' = \Omega - \bar{\Omega}_0$.

Assume to the contrary that Ω is ramified of order $t \geq 2$ over $p \in \Omega'$. Let c_p be the oriented boundary of a disk about p in Ω' , d_p a Jordan arc from $\pi(0)$ to c_p , and β the closed path from $\pi(0)$ along d_p to c_p , around c_p once, and back to $\pi(0)$ along d_p . We may assume that β does not pass through points over which Δ is ramified and intersects the curves α_i only a finite number of times.

Let β^* denote the lift of β from 0; β^* is a Jordan arc but not a closed curve in Δ . We have seen above that β^* is homotopic to an arc β'^* such that $\beta' = \pi(\beta'^*)$ is a product of curves α_i . Consider the closed curve $\gamma^* = \beta'^* \beta^{*-1}$ and a relatively compact, simply connected region K containing γ^* with $\pi(\partial K) \cap c_p = \emptyset$ (the points $\pi^{-1}(p)$ are isolated in Δ). Remove from K the at most finite number of points which lie over p and denote the resulting region by K_1 .

The collection of disjoint simple closed curves which comprise the components of $\pi^{-1}(c_p)$ in K_1 form a homology basis for K_1 . Hence γ^* , viewed as a singular cycle, is homologous to a linear combination of these curves. In $\Omega - \{p\}$ this implies that $\gamma = \pi(\gamma^*) = \beta' \beta^{-1}$ is homologous to ntc_p for some integer n , possibly zero. In other words the cycle $(nt+1)c_p$ in $\Omega - \Omega_0$, with $nt+1 \neq 0$ since $t \geq 2$, is homologous in $\Omega - \{p\}$ to the cycle β' in Ω_0 . Therefore $(nt+1)c_p$ must be homologous to a cycle in $\partial\Omega_0$ in contradiction to our choice of Ω_0 .

The proof of the converse of Theorem 1 is standard and will be omitted.

Proof of Theorem 2. Let P be a fundamental region. Since G is generated by the transformations which pair the sides of P , the sufficiency is immediate.

Assume then that G is finitely generated but that P has an infinite number of sides. \bar{P} is not compact in Δ for otherwise there would exist a sequence of points $\{z_j\}$ on distinct sides s_j of P which have a limit $p \in \Delta$ such that the sequence of conjugate points $\{z'_j\}$ on the conjugate sides s'_j also has a limit $p' \in \Delta$. If N is a neighborhood of p and N' of p' then for infinitely many distinct $S_j \in G$, $S_j(N') \cap N \neq \emptyset$, in violation of the discontinuity of G . In addition P is simply connected for otherwise there would be a relatively compact component of $\Delta - \bar{P}$ which would contain an image of P .

The hypotheses also imply that $S_i(P)$ is adjacent to P along s_i . Hence if $j \neq i$, $S_j(\bar{P})$ cannot intersect s_i at an interior point unless $S_j = S_i$.

In the remainder of the proof all arcs drawn in a region $\bar{R} \subset \bar{P}$ will be understood to be Jordan arcs which are contained in R except for their end points. In addition we can choose, and will only deal with, an infinite sequence of pairs $\{(s_i, s'_i)\}$ such that $S_i \neq S_j^{\pm 1}$ for all $j \neq i$.

CASE 1. There exists an arc τ in \bar{P} which divides P into two regions P_1, P_2 with the following property. There exists an infinite number of points $\{z_i\}$ in \bar{P}_1 , each z_i an interior point of a side of P , but no two points on the same side, such that the conjugate points $\{z'_i\}$ are in \bar{P}_2 .

Draw arcs γ_1 in \bar{P}_1 from z_1 to z_2 and γ'_1 in \bar{P}_2 from z'_2 to z'_1 . γ_1 divides P_1 into two regions at least one of which, say P_{11} , contains infinitely many points z_i in its boundary, and γ'_1 divides P_2 into two regions one of which, say P_{21} , contains infinitely many points conjugate to those z_i in \bar{P}_{11} . Eliminate all z_i which are not in P_{11} and which do not have their conjugates in P_{21} . Draw the arcs γ_2 in \bar{P}_{11} from z_3 to z_4 and γ'_2 in \bar{P}_{21} from z'_4 to z'_3 , etc. Thus we can find two infinite sequences of mutually disjoint arcs $\{\gamma_i\}$ in \bar{P}_1 and $\{\gamma'_i\}$ in \bar{P}_2 such that γ_i is an arc from z_{2i-1} to z_{2i} and γ'_i an arc from z'_{2i} to z'_{2i-1} .

In addition by a suitable choice of subsequence we may assume that either property A holds for all γ_i, γ_j or property A holds for no γ_i, γ_j in the infinite sequence $\{\gamma_i\}$:

PROPERTY A. γ_i separates γ_j from τ or γ_j separates γ_i from τ ($i \neq j$)

To prove this apply the following procedure inductively. Assume that by relabeling, all the arcs γ_j , $0 < j \leq n$, have property A with respect to all the arcs in $\{\gamma_i\}$, $1 \leq i \leq \infty$, but none of the arcs δ_j , $0 < j \leq m$, has property A with respect to any of the arcs in $\{\gamma_i, \delta_k\}$, $1 \leq i < \infty$, $0 < k \leq m$ ($m, n \geq 0$). Consider γ_{n+1} . If infinitely many arcs in $\{\gamma_i\}$ have property A with respect to γ_{n+1} (the arcs γ_i , $0 < i \leq n$, do), eliminate those arcs in $\{\gamma_i\}$ which do not, relabel, and move on to γ_{n+2} . If this is not the case, eliminate the finite number of arcs in $\{\gamma_i\}$ which do have property A with respect to γ_{n+1} , set $\delta_{m+1} = \gamma_{n+1}$, relabel, and move on to the new γ_1 .

Consider the sequence of mutually disjoint simple closed curves $\{\alpha_i = \pi(\gamma_i \cup \gamma'_i)\}$ in Ω . No α_i can be homotopic to 1 in $\Omega - \{p_i\}$. For otherwise $\gamma_i \cup S_{2i}(\gamma'_i)$ is a simple

closed curve through z_{2i-1} and z_{2i} and hence $S_{2i-1} = S_{2i}$, which is impossible. Therefore since $\Omega - \{p_i\}$ has finite topological type, there exist three of the curves α_i , say $\alpha_1, \alpha_2, \alpha_3$, such that α_1 and α_2 bound an annular region K in $\Omega - \{p_i\}$ and α_3 separates the contours of K . We can also choose K so that $\pi(\tau) \cap K = \emptyset$.

The arcs γ_1, γ_2 divide P_1 into three regions. If two of these regions lie over K then γ_3 is contained in one of them, but not in the other, so that γ_3 satisfies property A with only one of the arcs γ_1, γ_2 in contradiction to our selection of $\{\gamma_i\}$. We conclude that γ_1 and γ_2 bound a region R in P_1 which lies over K and contains γ_3 .

In P_2 , γ'_1 and γ'_2 bound one or two regions which lie over K . One of these regions contains γ'_3 and therefore γ'_3 bounds a region R'_1 with, say, γ'_1 which lies over $K(\pi(R'_1) \subset K)$. We can draw an arc δ' in R'_1 from z'_3 to one of z'_1, z'_2 , say to z'_1 , such that δ' does not separate γ'_1 from γ'_3 in R'_1 . In R draw an arc δ from z_1 to z_3 which does not otherwise intersect γ_3 .

In the annular region $K_1 \subset K$ which is bounded by α_1 and α_3 , the simple closed curve $\beta = \pi(\delta \cup \delta')$ does not separate α_1 and α_3 . Hence β is homotopic to 1 in K_1 which we have seen above, is impossible.

CASE 2. No such τ exists. In this case draw an arc γ_1 from z_1 to z'_1 . There exist infinitely many points $\{z_i\}$ such that γ_1 does not separate z_i from z'_i in P . Draw γ_2 from z_2 to z'_2 which is disjoint from γ_1 . Again there are infinitely many points $\{z_i\}$ such that neither γ_1 nor γ_2 separates z_i from z'_i . Thus we can find an infinite sequence of disjoint arcs $\{\gamma_i\}$ such that γ_i runs from z_i to z'_i .

The simple closed curves $\alpha_i = \pi(\gamma_i)$ are mutually disjoint and hence, as above, there are three of them, say $\alpha_1, \alpha_2, \alpha_3$, such that α_1 and α_2 bound an annular region K in $\Omega - \{p_i\}$ and α_3 separates the boundary components of K . γ_1 and γ_2 divide P into three regions, one or two of which lie over K and of these, one contains γ_3 . Hence one of the pairs $(\gamma_1, \gamma_2), (\gamma_1, \gamma_3), (\gamma_2, \gamma_3)$, say (γ_1, γ_2) , bounds a subregion R of P which lies over K . Interchange z_2 and z'_2 if necessary so that α_1 is homologous to α_2 . Draw an arc δ in \bar{R} from z'_1 to z'_2 ; then α_1 is homotopic to $\pi(\delta) \alpha_2 \pi(\delta)^{-1}$. But then the arc $S_2(\delta^{-1})$ in $\Delta - P$ from z_2 must terminate at z_1 which implies that $S_2 = S_1$, a contradiction.

Remark. If P also satisfies hypothesis (b) a much simpler proof can be given. The sequence $\{z_n\}$ can be chosen so that $\{\pi(z_n)\}$ approaches an ideal boundary component I of Ω . By using the fact that if α is a simple closed curve surrounding I then $\pi^{-1}(\alpha)$ divides P into a finite number of components, the arcs γ_i, γ'_i of Cases 1 and 2 can be chosen directly so that the curves α_i approach I in such a way that α_i bounds an annular region A_i in $\Omega - \{p_i\}$ with α_{i+1} that doesn't contain α_j for $j > i + 1$. It follows that the pairs $(\gamma_i, \gamma_{i+1}), (\gamma'_i, \gamma'_{i+1})$ (just the former in Case 2) each bound a region in P which lies over A_i , since α_{i+2} cannot be connected to α_i without crossing α_{i+1} . The proof is now completed as above.

BIBLIOGRAPHY

- [1] L. V. AHLFORS, *Finitely generated Kleinian groups*, Amer. J. Math. 86 (1964), 413–429.
- [2] L. BERS, *Automorphic forms and Poincaré series for infinitely generated Fuchsian groups*, Amer. J. Math. 87 (1965), 196–214.
- [3] C. EARLE, *Reduced Teichmüller spaces*, Trans. Amer. Math. Soc., to appear.
- [4] W. FENCHEL and J. NIELSEN, *Discontinuous Groups of non-Euclidean Motions*.
- [5] L. GREENBERG, *Fundamental polygons for Fuchsian groups*, to appear.
- [6] M. HEINS, *Fundamental polygons of Fuchsian and Fuchsoid groups*, Ann. Acad. Sci. Fenn. Ser. A, 337 (1964), 1–30. Bull. Amer. Math. Soc. 69 (1963), 747–751.
- [7] J. LEHNER, *Discontinuous Groups and Automorphic Functions*, Mathematical Surveys 8, Amer. Math. Soc., Providence (1964).
- [8] A. SELBERG, *On discontinuous groups in higher dimensional symmetric spaces*, Contributions to Function Theory, Bombay (1960), 147–164.

University of Minnesota

Received July 27, 1966