

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 6 (1951)  
**Heft:** 2

**Artikel:** The Osculating Conics of Steiner`s Hypocycloid  
**Autor:** Fabricius-Bjerre, Fr.  
**DOI:** <https://doi.org/10.5169/seals-15573>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)


**Download PDF:** 14.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## The Osculating Conics of Steiner's Hypocycloid

In the present note we shall give a simple construction of the osculating conic at an arbitrary point of Steiner's hypocycloid and determine the locus of the centres of these conics.

Let  $O$  be the centre of a fixed circle with radius  $3a$  and  $O_1$  the centre of a circle with radius  $a$ , which is rolling inside the first circle. The angle from an  $x$ -axis through  $O$  to the line  $OO_1$  is denoted by  $\theta$  and the corresponding point of Steiner's hypocycloid  $\mathfrak{H}$  is called  $P$  (figure). The line  $OO_1$  cuts the rolling circle in the points  $T$  and  $N$  such



that  $TP$  is the tangent and  $NP$  the normal to  $\mathfrak{H}$  at  $P$ . Since  $\sphericalangle (NO_1P) = 3\theta$  we have  $\sphericalangle (O_1TP) = \sphericalangle (O_1PT) = 3\theta/2$ . Furthermore, it is known that the radius of curvature  $\rho$  of  $\mathfrak{H}$  at  $P$  is

$$\rho = 4 PN = 8a \sin \frac{3}{2} \theta.$$

To determine the osculating conic of  $\mathfrak{H}$  at  $P$  we use some results from the affine geometry<sup>1)</sup>.

The *affine normal* (or axis of deviation) at  $P$  can be constructed in the following manner (l. c., p. 33): On the usual normal  $PN$  we find the centre of curvature  $P_1$  corresponding to  $P$  and again for the evolute of  $\mathfrak{H}$  the centre of curvature  $P_2$  corresponding to  $P_1$ . On the line  $P_1P_2$  which is parallel to the tangent  $TP$  we determine a point  $P_3$  such that  $P_3P_1 = P_3P_2/4$ . The line joining  $P$  and  $P_3$  is the required affine normal.

In consequence of a well-known property of the cycloids, the point  $P_2$  is situated on the line  $OP$ . Consider a normal to  $PN$  through  $O_1$  which cuts  $PN$  in a point  $P_1'$  and  $OP$  in a point  $P_2'$ . From the figure we derive that  $O_1P_1' = TP/2 = O_1P_2'/4$ . Hence the line  $PO_1$  passes through  $P_3$ , i. e.  $PO_1$  is the affine normal at  $P$ . We then have

*The affine normal at a point  $P$  of Steiner's hypocycloid is the line which joins  $P$  with the centre of the rolling circle.*

<sup>3)</sup> W. BLASCHKE, *Vorlesungen über Differentialgeometrie*, vol. 2 (Springer, Berlin 1923).

The envelope of the affine normal is the so-called affine evolute of  $\mathfrak{S}$ , and the point of contact  $C$  is the centre of the osculating conic at  $P$  (l. c., p. 28). Since  $C$  can be found by drawing a perpendicular from  $N$  to the line  $PO_1$ , the locus of  $C$  is a new hypocycloid where the diameter of the rolling circle is the segment  $NO_1$ .

*The locus of the centres of the osculating conics for the hypocycloid  $\mathfrak{S}$  is a new hypocycloid generated by a point of a circle with radius  $a/2$  which is rolling inside the fixed circle with radius  $3a^1$ .*

This hypocycloid consists of 6 equal arcs. Every second cusp coincides with a cusp of  $\mathfrak{S}$ .

All the osculating conics will be *ellipses*, because the centre  $C$  lies on the same side of the tangent  $TP$  as the arc of  $\mathfrak{S}$  which contains  $P$ . Of the ellipse that osculates  $\mathfrak{S}$  at  $P$  we have found the centre  $C$  and a semi-diameter  $CP$  with the length  $CP = 2a \sin^2(3\theta/2)$ . Let  $Q$  be the extreme point of the conjugate semi-diameter  $CQ$  parallel to  $TP$ . At  $P$  the osculating ellipse has the same radius of curvature  $\rho$  as  $\mathfrak{S}$ , and since the radius of curvature of the ellipse at  $P$  can be expressed by  $CQ^2/CP \sin C$ , where  $\sphericalangle C = 3\theta/2$ , we get

$$8a \sin \frac{3}{2} \theta = \frac{CQ^2}{2a \sin^2 \frac{3}{2} \theta \sin \frac{3}{2} \theta}$$

or 
$$CQ = 4a \sin^2 \frac{3}{2} \theta = 2CP.$$

*The length of the diameter parallel to the tangent  $TP$  is twice the diameter through  $P$ .*

We thus have obtained the required construction of the osculating ellipse at the point  $P$  of  $\mathfrak{S}$ .

In a similar way we may obtain the osculating ellipses of a general hypo- or epicycloid. The construction, however, will not be as simple as here, because the first theorem mentioned above can not be extended to other cycloids.

At last we notice that the reciprocal curve of  $\mathfrak{S}$  with respect to the circle with centre  $O$  and radius  $a$  is a *cubic  $\mathfrak{S}'$* , which consists of three open convex arcs situated in angles of  $60^\circ$  and having the same vertices as  $\mathfrak{S}$ . By a dualistic transformation an osculating conic will be transformed into an osculating conic. The point  $O$  being outside all the osculating ellipses of  $\mathfrak{S}$ , it is obvious that each osculating conic of the cubic  $\mathfrak{S}'$  will be a *hyperbola*.

FR. FABRICIUS-BJERRE, Copenhagen.

## La mode dans les mathématiques

La théorie élémentaire des séries insiste sur le rôle secondaire que jouent les premiers termes d'une série. Ce sont les termes de haut rang, leur allure, leur caractère, qui signifient tout.

Pourtant, la théorie des fonctions analytiques nous montre que ce point de vue, que nous inculquons à nos élèves de première année, peut induire en erreur. Le

<sup>1</sup>) J. LEMAIRE, *Hypocycloïdes et Epicycloïdes* (Paris 1929), p. 55.