

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 24 (1969)  
**Heft:** 4

**Artikel:** On Sc Functions  
**Autor:** Tauber, S.  
**DOI:** <https://doi.org/10.5169/seals-26649>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 21.05.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 9. Schlussbetrachtung

Ziel der Arbeit ist es, an Beispielen eine Methode vorzuführen, die es gestattet, die Inhaltsmasszahlen gewisser hyperbolischer Rotationskörper zu ermitteln. Die Auswahl der Beispiele erfolgte nach zwei Gesichtspunkten. Einmal sollten alle in Teil 3 ausgewerteten Integrale bei der Bestimmung von Masszahlen tatsächlich vorkommen. Zum andern aber wollten wir vor allem solche Drehkörper behandeln, die in der euklidischen Geometrie ein Gegenstück haben. Es ist besonders interessant, in diesen Fällen den Übergang von der hyperbolischen zur euklidischen Formel durchzuführen. Wir zeigen das am Beispiel des Torus. In Formel (16) werden die vorkommenden hyperbolischen Funktionen in Reihen entwickelt:

$$V = \pi^2 k^3 \left( \frac{\bar{a}}{k} + \frac{1}{3!} \left( \frac{\bar{a}}{k} \right)^3 + \dots \right) \left( 1 + \frac{1}{2} \left( \frac{2 \bar{R}}{k} \right)^2 + \dots - 1 \right)$$

Wächst jetzt  $k$  unbegrenzt, so ergibt sich das Torusvolumen der euklidischen Geometrie  $V = \pi^2 a \cdot 2 R^2$ .

H.ZEITLER, Weiden

# On $Sc$ Functions

## Introduction

In this paper we prove that the Dirac Delta and all its derivatives can be represented by sequences of constructed discontinuous functions. Although this result is stated in [1] it is not formally proved.

We then prove that by using this definition of the  $n$ -th derivative of the Dirac Delta its Laplace Transform is  $s^n$ . This result again can be considered as “classical” (see for example [3]) but is not proved either.

We feel that although the results are known the approach is new and our proof is rigorous which justifies the contents of this paper.

### Definition of the $n$ -th Derivative of a Function

Let  $\mathbf{V} = [v_1, v_2, \dots, v_n]$  be an  $n$ -dimensional vector. We say that the vector tends *basewise* to zero if the components  $v_k$  tend to zero *successively*. We write symbolically

$$\mathbf{V} \xrightarrow[B]{} 0 . \quad (1)$$

Geometrically speaking this means that the end point of  $\mathbf{V}$  describes a polygonal line whose sides are parallel to the axes of the basis.

We shall use the notation  $\Pi \mathbf{V} = \prod_{m=1}^n v_m$  for the product of the components of the vector.

Let  $f(t) \in C^n[b, c]$  be the class of functions that are defined and continuous as well as their derivatives up to and including the order  $n$  for  $b \leqq t \leqq c$ . Let  $a_k$ ,  $k = 1, 2, \dots, n$ , be such that  $(t + \alpha_{h,n}) \in [b, c]$ , where  $\alpha_{h,n}$  represents the sum of any  $h$  of the  $n$

numbers  $a_k$ , and  $b + \varepsilon \leq t \leq c - \varepsilon$ ,  $\varepsilon$  being a given positive number. Clearly

where,

$$a = [a_1, a_2, \dots, a_n], \quad \varphi_k = \sum_{m=1}^{\binom{n}{k}} f(t + \alpha_{k,m}), \quad \alpha_{k,m} = a_{s_1} + a_{s_2} + \dots + a_{s_k},$$

$$s_i, s_i = 1, 2, \dots, n, \quad s_i \neq s_j, \quad i, j = 1, 2, \dots, k.$$

Since  $\varphi_k$  is symmetric with respect to the  $a_k$ 's,  $D^n f(t)$  is independent of the order in which the different  $a_k$ 's tend to zero, this is why no specific order is necessary when a vector tends basewise to zero.

If in  $a \rightarrow 0$  we make a change of variables in the  $a_k$ 's, change defined by

$$\mathbf{b} = H \mathbf{a}, \quad (3)$$

where  $H$  is a  $n \times n$  matrix, this change of variables corresponds to a rotation of the reference system. When  $\mathbf{a} \xrightarrow[B]{} 0$ , the last leg of the polygonal line described by the endpoint of  $\mathbf{a}$  is a straight line. With respect to the new reference system, when  $\mathbf{a}$  tends to zero basewise then  $\mathbf{b}$  tends to zero, although *not* basewise. Under these conditions, considering (3) all the  $b_k$ 's, components of  $\mathbf{b}$  will tend to zero *simultaneously*.

We may thus assume without loss of generality that in (2) all the  $a_k$ 's tend to zero simultaneously. In addition we may assume that all the  $a_k$ 's are equal to  $a$ . This corresponds to a special choice of the matrix  $H$  in (3) that would make all the  $b_k$ 's equal.

It follows that we can define the  $n$ -th derivative of the function  $f(t)$  by the expression

$$D^n f(t) = \lim_{a \rightarrow 0} a^{-n} \sum_{k=0}^n (-1)^{k+n} \binom{n}{k} f(t + k a). \quad (4)$$

If  $0 \leq \theta \leq 1$ , and  $0 \leq \beta_k \leq 1$ ,  $k = 1, 2, \dots, n$ , we can write according to the mean-value theorem

$$f[f + (k + \theta) a] = f(t + k a) + a \theta Df[t + (k + \beta_k \theta) a],$$

so that by substitution into (4) we obtain

$$D^n f(t) = \lim_{a \rightarrow 0} a^{-n} \sum_{k=0}^n (-1)^{n+k} [f(t + \theta) a] - a \theta Df[t + (k + \beta_k \theta) a],$$

where the second term in the sum tends to zero with  $a$ .

We can thus write for the  $n$ -th derivative of  $f(t)$

$$D^n f(t) = \lim_{a \rightarrow 0} (-1)^n a^{-n} \sum_{k=0}^n (-1)^k \binom{n}{k} f[t + (k + \theta) a]. \quad (5)$$

This is the form we are going to use in the next section.

*Remark.* (5) can easily be checked by writing

$$\begin{aligned} f[t + (k + \theta) a] &= f(t) + \sum_{m=1}^{n-1} (m!)^{-1} a^m (k + \theta)^m D^m f(t) \\ &\quad + (k + \theta)^n a^n D^n f[t + \beta_k (k + \theta) a]/n!, \quad 0 \leq \beta_k \leq 1, \end{aligned}$$

thus substituting into (5)

$$\begin{aligned} D^n f(t) &= \lim_{a \rightarrow 0} (-1)^n a^{-n} \sum_{k=0}^n (-1)^k \binom{n}{k} [f(t) + \sum_{m=0}^{n-1} (m!)^{-1} a^m (k + \theta)^m D^m f(t) \\ &\quad + (n!)^{-1} a^n (k + \theta)^n D^n f[t + \beta_k (k + \theta) a]] \\ &= \lim_{a \rightarrow 0} (-1)^n a^{-n} \left\{ f(t) \sum_{k=0}^n (-1)^k \binom{n}{k} + \sum_{m=1}^{n-1} (m!)^{-1} a^m D^m f(t) \sum_{k=0}^n (-1)^k \binom{n}{k} (k + \theta)^m \right. \\ &\quad \left. + (n!)^{-1} a^n D^n f[t + \beta_k (k + \theta) a] \sum_{k=0}^n (-1)^k \binom{n}{k} (k + \theta)^n \right\}. \end{aligned}$$

Since

$$\sum_{k=0}^n (-1)^k \binom{n}{k} (k + \theta)^m = (-1)^n n! \delta_n^m,$$

where  $\delta_n^m$  is the Kronecker Delta, it follows that all the terms on the right hand side, except the last one, cancel out, so that

$$D^n f(t) = \lim_{a \rightarrow 0} (-1)^n a^{-n} (n!)^{-1} a^n D^n f[t + \beta_k (k + \theta) a] n! (-1)^n,$$

which clearly is an identity.

### 3. The accordeon function

We shall use the following notation for the Heaviside-step function:

$$u(t - T) = \begin{cases} 0 & \text{for } t < T, \\ 1 & \text{for } t > T, \end{cases} \quad f(T^-) = 0, \quad f(T^+) = 1,$$

so that for  $T < \theta$ , and  $f(t) \in C^0[T, \theta]$ ,

$$\varphi(t) = f(t) [u(t - T) - u(t - \theta)] = \begin{cases} 0 & \text{for } t < T, \\ f(t) & \text{for } T < t < \theta, \\ 0 & \text{for } \theta < t, \end{cases}$$

$$\varphi(T^-) = 0, \quad \varphi(T^+) = f(T^+), \quad \varphi(\theta^-) = f(\theta^-), \quad \varphi(\theta^+) = 0.$$

Under these conditions, with  $a > 0$ , we define the following *accordeon function*:

$$\left. \begin{aligned} \text{Ac}(t, n, a) &= a^{-n-1} \sum_{m=0}^n (-1)^m \binom{n}{m} [u(t - m a) - u(t - (m+1) a)], \\ &= a^{-n-1} \sum_{m=0}^{n+1} (-1)^m u(t - m a) \left[ \binom{n}{m} + \binom{n}{m-1} \right] \\ &= a^{-n-1} \sum_{m=0}^{n+1} (-1)^m \binom{n+1}{m} u(t - m a), \end{aligned} \right\} \quad (6)$$

where use has been made of the classical properties of the binomial coefficients and  $\binom{n}{-1} = 0$ .

If we let  $a \rightarrow 0$  in (6) we obtain a generalized function in the sense of MIKUSINSKI (cf. [2] and [3]). It is easier in this case not to use the notion of distribution in the sense of SCHWARTZ. We shall call this generalized function a *squeezed accordéon* and shall write

$$\lim_{a \rightarrow 0} \text{Ac}(t, n, a) = \text{Sc}(t, n). \quad (7)$$

For any function  $f(t)$  defined and continuous over a sufficiently large neighbourhood of  $t = 0$  we have, using the classical notation for the inner product

$$\left. \begin{aligned} \langle \text{Sc}(t, n), f(t) \rangle &= \int_{-\infty}^{+\infty} f(t) \text{Sc}(t, n) dt \\ &= \int_{-\infty}^{+\infty} f(t) \left[ \lim_{a \rightarrow 0} a^{-n-1} \sum_{m=0}^{n+1} (-1)^m u(t - m a) \binom{n+1}{m} \right] dt. \end{aligned} \right\} \quad (8)$$

Since the integration is independent of  $a$  and of  $m$  we can change the order of the operations and write, using (6)

$$\langle \text{Sc}(t, n), f(t) \rangle = \lim_{a \rightarrow 0} a^{-n-1} \sum_{m=0}^n (-1)^m \binom{n}{m} \int_{-\infty}^{+\infty} f(t) [u(t - m a) - u(t - (m+1) a)] dt.$$

The integral can be written

$$\begin{aligned} \int_{-\infty}^{+\infty} f(t) [u(t - m a) - u(t - (m+1) a)] dt &= \int_{ma}^{(m+1)a} f(t) dt = a f[(m + \theta_m) a] \\ &= a f[(m + \theta + \beta_m) a] = a [f[(m + \theta) a] + a \beta_m Df[(m + \theta + \eta_m \theta_m) a]], \end{aligned}$$

where  $0 \leq \theta_m \leq 1$ ,  $\theta$  is a fixed number such that

$$0 \leq \theta < \min(\theta_1, \theta_2, \dots, \theta_n), \quad \theta_m = \theta + \beta_m, \quad \text{thus } 0 \leq \beta_m \leq 1, \quad 0 \leq \eta_m \leq 1.$$

It follows that

$$\langle \text{Sc}(t, n), f(t) \rangle = \lim_{a \rightarrow 0} a^{-n} \sum_{m=0}^n (-1)^m \binom{n}{m} \{f[(m + \theta) a] + a \beta_m Df[(m + \theta + \eta_m \beta_m) a]\},$$

where the second term in the sum tends to zero with  $a$  so that

$$\langle \text{Sc}(t, n), f(t) \rangle = \lim_{a \rightarrow 0} a^{-n} \sum_{m=0}^n (-1)^m \binom{n}{m} f[(m + \theta) a],$$

which according to (5) gives

$$\langle \text{Sc}(t, n), f(t) \rangle = (-1)^n D^n f(0). \quad (9)$$

(9) shows that  $\text{Sc}(t, n)$  is identical to the  $n$ -th derivative of the Dirac Delta, i.e.  $\text{Sc}(t, n) = \delta^{(n)}(t)$ , as it is usually considered (cf. [4]). In particular for  $n = 0$ ,

$$\langle \text{Sc}(t, 0), f(t) \rangle = f(0) = \langle \delta(t), f(t) \rangle, \quad (10)$$

i.e.  $\text{Sc}(t, 0) = \delta(t)$ , the Dirac Delta.

#### 4. Laplace Transforms

We clearly have

$$\mathcal{L}[\text{Sc}(t, n)] = \int_{0^+}^{+\infty} e^{-st} \text{Sc}(t, n) dt, \quad (11)$$

where the integration starts on the positive side of zero. Thus

$$\mathcal{L}[\text{Sc}(t, n)] = \int_{0^+}^{+\infty} e^{-st} \left[ \lim_{a \rightarrow 0} a^{-n-1} \sum_{m=0}^{n+1} (-1)^m \binom{n+1}{m} u(t - m a) \right] dt.$$

Since the integration is independent of  $a$  and  $m$  we may change the order of operations, i.e.

$$\begin{aligned} \mathcal{L}[\text{Sc}(t, n)] &= \lim_{a \rightarrow 0} a^{-n-1} \sum_{m=0}^{n+1} \binom{n+1}{m} (-1)^m \int_{0^+}^{+\infty} e^{-st} u(t - m a) dt \\ &= \lim_{a \rightarrow 0} a^{-n-1} \sum_{m=0}^{n+1} \binom{n+1}{m} (-1)^m [u(t - m a)] \\ &= \lim_{a \rightarrow 0} a^{-n-1} \sum_{m=0}^{n+1} \binom{n+1}{m} (-1)^m e^{-mas}/s \\ &= \lim_{a \rightarrow 0} (1 - e^{-as})^{n+1}/s a^{n+1} = s^n. \end{aligned}$$

It follows that

$$\mathcal{L}[\text{Sc}(t, n)] = s^n,$$

or,

$$\text{Sc}(t, n) = \delta^{(n)}(t) = \mathcal{L}^{-1}s^n.$$

Thus the  $\text{Sc}(t, n)$  function, i.e. the  $n$ -th derivative of the Dirac Delta is the inverse Laplace Transform of  $s^n$ . This result is considered classical and can be found for example in [4].  
 S. TAUBER, Portland State University USA

## REFERENCES

- [1] VAN DER POL and BREMMER, *Operational Calculus* (Cambridge Univ. Press, Cambridge 1964).
- [2] J. MIKUSINSKI, *Operational Calculus* (Pergamon Press, N.Y. 1959).
- [3] A.H. ZEMANIAN, *Distribution Theory and Transform Analysis* (McGraw-Hill, N.Y. 1965).
- [4] W. KAPLAN, *Operational Methods for Linear Systems* (Addison Wesley, Reading Mass. 1962).

## Aufgaben

**Aufgabe 577.** K. RADZISZEWSKI (Ann. Univ. Marie Curie-Sklodowska A 10, 57–59, 1956) hat bewiesen: Es sei  $P$  der Flächeninhalt des Rechtecks, das einem gegebenen Oval umschrieben ist und das eine Seite in der Richtung  $\theta$  hat. Der Flächeninhalt des Ovals sei  $S$ . Dann ist

$$\frac{4}{\pi} S \leq \bar{P} = \frac{1}{2\pi} \int_0^{2\pi} P d\theta$$

mit Gleichheit nur für den Kreis. Man beweise: Es sei  $S^*$  der Flächeninhalt der Fusspunkt-kurve des Ovals für einen beliebigen inneren Punkt. Dann ist

$$\bar{P} \leq \frac{4}{\pi} S^*$$

mit Gleichheit nur, wenn das Oval durch eine Rotation von  $90^\circ$  in sich übergeführt werden kann.  $S^*$  hat ein einziges Minimum, wenn der Aufpunkt im Inneren variiert. Für glatte Ovale wird das Minimum im Krümmungsschwerpunkt angenommen.

H. GUGGENHEIMER, Polytechnic Institute of Brooklyn, USA

*Lösung des Aufgabenstellers:* Das Oval habe die Stützfunktion  $h(\theta)$ , gegeben als Funktion des Tangentenwinkels. Dann ist

$$\bar{P} = \frac{2}{\pi} \int_0^{2\pi} h(\theta) h\left(\theta + \frac{\pi}{2}\right) d\theta.$$

Wenn der Nullpunkt des Koordinatensystems ein innerer Punkt des Ovals ist, so ist die Fusspunkt-Kurve die Kurve deren Polargleichung  $r(\phi) = h(\theta)$  ist,  $\theta = \phi + \pi/2$ . Daher ist

$$S^* = \frac{1}{2} \int_0^{2\pi} h^2(\theta) d\theta.$$

Die gefragte Ungleichung folgt sofort aus der Schwarzschen Ungleichung für das Integral  $P$ . Gleichheit besteht, wenn  $h(\theta) = h(\theta + \pi/2)$  für alle  $\theta$ .

Eine Translation des Aufpunktes resultiert in einer Änderung der Stützfunktion

$$h(\theta) \rightarrow h(\theta) + a \cos \theta + b \sin \theta.$$