

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 27 (1972)  
**Heft:** 1

**Rubrik:** Kleine Mitteilungen

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Kleine Mitteilungen

### Functionals Close to Each Other

Bishop and Phelps [1] proved the fundamental theorem that every real or complex Banach space is subreflexive, that is the functionals (real or complex) attaining their suprema on the unit sphere of the space are dense in the dual space. (For an extension see [2].) The proof is based on a very useful lemma of Phelps [3, 4], stating an intuitively obvious fact. We shall give a completely elementary proof of this lemma.

*Let  $E$  be a real normed space with unit ball  $B$  and dual  $E'$ . Suppose  $f, g \in E'$ ,  $\|f\| = \|g\| = 1$  and  $f^{-1}(0) \cap B \subset g^{-1}[-\varepsilon, \varepsilon]$  ( $\varepsilon < 1/2$ ). Then  $\|f - g\| \leq 2\varepsilon$  or  $\|f + g\| \leq 2\varepsilon$ .*

*Proof.* If  $f = \pm g$  there is nothing to prove. If not, choose  $a, b \in E$  such that


$$f(a) = g(b) = 1 \quad \text{and} \quad f(b) = g(a) = 0.$$

Denote by  $M$  the plane spanned by  $a$  and  $b$ . Put  $N = f^{-1}(0) \cap g^{-1}(0)$ . Evidently  $E$  is the direct sum of  $N$  and  $M$ . Denote by  $D$  the projection of  $B$  into  $M$  parallel to  $N$ .

Choosing  $D$  as the unit ball, define a norm  $\|\cdot\|'$  in  $M$ . Putting  $f_0 = f|_M$  and  $g_0 = g|_M$ , by construction we have  $\|\alpha f_0 + \beta g_0\|' = \|\alpha f + \beta g\|$  ( $\alpha, \beta \in \mathbb{R}$ ), so we can suppose that  $E$  is two-dimensional,  $D = B$ .

$f^{-1}(0) \cap B \subset g^{-1}[-\varepsilon, \varepsilon]$  implies  $\|b\| \geq \varepsilon^{-1}$ , i.e.  $B$  intersects the line determined by  $b$  between  $-\varepsilon b$  and  $\varepsilon b$ , say  $\partial B$  intersects the line at  $-b_0$  and  $b_0$ .

Take two parallel lines through  $b_0$  and  $-b_0$  whose strip contains  $B$ . As  $\|f\| = \|g\| = 1$ , the strip must contain either  $a + b$  or  $a - b$ . By symmetry one can suppose that it contains  $a + b$ . But then  $B$  is contained in the striped region in the Figure.



Comparing this region with the lines  $f - g = \pm 1$ , one sees immediately that  $\|f - g\| \leq \max(\eta, \lambda)$ . As  $\eta \leq 2\varepsilon$  and  $\lambda \leq 2\varepsilon/(1+\varepsilon)$ , we have  $\|f - g\| \leq 2\varepsilon$ .

*Remarks.* 1. It is evident from the Figure that the result is best possible.

2. The figure also shows that if  $f^{-1}(0) \cap B \subset g^{-1}[-\varepsilon, \varepsilon]$  and  $g^{-1}(0) \cap B \subset f^{-1}[-\varepsilon, \varepsilon]$  then  $\|f - g\| \leq 2\varepsilon/(1 + \varepsilon)$  or  $\|f + g\| \leq 2\varepsilon/(1 + \varepsilon)$ .

Béla Bollobás, Trinity College, Cambridge

#### REFERENCES

- [1] E. BISHOP and R. R. PHELPS, *A Proof that Every Banach Space is Subreflexive*, Bull. Am. math. Soc. 67, 97–98 (1961).
- [2] B. BOLLOBÁS, *An Extension to the Theorem of Bishop and Phelps*, Bull. Lond. math. Soc. 2, 181–182 (1970).
- [3] R. R. PHELPS, *Subreflexive Normed Linear Spaces*, Arch. Math. 8, 444–450 (1957).
- [4] R. R. PHELPS, *A Representation Theorem for Bounded Convex Sets*, Proc. Am. math. Soc. 11, 976–983 (1960).

## A Distribution Property of the Sequence of Lucas Numbers

Let  $\{L_n\}$  ( $n = 1, 2, \dots$ ) be the Lucas sequence: 1, 3, 4, 7, 11,  $\dots$ . Then we have the following lemmas:

**Lemma 1.**  $L_n$  divides  $L_m$  if and only if  $m = (2k - 1)n$ ;  $n > 1$ ;  $k = 1, 2, \dots$  [1].

**Lemma 2.** If  $L_m$  is divided by  $L_n$ ,  $m > n$ , then the remainder  $R$  is zero, or  $R$  is a Lucas number, or  $L_n - R$  is a Lucas number [2].

**Lemma 3.** Let  $p$  be a prime number  $\geq 2$  and with the property that it divides some Lucas number. Suppose that  $L_t$  is the smallest Lucas number with  $L_t \equiv 0 \pmod{p}$ , where  $t$  is an integer  $> 1$ . Then there does not exist an  $L_q$  with  $L_{(2k-1)t} < L_q < L_{(2k+1)t}$  such that  $L_q \equiv 0 \pmod{p}$ .

*Proof.* By lemma 1 we have  $L_{(2k-1)t} \equiv 0 \pmod{p}$ . Now suppose that there exists an  $L_q$  with  $L_{(2k-1)t} < L_q < L_{(2k+1)t}$  such that  $L_q \equiv 0 \pmod{p}$ . Let  $L_q$  be divided by  $L_t$ , then, by lemma 2, the remainder  $R = 0$  or  $R$  is a Lucas number or  $L_t - R$  is a Lucas number. But  $R = 0$  is impossible according to lemma 1, for  $R = 0$  implies that  $L_t$  divides  $L_q$ , which in its turn implies that  $q$  must be an odd multiple of  $t$ . If  $R$  would be a Lucas number, then  $R \equiv 0 \pmod{p}$ , for by the division algorithm  $R = -aL_t + L_q$ , where  $a$  is an integer, and  $L_t, L_q \equiv 0 \pmod{p}$ . Now  $R < L_t$  and  $R \equiv 0 \pmod{p}$  yields a contradiction. Similarly, if  $L_t - R$  would be a Lucas number, then by the same argument we see that  $L_t - R \equiv 0 \pmod{p}$  and  $L_t - R < L_t$ , which again yields a contradiction.

**Definition.** The sequence  $\{x_n\}$  ( $n = 1, 2, \dots$ ) of integers is said to be uniformly distributed mod  $m$  where  $m \geq 2$  is an integer, provided that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \cdot A(N, j, m) = \frac{1}{m}, \quad (*)$$

for each  $j = 0, 1, \dots, m - 1$ , where  $A(N, j, m)$  is the number of  $x_n$ ,  $n = 1, 2, \dots, N$ , that are congruent to  $j \pmod{m}$ .

**Theorem 1.** Let  $\{L_n\}$  ( $n = 1, 2, \dots$ ) be the Lucas sequence. Then  $\{L_n\}$  is not uniformly distributed mod  $p$  for any prime  $p \geq 2$ .

*Proof.* Let  $p$  be a prime  $\geq 2$ . We distinguish two cases.

(1) If  $p$  does not divide any Lucas number, then  $A(N, 0, p) = 0$  which violates (\*) for  $m = p$  and  $j = 0$ .

(2) Let  $p$  divide the Lucas number  $L_t$ , where the index  $t$  is the least possible. Then by virtue of lemma 1 and lemma 3  $p$  divides all  $L_{(2k-1)t}$  ( $k = 1, 2, \dots$ ) and no other Lucas numbers. Consequently

$$A(N, 0, p) = \frac{N - r}{2t} \quad \text{or} \quad \frac{N - r}{2t} + 1 \quad (0 \leq r < 2t)$$

and therefore

$$\lim_{N \rightarrow \infty} \frac{1}{N} A(N, 0, p) = \lim_{N \rightarrow \infty} \left( \frac{1}{2t} - \frac{r}{2Nt} \right) = \frac{1}{2t} \quad \text{as } N \rightarrow \infty.$$

But  $2t$  is even and thus  $\neq p$  if  $p > 2$ . Moreover, the Lucas numbers are not uniformly distributed mod 2. This completes the proof.

**Theorem 2.** Let  $\{L_n\}$  ( $n = 1, 2, \dots$ ) be the Lucas sequence. Then  $\{L_n\}$  is not uniformly distributed mod  $m$  for any composite integer  $m \geq 2$ .

*Proof.* If we assume that  $\{L_n\}$  is uniformly distributed mod  $m$  for some composite integer  $m \geq 2$ , then  $\{L_n\}$  would be uniformly distributed mod  $p$ , where  $p$  is a prime factor of  $m$ , according to theorem 5.1 of [3], which says that a sequence of integers which is uniformly distributed mod  $m$ , where  $m$  is composite, is also uniformly distributed with respect to any positive divisor of  $m$ . This contradicts theorem 1.

Consequently, from theorem 1 and theorem 2 we have:

**Theorem 3.** The sequence of Lucas numbers is not uniformly distributed mod  $m$  for any integer  $m \geq 2$ .

L. Kuipers and Jau-Shyong Shiue, Southern Illinois University, USA

#### REFERENCES

- [1] L. CARLITZ, *A Note on Fibonacci Numbers*, Fibonacci Quart. 2, No. 1, 15–28 (1964).
- [2] VERNER E. HOGGATT, JR., *Fibonacci and Lucas Numbers*, Houghton Mifflin Mathematics Enrichment Series 1969, 46.
- [3] I. NIVEN, *Uniform Distribution of Sequences of Integers*, Trans. Am. math. Soc. 98, 52–61 (1961).

#### Note on Arithmetical Progressions with Equal Products of Five Terms

It is not known if there exist two arithmetical progressions each formed by five positive integers and such that the products of their terms are equal (cf. [2]). This problem leads to the equation

$$(a - 2r)(a - r)a(a + r)(a + 2r) = (b - 2R)(b - R)b(b + R)(b + 2R)$$

or

$$a(a^2 - r^2)(a^2 - 4r^2) = b(b^2 - R^2)(b^2 - 4R^2),$$

where we look for solutions in positive integers  $a, b, r, R$  such that  $a - 2r > 0$ ,  $b - 2R > 0$ .

We are giving here all solutions of this equation in integers in the case when  $a = b$ , i.e. when the middle terms of our progressions are equal, and the progressions consist of integers.

*Theorem.* All integer solutions of the equation

$$(a^2 - r^2)(a^2 - 4r^2) = (a^2 - R^2)(a^2 - 4R^2) \quad (1)$$

are of the form :

- a)  $a, r$  arbitrary integers,  $R = \pm r$ ,
- b)  $r = \pm(u^2 + 4uv - v^2)\varrho$ ,  $R = (2u^2 - 2uv - 2v^2)\varrho$ ,  $a = 2(u^2 + v^2)\varrho$ ,
- c)  $r = \pm(2u^2 - 2uv - 2v^2)\varrho$ ,  $R = (u^2 + 4uv - v^2)\varrho$ ,  $a = 2(u^2 + v^2)\varrho$ ,

where  $u, v, \varrho$  are integers.

*Proof.* The equation (1) can be written as

$$4(r^4 - R^4) = 5a^2(r^2 - R^2).$$

On excluding the case a), we get

$$4(r^2 + R^2) = 5a^2.$$

Now  $a = 2A$ , and

$$r^2 + R^2 = 5A^2. \quad (2)$$

All integer solutions of this equation are well-known (cf. [1], p. 48, Ex. 1) :

$$r = \pm p\varrho, R = q\varrho, A = s\varrho \text{ and } r = \pm q\varrho, R = p\varrho, A = s\varrho$$

where  $p = u^2 + 4uv - v^2$ ,  $q = 2u^2 - 2uv - 2v^2$ ,  $s = u^2 + v^2$  and  $u, v, \varrho$  are integers. Putting here  $a = 2A$  we obtain solutions b) and c) and the Theorem is proved.

*Remark.* It may be easily seen that the equation (1) has no solutions giving arithmetical progressions with positive terms. In fact, assume  $R > r > 0$  and  $a - 2R > 0$ . Then  $2A - 2R = a - 2R > 0$ ,  $A > R$  and  $r^2 + R^2 < 2R^2 < 5R^2$ , contrary to (2). Thus if  $R > r > 0$ , then  $a - 2R \leq 0$ .

K. Szymiczek, Katowice, Poland

#### REFERENCES

- [1] L. E. DICKSON, *Introduction to the Theory of Numbers* (Dover 1957).
- [2] J. GABOVIČ, *On Arithmetical Progressions with Equal Products of Their Terms* (Russian), Coll. Math. 15, 45–48 (1966).

## Nichtnegative Matrizen mit konstanter Zeilensumme

Mit  $M_n(R)$  bezeichnen wir die Menge der quadratischen Matrizen der Ordnung  $n$  über dem Körper  $R$  der reellen Zahlen.

Eine Matrix  $A = (a_{kl}) \in M_n(R)$  heisst *nichtnegativ*, wenn jedes ihrer Elemente nichtnegativ ist. Bekannt ist der folgende

**Satz:** Die Matrix  $A$  sei nichtnegativ und

$$0 < \sum_{l=1}^n a_{kl} = r \text{ für } k = 1, 2, \dots, n. \quad (1)$$

So gilt  $a(r)$ :  $A$  hat den Eigenwert  $r$ ,  $b(r)$ : alle Eigenwerte sind dem Betrage nach nicht grösser als  $r$ , und  $c(r)$ : Eigenwerten mit dem Betrage  $r$  entsprechen lineare Elementarteiler.

Diesen Satz erhält man z. B. aus einer entsprechenden Aussage über stochastische Matrizen (s. etwa GANTMACHER [1], S. 72ff.). Der Beweis kann aber besonders einfach geführt werden, wenn man den folgenden Satz aus der Theorie der Differenzengleichungen als bekannt voraussetzt (s. HAHN [2], S. 52):

Die Nulllösung des Systems linearer Differenzengleichungen mit konstanten Koeffizienten

$$x(t+1) = B x(t) \quad (2)$$

ist genau dann stabil, wenn für  $B$  die Aussagen  $b(1)$  und  $c(1)$  gelten.

Dabei heisst die Nulllösung des Systems (2) *stabil*, wenn sich zu jedem positiven  $\varepsilon$  ein positives  $\delta$  finden lässt, so dass  $|x(0)| < \delta$

$$|x(t)| < \varepsilon, \quad t = 1, 2, \dots,$$

nach sich zieht.  $|\cdot|$  bedeutet hier eine beliebige Vektornorm. Wir zeigen, dass die Nulllösung von (2) stabil ist, wenn  $A = r B$  nichtnegativ ist und (1) erfüllt.

Setzen wir  $|x| = \max_i |x_i|$  und bezeichnet  $Z(A)$  die zu dieser Vektornorm passende Zeilennorm von  $A$  (s. etwa ZURMÜHL [3], S. 202ff.), so ist

$$|x(t+1)| \leq Z(B) |x(t)| = \frac{1}{r} r |x(t)| = |x(t)|,$$

und daher

$$|x(t)| \leq |x(0)|, \quad t = 1, 2, \dots,$$

wie behauptet.

Also gelten die Aussagen  $b(1)$  und  $c(1)$  für  $B$ , und daher  $b(r)$  und  $c(r)$  für  $A$ .  $a(r)$  folgt daraus, dass  $A$  den Eigenvektor  $\text{col}(1, 1, \dots, 1)$  hat.

H. Stettner, TH Graz

### LITERATUR

- [1] F. R. GANTMACHER, *Matrizenrechnung II* (VEB Deutscher Verlag der Wissenschaften, Berlin 1959).
- [2] W. HAHN, *Stability of Motion* (Springer, Berlin 1968).
- [3] R. ZURMÜHL, *Matrizen* (Springer, Berlin 1964).