

Zeitschrift:	Elemente der Mathematik
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	27 (1972)
Heft:	2
Artikel:	A new method of evaluating the sums of [formula] and related series
Autor:	Stark, Eberhard
DOI:	https://doi.org/10.5169/seals-28628

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

REFERENCES

- [1] B. L. VAN DER WAERDEN, Elem. Math. 25, 73 (1970). See also: W. LÜSSY and E. TROST, Elem. Math. 25, 82 (1970); H. IRMINGER, Elem. Math. 25, 135 (1970).
- [2] J. WASER and V. SCHOMAKER, J. Amer. Chem. Soc., 67, 2014 (1945).
- [3] L. PAULING, *The Nature of the Chemical Bond*, 2nd Ed. (Cornell University Press, Ithaca, N.Y., 1940).
- [4] J. WASER, Proposition 10, Ph. D. Dissertation, California Institute of Technology, Pasadena, 1944. Oral Examination, September 5, 1944.
- [5] J. H. BURNS and J. WASER, J. Amer. Chem. Soc. 79, 859 (1956).
- [6] J. DONOHUE, Acta Crystallogr. 15, 708 (1962).
- [7] G. A. SIM, J. Chem. Soc. London, 5974 (1965).
- [8] E. RUCH, personal communication to J. D. D., February, 1970.
- [9] L. OOSTERHOFF, personal communication to J. D. D., May, 1971.
- [10] See, for example, A. C. ATKEN, *Determinants and Matrices*, 7th Edition, p. 123 (Oliver & Boyd, Edinburgh and London 1951).
- [11] See, for example, M. LAGALLY, *Vorlesungen über Vektorrechnung* (Akademische Verlagsgesellschaft, Leipzig 1944), p. 30.

A New Method of Evaluating the Sums of

$$\sum_{k=1}^{\infty} (-1)^{k+1} k^{-2p}, \quad p = 1, 2, 3, \dots \text{ and Related Series}$$

The decisive tool in our attempt to evaluate the sum $\sum_{k=1}^{\infty} (-1)^{k+1} k^{-2p}$ for fixed $p \in \mathbb{N} = \{1, 2, 3, \dots\}$ is the kernel of Dirichlet in both of its representations (for all real x)

$$D_n(x) = \frac{\sin(2n+1)\frac{x}{2}}{2 \sin \frac{x}{2}} = \frac{1}{2} + \sum_{k=1}^n \cos kx \quad (n \in \mathbb{N}). \quad (1)$$

First of all, let us consider

$$C_p(k) = \int_0^\pi t^{2p} \cos kt dt, \quad k \in \mathbb{N};$$

a twofold integration by parts gives the recursive formula

$$C_p(k) = \frac{2p}{k^2} \left\{ (-1)^k \pi^{2p-1} - (2p-1) C_{p-1}(k) \right\} \quad (p \in \mathbb{N})$$

$$C_0(k) = 0$$

and hence, as is immediately verified¹⁾,

$$C_p(k) = (-1)^k \pi(2p)! \sum_{j=1}^p (-1)^{j+1} \frac{\pi^{2(p-j)}}{(2(p-j)+1)!} \frac{1}{k^{2j}} \quad (p \in \mathbb{N}). \quad (2)$$

¹⁾ Formula 3.529.1 in [4] is obviously incorrect.

Now, concerning the even moments of (1), i.e.,

$$M(2p; n) = \int_0^\pi t^{2p} D_n(t) dt \quad (p \in \mathbb{N}),$$

we have, on the one hand, by the polynomial representation

$$M(2p; n) = \frac{\pi^{2p+1}}{2(2p+1)} + \sum_{k=1}^n C_p(k); \quad (3)$$

and on the other,

$$M(2p; n) = 2^{2p} \int_0^{\pi/2} \frac{y^{2p}}{\sin y} \sin(2n+1)y dy.$$

The second mean-value theorem (cf. e.g. [6, p. 163]) yields

$$\int_0^{\pi/2} \frac{y^{2p}}{\sin y} \sin(2n+1)y dy = \left(\frac{\pi}{2}\right)^{2p} \frac{\cos(2n+1)\xi}{2n+1} \quad (0 \leq \xi \leq \frac{\pi}{2}, n \in \mathbb{N})$$

since $\sin(2n+1)y$ is continuous, whereas $y^{2p}/\sin y$ is non-negative and increasing for $y \in [0, \pi/2]$. Thus, uniformly for all $p \in \mathbb{N}$,

$$M(2p; n) = O(n^{-1}) \quad (n \rightarrow \infty).$$

This gives, using (3) and (2),

$$\lim_{n \rightarrow \infty} \left\{ \frac{\pi^{2p}}{2(2p+1)} - \sum_{k=1}^n (-1)^{k+1} (2p)! \sum_{j=1}^p (-1)^{j+1} \frac{\pi^{2(p-j)}}{(2(p-j)+1)!} \frac{1}{k^{2j}} \right\} = 0$$

or

$$\sum_{k=1}^{\infty} (-1)^{k+1} \sum_{j=1}^p (-1)^{j+1} \frac{\pi^{2(p-j)}}{(2(p-j)+1)!} \frac{1}{k^{2j}} = \frac{\pi^{2p}}{2(2p+1)!} \quad (p \in \mathbb{N}).$$

After extracting the term for $j = p$, the desired formula reads

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2p}} &= (-1)^{p+1} \frac{\pi^{2p}}{2(2p+1)!} \\ &\quad - \sum_{k=1}^{\infty} (-1)^{k+1} \sum_{j=1}^{p-1} (-1)^{p+j} \frac{\pi^{2(p-j)}}{(2(p-j)+1)!} \frac{1}{k^{2j}}. \end{aligned} \quad (4)$$

From (4), by recursion for $p = 1, 2, 3, \dots$ the sums are easily calculated: in case $p = 1$ it follows directly that

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2} = \frac{\pi^2}{12}; \quad (5)$$

for $p = 2$, together with (5),

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^4} = -\frac{\pi^4}{240} + \frac{\pi^2}{6} \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2} = -\frac{\pi^4}{240} + \frac{\pi^4}{72},$$

so that

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^4} = \frac{7}{720} \pi^4; \quad (6)$$

continuing in this way, for $p = 3$ now using both (5) and (6), one obtains

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^6} = \frac{31}{30240} \pi^6, \dots$$

The standard proof (see e.g. [2, p. 245ff], [1, p. 244]) by means of the theory of Bernoulli polynomials leads to

$$\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2p}} = \frac{(-1)^{p+1} \pi^{2p}}{2(2p)!} (2^{2p} - 2) B_{2p} \quad (p \in N) \quad (7)$$

where B_{2p} are the Bernoulli numbers. These, in turn, may be deduced from the symbolic equation (i.e. expand (8) and set formally $B^n = B_n$)

$$(B + 1)^n - B^n = 0, \quad B_0 = B^0 = 1, \quad n = 1, 2, 3, \dots, \quad (8)$$

cf. [1, p. 233], [2, p. 185].

Of course, neither (4) nor (7) are actually closed expressions for the sum since the Bernoulli numbers are obtainable only via the recurrence formula (8). But (4) has the advantage of avoiding the use of B_{2p} ; moreover a comparison of (7) and (4) gives another way of evaluating these numbers.

In view of the obvious relations, cf. [2, p. 246],

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{1}{k^{2p}} &= \frac{2^{2p}}{2^{2p} - 2} \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2p}}, \\ \sum_{k=1}^{\infty} \frac{1}{(2k-1)^{2p}} &= \frac{2^{2p}-1}{2^{2p}-2} \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2p}} \end{aligned} \quad (p \in N)$$

the sums of these series may also be derived immediately from (4).

In the particular case $p = 1$, there is a proof in [5] using another kernel, namely that of Fejér, whereas the method employed in [3] is implicitly based upon the kernel of de La Vallée Poussin. This reveals that these kernels originating from approximation theory, also play an interesting rôle in a quite different branch of analysis.

Eberhard L. Stark, Technological University of Aachen, Germany²⁾

REFERENCES

- [1] C. JORDAN, *Calculus of Finite Differences*, 2nd ed. (New York 1950).
- [2] K. KNOPP, *Theorie und Anwendungen der unendlichen Reihen*, 4. Aufl. (Berlin 1947).
- [3] Y. MATSUOKA, *An Elementary Proof of the Formula* $\sum_{k=1}^{\infty} 1/k^2 = \pi^2/6$, Amer. Math. Monthly 68, 485–487 (1961).
- [4] I. M. RYSHIK and I. S. GRADSTEIN, *Summen-, Produkt- und Integraltafeln*, 2nd ed. (Berlin 1963).
- [5] E. L. STARK, *Another Proof of the Formula* $\sum_{k=1}^{\infty} 1/k^2 = \pi^2/6$, Amer. Math. Monthly 76, 552–553 (1969).
- [6] D. V. WIDDER, *Advanced Calculus*, 2nd ed. (Englewood Cliffs 1961).

²⁾ The author is indebted to Professor P. L. BUTZER for many helpful remarks. A support of the *Deutsche Forschungsgemeinschaft* is greatly acknowledged.