**Zeitschrift:** Elemente der Mathematik
**Herausgeber:** Schweizerische Mathematische Gesellschaft
**Band:** 29 (1974)
**Heft:** 1

**Artikel:** Rationals not expressible as a sum of three unit fractions
**Autor:** Webb, William A.
**DOI:** https://doi.org/10.5169/seals-29889

**Nutzungsbedingungen**
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

**Conditions d'utilisation**
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

**Terms of use**
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 18.02.2026

**ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch**

ELEMENTE DER MATHEMATIK
Revue de mathématiques élémentaires - Rivista di matematica elementare
*Zeitschrift zur Pflege der Mathematik*
*und zur Förderung des mathematisch-physikalischen Unterrichts*
Publiziert mit Unterstützung des Schweizerischen Nationalfonds
zur Förderung der wissenschaftlichen Forschung

El. Math. Band 29 Heft 1 Seiten 1–24 10. Januar 1974

# Rationals Not Expressible as a Sum of Three Unit Fractions

## I. Introduction

Let $k$ be a positive integer and define

$$\delta_k = \left\{ n : \frac{k}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z};\ x, y, z \text{ positive integers} \right\} \tag{1.1}$$

and

$$\lambda_k = \sup_{n \notin \delta_k} n\,. \tag{1.2}$$

It has been conjectured by Schinzel [5] that $\lambda_k < \infty$ for all $k$; some specific cases of this conjecture are due to Erdös [3] – $k = 4$; Sierpiński [5] – $k = 5$; and Aigner [1] – $k = 6, 7$.

Since the set of fractions expressible as a sum of three unit fractions is nowhere dense ([1, 7]) we must have $\limsup \frac{k}{\lambda_k} = 0$, that is $\lambda_k$ must approach $\infty$ faster than $k$ does. However, for small $k$, $\lambda_k$ also seems small. For example if the conjectures are correct $\lambda_4 = 1$, $\lambda_5 = 1$, $\lambda_6 = 1$ and $\lambda_7 = 2$, where the magnitude of $7/2$ alone implies that $2 \notin \delta_7$. The first case where $\lambda_k > k$ is $k = 8$. Aigner [1] noted that $17 \notin \delta_8$ so $\lambda_8 \geq 17$. We will see later that actually $\lambda_8 \geq 241$.

In this paper we will look for examples of $n \notin \delta_k$ for $k \geq 6$.

## II. General Methods

One of the principal tools we will use in our search for $n \notin \delta_k$ is the following lemma.

*Lemma* 2.1 $a/b = 1/x + 1/y$ if and only if there exist positive divisors $d_1$ and $d_2$ of $b$ such that $a \mid d_1 + d_2$.

This is a modified form of a result of Aigner [1] Satz 6. For a complete proof of some generalizations of Lemma 2.1 see [7].

If $k/n = 1/x + 1/y + 1/z$, $0 < x \leq y \leq z$, then $x \leq \frac{3n}{k}$. Hence, in checking if $n \in \delta_k$ it suffices to apply Lemma 2.1 to $k/n - 1/x$ for each $x$ such that $\frac{n}{k} \leq x \leq \frac{3n}{k}$.

Also, since $n \in \delta_k$ implies $n\,m \in \delta_k$ for all $m \geq 1$, we can limit ourselves to the case where $n$ is a prime.

The following result shows that we may restrict the values of $x$ we must check even further.

*Theorem* 2.2 Let $k \geq 4$, $p$ a prime such that $(k, p) = 1$, and let $x \leq y \leq z$ be positive integers such that

$$\frac{k}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}; \tag{2.1}$$

then $x \leq (2p+2)/k$. Moreover, if $x > 2p/k$ then either $k \mid 2p+1$ and $x = (2p+1)/k$ or $k \mid p+1$ and $x = (2p+2)/k$.

Proof. Now by Lemma 2.1

$$\frac{k}{p} - \frac{1}{x} = \frac{kx-p}{px} = \frac{a}{b} = \frac{1}{y} + \frac{1}{z}; \tag{2.2}$$

if and only if there exist $d_1, d_2 \mid px$ such that $d_1 + d_2 = s(kx - p)$ for some integer $s$; in which case

$$\frac{a}{b} = \frac{1}{s\left(\frac{b}{d_1}\right)} + \frac{1}{s\left(\frac{b}{d_2}\right)}. \tag{2.3}$$

Hence, $x \leq s\,b/d_1 \leq s\,b/d_2$ since we may assume $d_1 \geq d_2$.

Now, assume that $x > 2p/k$. This implies that $kx - p > p$. We know that $x \leq 3p/k \leq 3p/4$ so $p \nmid x$. Hence $(p, kx) = 1$ and $(p, kx - p) = 1$.

If $d \leq e$ are any two positive divisors of $x$, then $kx - p \nmid d + e$. This may be seen as follows:

Assume $kx - p \mid d + e$. Then $d + e \leq 2x \leq \frac{3}{2}p \leq \frac{3}{2}(kx - p)$, so $kx - p = d + e$. Also, if $e < x$, $d + e \leq x < kx - p$; so $e = x$. Hence, $d = (k-1)x - p$ which implies $d \mid p$, and so $d = 1$ since $(p, x) = 1$. Therefore

$$x = \frac{p+1}{k-1} \leq \frac{2p}{k}, \text{ a contradiction}.$$

Since we have that $d_1, d_2 \mid px$ and $kx - p \mid d_1 + d_2$, by the above, we cannot have $d_1, d_2 \mid x$. Similarly we cannot have $d_1 = p\,d_3$ and $d_2 = p\,d_4$ where $d_3, d_4 \mid x$, since this would mean that $kx - p \mid p\,(d_3 + d_4)$ which implies $kx - p \mid d_3 + d_4$ again contradicting what we have shown above. Therefore, and since $d_1 \geqslant d_2$, we must have $d_1 = p\,d$ where $d \mid x$ and $d_2 \mid x$.

Now $p\,d + d_2 = s\,(kx - p) > s\,p$ and $d_2 \leq x < p$ which implies $d \geq s$. Hence,

$$\frac{s\,b}{d_1} = \frac{s\,p\,x}{d_1} = \frac{s\,p\,x}{p\,d} = \frac{s\,x}{d} \leq x; \tag{2.4}$$

but $x \leq s\,b/d_1$ and so $x = s\,x/d = s\,b/d_1 = y$. Thus from (2.2), (2.3) and (2.4) we see that

$$\frac{k}{p} - \frac{2}{x} = \frac{kx - 2p}{px} = \frac{1}{z}, \tag{2.5}$$

and so $kx - 2p \mid px$.

Clearly $(p, kx - 2p) = 1$, so $kx - 2p \mid x$ which implies $kx - 2p \mid 2p$ which implies $kx - 2p = 1$ or 2. If $kx - 2p = 1$ then $k \mid 2p + 1$ and $x = (2p + 1)/k$. If $kx - 2p = 2$, then (2.5) holds if and only if $x$ is even, which implies $x = (2p + 2)/k$ and $k \mid p + 1$.

This completes the proof of our theorem.

This theorem says that in looking for a solution of (2.1) we need only check $x \leqslant (2p + 2)/k$. The theorem is best possible as the cases $k \mid 2p + 1$ and $k \mid p + 1$ show; since if $k \mid 2p + 1$ then

$$\frac{k}{p} = \frac{1}{(2p+1)/k} + \frac{1}{(2p+1)/k} + \frac{1}{p\,(2p+1)/k},$$

and if $k \mid p + 1$ then

$$\frac{k}{p} = \frac{1}{2\,(p+1)/k} + \frac{1}{2\,(p+1)/k} + \frac{1}{p\,(p+1)/k}.$$

Although these examples show that $x$ may be as large as $(2p + 2)/k$ in a solution of (2.1), in all of the examples so far encountered, there have been other solutions of (2.1) with a smaller value of $x$. For example in the case $k \mid p + 1$ mentioned above, we may take $x = (p + 1)/k$ since

$$\frac{k}{p} = \frac{1}{(p+1)/k} + \frac{1}{2p\,(p+1)/k} + \frac{1}{2p\,(p+1)/k}.$$

If $x_0$ denotes the smallest value of $x$ such that (2.1) is solvable, how close to $(2p + 2)/k$ can $x_0$ be? If $k = 12$, $p = 727$ then $x_0 = 108$;

$$\frac{12}{727} = \frac{1}{108} + \frac{1}{138} + \frac{1}{1805868}.$$

Since $108 > (1.78)\,\dfrac{727}{12}$, we see that the bound $2p/k$ cannot be improved too much. Probably even better examples can be found.

## III. Numerical Results

In this section we will give some examples of primes $p$ for which (2.1) has no solution. Using Lemma 2.1 it can be shown that for a fixed $k$, (2.1) is solvable if $p$ is an element of certain residue classes. For small values of $k$ we can restrict our search for counterexamples to a very few residue classes. For example, for $k = 5$ Stewart [6] has shown that if (2.1) is unsolvable then $p \equiv 1 (\mathrm{mod}\ 278460)$. Using this he has shown that (2.1) is solvable for all $p < 1{,}057{,}438{,}801$. Yamamoto [8] has shown that for $k = 4$, (2.1) is solvable for all $p < 10^7$. This extends work done by Aigner [1],

Bernstein [2], Palamà [4]. As $k$ becomes larger this procedure becomes less effective since it eliminates a smaller fraction of the integers from consideration, and it multiplies the number of cases to be considered. For example consider the cases $k = 5$ and $k = 10$. If $k = 5$ we need consider only one residue class modulo 278460. But if $k = 10$, although we can easily eliminate the residue classes 2, 4, 5, 6, 7, 8, 9, 10 (mod 10), we are left with the possibilities $p \equiv 1, 3, 7 \pmod{10}$. It is not hard to show that actually we may restrict ourselves to the residue classes 1 (mod 10), 3 (mod 140), 43 (mod 140) and 7 (mod 60). Furthermore we can replace the residue class 1 (mod 10) by the residue classes 1, 11, 31, 41, 61 (mod 90) which would reduce the number of numbers considered slightly, but we would have five separate cases instead of only one. It is probably possibly to find a procedure somewhat more efficient for $k = 10$ than this; but as we shall see in Theorem 3.2, since there are several primes $p$ for which (2.1) is unsolvable with $k = 10$ and these $p$ fall in three different residue classes modulo 10, it is impossible to do nearly as well as we can in the case $k = 5$.

The procedure used in this paper is to remove as many residue classes as seems practical by the above method, and then use a computer to search for counter-examples in the remaining residue classes. The method used in the computer search is essentially that given near the beginning of section 2.

As an example we take $k = 6$, and assume (2.1) is not solvable. It is easily seen that $p = 6n + 1$. We now eliminate various residue classes for $n$ as follows.

$$\frac{6}{6n+1} - \frac{1}{n+1} = \frac{5}{(6n+1)(n+1)}. \tag{3.1}$$

Now apply Lemma 2.1.

| $n$ | $d_1$ | $d_2$ | |
|---|---|---|---|
| $\equiv 4 \pmod 5$ | | | $5 \mid n+1$ |
| $\equiv 3 \pmod 5$ | $n+1$ | 1 | |
| $\equiv 2 \pmod 5$ | $(6n+1)(n+1)$ | 1 | |
| $\equiv 1 \pmod 5$ | $(6n+1)(n+1)$ | 1 | |
| $\equiv 5 \pmod{10}$ | $(n+1)/2$ | 2 | |

Hence $n \equiv 0 \pmod{10}$ and $p = 60m + 1$. Then

$$\frac{6}{60m+1} - \frac{1}{10m+2} = \frac{11}{2(60m+1)(5m+1)}. \tag{3.2}$$

Apply Lemma 2.1 again.

| $m$ | $d_1$ | $d_2$ | |
|---|---|---|---|
| $\equiv 10 \pmod{11}$ | $2(60m+1)(5m+1)$ | 1 | |
| $\equiv 8 \pmod{11}$ | $(60m+1)(5m+1)$ | 2 | |
| $\equiv 7 \pmod{11}$ | $(60m+1)(5m+1)$ | 2 | |
| $\equiv 6 \pmod{11}$ | $5m+1$ | 2 | |
| $\equiv 5 \pmod{11}$ | $2(60m+1)(5m+1)$ | 1 | |
| $\equiv 4 \pmod{11}$ | $5m+1$ | 1 | |
| $\equiv 3 \pmod{11}$ | $2(5m+1)$ | 1 | |
| $\equiv 2 \pmod{11}$ | | | $11 \mid 5m+1$ |

Thus $m \equiv 0$, 1 or 9 (mod 11) and so $p \equiv 1$, 61 or 541 (mod 660).

Using the methods described above we have the following results.

*Theorem* 3.1 The equations

$$\frac{6}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \tag{3.3}$$

and

$$\frac{7}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \tag{3.4}$$

are solvable for all $3 \leq p \leq 100{,}000$.

*Theorem* 3.2. The following table lists values of $p > k/3$ for which

$$\frac{k}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \tag{3.5}$$

is not solvable.

| $k$ | $p$ |
|---|---|
| 8 | 11, 17, 131, 241 |
| 9 | 5, 11, 19 |
| 10 | 7, 11, 43, 61, 67, 181 |
| 11 | 37 |
| 12 | 5, 13, 29, 37, 73, 97, 193, 433, 577, 1129, 1657, 1873, 2521, 2593, 3433, 10369, 12049, 12241 |
| 13 | 5, 7, 53, 61, 67, 79, 211, 281 |
| 14 | 5, 17, 19, 29, 59, 257, 353 |
| 15 | 17, 19, 23, 31, 47, 53, 61, 79, 113, 137, 151, 197, 233, 271, 541, 1103, 1171, 1367, 4201 |
| 16 | 7, 11, 13, 17, 23, 37, 73, 97, 113, 131, 167, 193, 241, 257, 421, 577, 593, 641, 769, 1201, 1489, 2113, 2521, 2689, 3169, 3361, 4801, 4993 |
| 17 | 7, 13, 19, 23, 41, 53, 71, 73, 157, 281, 421, 1123, 2081 |
| 18 | 7, 11, 13, 19, 23, 29, 31, 37, 41, 47, 59, 61, 73, 109, 113, 131, 137, 149, 181, 193, 223, 239, 281, 379, 389, 397, 443, 457, 541, 599, 613, 661, 761, 811, 821, 911, 1009, 1297, 1381, 2269, 2819 |

For $8 \leq k \leq 11$ the table is complete for $p \leq 25000$, for $13 \leq k \leq 18$ the table is complete for $p \leq 5000$ and for $k = 12$ it is complete for $p \leq 100000$. The case $k = 12$ was carried further than the others since it is the first case for which (3.5) is unsolvable for some relatively large primes. It is quite likely that more counterexamples for $13 \leq k \leq 18$ can be found, if we check for $p > 5000$. The total computer time used in verifying theorems 3.1 and 3.2 was approximately 20 minutes using an IBM 360/67.

These examples show that $\lambda_k$ gets relatively large for even some small values of $k$ ($\lambda_{12} \geq 12241$), something that was not evident for $k \leq 7$ studied previously. However, in the case $k = 12$ which was carried out further than the others, there were no more counterexamples for $12241 < p \leq 100000$. Thus these examples

appear to give some evidence both for and against the general conjecture that $\lambda_k < \infty$ for all $k$.

The author wishes to thank John Steinig for several valuable suggestions.

William A. Webb, Washington State University, Pullman, Washington

REFERENCES

[1] A. Aigner, *Brüche als Summe von Stammbrüchen*, J. reine angew. Math. *214/215*, 174–179, (1964).

[2] L. Bernstein, *Zur Lösung der diophantischen Gleichung* $m/n = 1/x + 1/y + 1/z$, *insbesondere im Falle* $m = 4$, J. reine angew. Math. *211*, 1–10 (1962).

[3] P. Erdös, *On a diophantine equation*, Mat. Lapok, *1*, 192–210 (1950).

[4] G. Palamà, *Su di una congettura di Sierpiński relativa alla possibilità in numeri naturali della* $5/n = 1/x_1 + 1/x_2 + 1/x_3$, Boll. Un. Mat. Ital., *13*, 65–72 (1958).

[5] W. Sierpiński, *Sur les décompositions de nombres rationnels en fractions primaires*, Mathesis *65*, 16–32 (1956).

[6] B. Stewart, *Theory of Numbers*, 2nd ed., 198–207, (New York, 1964).

[7] B. Stewart and W. Webb, *Sums of Fractions with Bounded Numerators*, Can. J. Math. *18*, 999–1003 (1966).

[8] K. Yamamoto, *On the Diophantine Equation* $4/n = 1/x + 1/y + 1/z$., Mem. Fac. Sci. Kyushu U., [A] *19*, 37–47 (1965).

# Eine bemerkenswerte Abbildung der Punkte des Raumes auf die Kreise einer Ebene

## Einleitung

Die im folgenden untersuchte Abbildung hat gewisse Ähnlichkeiten mit der *Zyklographie* [1, 2]. Bei ihr werden die Punkte des dreidimensionalen Raumes $R^3$ auf die Kreise einer im $R^3$ enthaltenen, waagrecht gedachten Ebene abgebildet, wobei statt der in der Zyklographie für den Abbildungsvorgang verwendeten Drehkegel mit lotrechter Achse Drehparaboloide mit lotrechter Achse und festem Parameter ($= 1/2$) treten.

Ein wesentlicher Unterschied gegenüber der Zyklographie liegt darin, dass die Abbildung ohne Orientierung der Kreise auskommt und trotzdem umkehrbar eindeutig ist. Durch zyklographische Deutung der Bildkreise wird im Raum eine zweieindeutige Punkttransformation induziert, die einen klaren Einblick in das Wesen der neuen Abbildung gewährt.

Einer Geraden des $R^3$ entspricht die Menge der eine Parabel der Bildebene doppelt berührenden Kreise, während die Scharen der einen Mittelpunktskegelschnitt doppelt berührenden Kreise von Parabeln mit lotrechter Achse herrühren. Diese Tatsachen lassen die konstruktive Lösung verschiedener damit zusammenhängender Aufgaben zu.