

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 29 (1974)  
**Heft:** 3

**Artikel:** On 1-factorability and edge-colorability of cartesian products of graphs  
**Autor:** Himmelwright, P.E. / Williamson, J.E.  
**DOI:** <https://doi.org/10.5169/seals-29897>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 25.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Hence  $G_0 \neq G$ ,  $G_{\{0,\infty\}} \neq G_0$ . It is also clear that  $(G_{\{0,\infty\}}, k - \{0\}, *)$  is transitive, for if  $x \neq 0$  and  $y \neq 0$ , then

$$\begin{bmatrix} y & 0 \\ 0 & x \end{bmatrix} * x = y.$$

Hence by Theorem 3,  $(G, X, *)$  is 3-fold transitive. We note that  $(G, X, *)$  is not 4-fold transitive, for then  $(G_{\{0,\infty\}}, k - \{0\}, *)$  would be 2-fold transitive.

David P. Sumner, University of South Carolina, USA

### Acknowledgement

I would like to thank D. J. Foulis and L. N. Mann for conducting the seminar on transformation groups during which this note was first written.

### REFERENCE

- [1] H. WIELANDT, *Finite Permutation Groups*, trans. by R. Bercov, Academic Press, New York, 1964.

## On 1-Factorability and Edge-Colorability of Cartesian Products of Graphs

There is no characterization of 1-factorable graphs. Thus, it is natural that many of the results on this topic have been the determination of classes of 1-factorable graphs. The object of this paper is to present a sufficient condition for the 1-factorability of the cartesian product of two graphs. We begin with some notation and definitions.

The vertex set of a graph  $G$  will be denoted by  $V(G)$  and its edge set by  $E(G)$ . In this paper we consider only finite, undirected graphs without loops or multiple edges. Let  $G$  and  $H$  be two nonempty graphs for which  $V(G) = V(H)$  and  $E(G) \cap E(H) = \emptyset$ ; then the graph  $G'$  is the *sum* of  $G$  and  $H$ , written  $G' = G + H$ , if  $V(G') = V(G)$  and  $E(G') = E(G) \cup E(H)$ . A *1-factor* of a graph  $G$  is a spanning 1-regular subgraph of  $G$ . A graph is *1-factorable* if it can be expressed as a sum of edge-disjoint 1-factors. The *cartesian product* (or *product*) of the graph  $G$  with the graph  $H$ , denoted by  $G \times H$ , is defined by:  $V(G \times H) = V(G) \times V(H)$ ;  $E(G \times H) = \{[(u_1, v_1), (u_2, v_2)] \mid u_1 = u_2 \text{ and } v_1 v_2 \in E(H), \text{ or } v_1 = v_2 \text{ and } u_1 u_2 \in E(G)\}$ .

An assignment of  $n$  colors to the edges of a nonempty graph  $G$  so that adjacent edges are colored differently is an  *$n$ -edge-coloring* of  $G$ . The minimum  $n$  for which a graph  $G$  is  $n$ -edge-colorable is its *edge-chromatic number*  $\chi_1(G)$ . By a theorem of Vizing [2], the edge-chromatic number  $\chi_1(G)$  of a graph  $G$  is bounded by:  $\Delta(G) \leq \chi_1(G) \leq \Delta(G) + 1$ , where  $\Delta(G)$  is the maximum degree of  $G$ . If  $G$  is regular, then  $G$  is 1-factorable if and only if  $\chi_1(G) = \Delta(G)$ . Hence any theorem concerning the 1-factorability of regular graphs has as an immediate corollary a result concerning edge-colorability, which is useful since there is also no characterization of those graphs which are  $\Delta(G)$ -edge-colorable. For other notations and definitions, we follow [1].

If  $K_2$  denotes the complete graph on two vertices, then  $K_2 \times H$ , where  $H$  is any regular graph, is shown to be 1-factorable in the following lemma.

**Lemma:** If  $H$  is a regular graph, then  $K_2 \times H$  is 1-factorable.

*Proof.* If  $H$  is 1-factorable, then the result follows immediately. Hence we consider the case that  $H$  is not 1-factorable. If  $H$  is an  $r$ -regular graph, then by a previous remark,  $\chi_1(H) = r+1$ . Let an  $(r+1)$ -edge-coloring of  $H$  be given and let  $C_1, C_2, \dots, C_{r+1}$  be the edge-color classes of  $E(H)$ . Now  $K_2 \times H$  contains two disjoint copies of  $H$ . Let the  $(r+1)$ -edge-coloring of  $H$  be applied to these disjoint copies, and assign to each edge  $[(u_1, v), (u_2, v)]$  of  $K_2 \times H$  the only color among the  $r+1$  colors which was assigned to no edge of  $H$  incident with  $v$ . Hence  $K_2 \times H$  may be  $(r+1)$ -edge-colored. But  $K_2 \times H$  is  $(r+1)$ -regular. Hence  $\chi_1(K_2 \times H) = r+1$ , and  $K_2 \times H$  is 1-factorable.

We now state and prove the main result.

**Theorem:** If  $G$  is a 1-factorable graph and  $H$  is a regular graph, then  $G \times H$  is a 1-factorable graph.

*Proof:* Let  $G$  be a 1-factorable,  $r$ -regular graph of order  $p_1$  with 1-factors  $G_1, G_2, \dots, G_r$ , and let  $H$  be an  $s$ -regular graph of order  $p_2$ . Then consider the subgraphs  $G_1 \times H, G_2 \times \bar{K}_{p_2}, \dots, G_r \times \bar{K}_{p_2}$  of  $G \times H$ , where  $\bar{K}_{p_2}$  denotes the graph consisting of  $p_2$  isolated vertices. Note that these subgraphs are mutually edge-disjoint subgraphs spanning  $G \times H$ , and  $G \times H = G_1 \times H + \sum_{i=2}^r G_i \times \bar{K}_{p_2}$ . Moreover, the subgraphs  $G_2 \times \bar{K}_{p_2}, \dots, G_r \times \bar{K}_{p_2}$  are 1-regular and thus are 1-factors of  $G \times H$ . Hence if  $G_1 \times H$  is 1-factorable,  $G \times H$  is 1-factorable. Now  $G_1 \times H$  is a spanning  $(s+1)$ -regular subgraph of  $G \times H$  consisting of  $p_1/2$  components each of which is isomorphic to  $K_2 \times H$ . By the Lemma,  $K_2 \times H$  is 1-factorable and of regularity  $s+1$ . Let the 1-factors of  $K_2 \times H$  be  $F_1, F_2, \dots, F_{s+1}$  in a 1-factorization of  $K_2 \times H$ . Select in every component of  $G_1 \times H$ , the same 1-factor  $F_k$ , where  $1 \leq k \leq s+1$ , and designate the resultant subgraph of  $G_1 \times H$  by  $F'_k$ . Then by the choice of  $F'_k$  it follows that  $F'_k$  is a spanning 1-regular subgraph of  $G_1 \times H$ , and hence a 1-factor of  $G_1 \times H$ . In a like manner mutually edge-disjoint 1-factors  $F'_1, F'_2, \dots, F'_{s+1}$  of  $G_1 \times H$  can be obtained from each of  $F_1, F_2, \dots, F_{s+1}$ , respectively. Therefore  $G_1 \times H$  is 1-factorable, which implies that  $G \times H$  is also 1-factorable as previously indicated.

*Corollary:* If  $G$  and  $H$  are regular graphs, and  $\chi_1(G) = \Delta(G)$ , then  $\chi_1(G \times H) = \Delta(G) + \Delta(H)$ .

We remark that the theorem gives a sufficient condition for 1-factorability which is, however, not a necessary condition, since 1-factorable products of two non-1-factorable graphs are known. An example of this is the cartesian product of the Petersen graph with a triangle.

P. E. Himelwright and J. E. Williamson,  
Grand Valley State College, and Southern Illinois University, USA

## REFERENCES

- [1] M. BEHZAD and G. CHARTRAND, *Introduction of the Theory of Graphs*, Allyn and Bacon, Inc. 1972.
- [2] V. G. VIZING, *On an estimate of the chromatic class of a p-graph* (Russian), Diskret. Analiz. 3, 25-30 (1964).