

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 29 (1974)
Heft: 4

Artikel: A note on Bernoulli numbers and polynomials
Autor: Carlitz, L.
DOI: <https://doi.org/10.5169/seals-29900>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A note on Bernoulli numbers and polynomials

Put

$$S_k = S_k(n) = \sum_{a=0}^{n-1} a^k.$$

It is well known that

$$S_1^2 = S_3, \quad 2S_1^4 = S_5 + S_7. \quad (1)$$

The general formula of this type was found by Stern [1, p. 20]:

$$2^{m-1} S_1^m = \sum_{2j < m} \binom{m}{2j+1} S_{2m-2j-1}. \quad (2)$$

We recall that [2, Ch. 2]

$$S_k(n) = \frac{B_{k+1}(n) - B_{k+1}}{k+1}, \quad (3)$$

where $B_{k+1}(n)$ is the Bernoulli polynomial of degree $k+1$ and $B_{k+1} = B_{k+1}(0)$. The Bernoulli polynomial may be defined by

$$\frac{ze^{xz}}{e^z - 1} = \sum_{k=0}^{\infty} B_k(x) \frac{z^k}{k!}. \quad (4)$$

Substituting from (3) in (2) we get

$$2^{m-1} S_1^m = \sum_{2j < m} \binom{m}{2j+1} \frac{B_{2m-2j}(n) - B_{2m-2j}}{2m-2j}. \quad (5)$$

Since $S_1 = n(n-1)/2$, it is clear that (5) is a polynomial identity for $n = 1, 2, 3, \dots$. Therefore we may write

$$(x(x-1))^m = 2 \sum_{2j < m} \binom{m}{2j+1} \frac{B_{2m-2j}(x) - B_{2m-2j}}{2m-2j}.$$

Since $B'_m(x) = mB_{m-1}(x)$, it follows that

$$m(2x-1)(x(x-1))^{m-1} = 2 \sum_{2j < m} \binom{m}{2j+1} B_{2m-2j-1}(x). \quad (6)$$

Conversely integration of (6) gives (5).

A slightly simpler formula is

$$(m+1)(x-1/2)^m = \sum_{2j \leq m} \binom{m+1}{2j+1} 2^{-2j} B_{m-2j}(x). \quad (7)$$

To prove (7) we make use of (4). Clearly

$$\sum_{k=0}^{\infty} B_k(x) \frac{z^k}{k!} = \frac{ze^{(x-1/2)z}}{e^{z/2} - e^{-z/2}},$$

so that

$$ze^{(x-1/2)z} = 2 \sum_{k=0}^{\infty} B_k(x) \frac{z^k}{k!} \sum_{j=0}^{\infty} \frac{(z/2)^{2j+1}}{(2j+1)!}.$$

Equating coefficients of z^{m+1} , we get (7).

Next we recall the expansion [2, p. 28]

$$B_k(x) = \sum_{s=0}^k \binom{k}{s} 2^{-s} D_s (x - 1/2)^{k-s} \quad (8)$$

where

$$D_k = 2^k B_k(1/2) = 2 (1 - 2^{k-1}) B_k. \quad (9)$$

It follows from (9) that

$$D_{2k+1} = 0, \quad (-1)^k D_{2k} > 0 \quad (k = 0, 1, 2, \dots). \quad (10)$$

The first few values are

$$D_0 = 1, \quad D_2 = -\frac{1}{3}, \quad D_4 = \frac{7}{15}, \quad D_6 = -\frac{31}{21}.$$

Thus (7) and (8) are an inverse pair. It is convenient to consider separately even and odd values of m and k . Replacing m by $2m$, (7) becomes

$$(2m+1) (x - 1/2)^{2m} = \sum_{j=0}^m \binom{2m+1}{2j} 2^{-2m+2j} B_{2j}(x). \quad (11)$$

Similarly, by (8) and (10)

$$B_{2k}(x) = \sum_{s=0}^k \binom{2k}{2s} 2^{-2k+2s} D_{2k-2s} (x - 1/2)^{2s}. \quad (12)$$

Substituting from (12) in (11), we get

$$(2m+1) (x - 1/2)^{2m} = \sum_{j=0}^m \binom{2m+1}{2j} 2^{-2m} \sum_{s=0}^j \binom{2j}{2s} 2^{2s} D_{2j-2s} (x - 1/2)^{2s},$$

so that

$$\sum_{j=s}^m \binom{2m+1}{2j} \binom{2j}{2s} 2^{2s} D_{2j-2s} = (2m+1) 2^{2m} \delta_{m,s}. \quad (13)$$

For odd values of the parameters, (7) and (8) yield

$$(2m+2) (x - 1/2)^{2m+1} = \sum_{j=0}^m \binom{2m+2}{2j+1} 2^{-2m+2j} B_{2j+1}(x), \quad (14)$$

and

$$B_{2k+1}(x) = \sum_{s=0}^k \binom{2k+1}{2s+1} 2^{-2k+2s} D_{2k-2s} (x - 1/2)^{2s+1}, \quad (15)$$

respectively. Hence

$$(2m+2) (x - 1/2)^{2m+1} = \sum_{j=0}^m \binom{2m+2}{2j+1} 2^{-2m} \sum_{s=0}^j \binom{2j+1}{2s+1} 2^{2s} D_{2j-2s} (x - 1/2)^{2s+1},$$

so that

$$\sum_{j=s}^m \binom{2m+2}{2j+1} \binom{2j+1}{2s+1} 2^{2s} D_{2j-2s} = (2m+2) 2^{2m} \delta_{m,s}. \quad (16)$$

The formulas (13), (16) evidently imply the following matrix formulas:

$$\left[\begin{pmatrix} 2m+1 \\ 2s \end{pmatrix} \right] \left[\begin{pmatrix} 2m \\ 2s \end{pmatrix} 2^{2s} D_{2m-2s} \right] = \left[\begin{pmatrix} 2m+1 \\ 2s \end{pmatrix} 2^{2m} \delta_{m,s} \right], \quad (17)$$

$$\left[\begin{pmatrix} 2m+2 \\ 2s+1 \end{pmatrix} \right] \left[\begin{pmatrix} 2m+1 \\ 2s+1 \end{pmatrix} 2^{2s} D_{2m-2s} \right] = \left[\begin{pmatrix} 2m+2 \\ 2s+1 \end{pmatrix} 2^{2m} \delta_{m,s} \right], \quad (18)$$

where $m, s = 0, 1, 2, \dots, N-1$ and N is either a positive integer or infinity. Thus in particular we have found inverses of the matrices

$$\left[\begin{pmatrix} 2m+1 \\ 2s \end{pmatrix} \right], \quad \left[\begin{pmatrix} 2m+2 \\ 2s+1 \end{pmatrix} \right] \quad (m, s = 0, 1, 2, \dots, N-1). \quad (19)$$

For applications it is convenient to state the following result.

Theorem 1. *The set of equations*

$$(m+1)x_m = \sum_{2j \leq m} \binom{m+1}{2j+1} 2^{-2j} y_{m-2j} \quad (m = 0, 1, 2, \dots) \quad (20)$$

is equivalent to the set

$$y_m = \sum_{2j \leq m} \binom{m}{2j} 2^{-2j} D_{2j} x_{m-2j} \quad (m = 0, 1, 2, \dots). \quad (21)$$

Separating even and odd values of m we have

Theorem 2. *The set of equations*

$$(2m+1)x_m = \sum_{j=0}^m \binom{2m+1}{2j} 2^{2j-2m} y_j \quad (m = 0, 1, 2, \dots) \quad (22)$$

is equivalent to the set

$$y_m = \sum_{j=0}^m \binom{2m}{2j} 2^{2j-2m} D_{2m-2j} x_j \quad (m = 0, 1, 2, \dots). \quad (23)$$

The set of equations

$$(2m+2)x_m = \sum_{j=0}^m \binom{2m+2}{2j+1} 2^{2j-2m} y_j \quad (m = 0, 1, 2, \dots) \quad (24)$$

is equivalent to the set

$$y_m = \sum_{j=0}^m \binom{2m+1}{2j+1} 2^{2j-2m} D_{2m-2j} x_j \quad (m = 0, 1, 2, \dots). \quad (25)$$

L. Carlitz, Duke University, USA

REFERENCES

- [1] P. BACHMANN, *Niedere Zahlentheorie*, Vol. 2, Teubner, Leipzig, 1910.
- [2] N. E. NÖRLUND, *Vorlesungen über Differenzrechnung*, Springer, Berlin, 1924.