

Zeitschrift:	Elemente der Mathematik
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	29 (1974)
Heft:	4
Artikel:	A new proof of the reciprocity theorem for Dedekind sums
Autor:	Berndt, Bruce C.
DOI:	https://doi.org/10.5169/seals-29901

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A new proof of the reciprocity theorem for Dedekind sums

If

$$\left((x) \right) = \begin{cases} x - [x] - \frac{1}{2}, & \text{if } x \text{ is not an integer,} \\ 0, & \text{otherwise,} \end{cases}$$

the classical Dedekind sum $s(h, k)$ is defined by

$$s(h, k) = \sum_{n=1}^{k-1} \left((h n/k) \right) \left((n/k) \right),$$

where k is a positive integer, and h is any integer. The most famous and useful property of these sums is the following result.

Reciprocity Theorem. If h and k are positive integers, and $(h, k) = 1$, then

$$s(h, k) + s(k, h) = -\frac{1}{4} + \frac{1}{12} \left(\frac{h}{k} + \frac{k}{h} + \frac{1}{hk} \right). \quad (1)$$

The beautiful, new monograph [1] by Rademacher and Grosswald presents four distinct proofs of (1). In this note, we give a completely new, short proof of (1).

Our proof is based on the classical Poisson summation formula. If f is of bounded variation on $[a, b]$,

$$\frac{1}{2} \sum_{n=a}^b' \{f(n+0) + f(n-0)\} = \int_a^b f(x) dx + 2 \sum_{n=1}^{\infty} \int_a^b f(x) \cos(2\pi n x) dx, \quad (2)$$

where the dash ' on the summation sign indicates that for the terms corresponding to $n = a$ and $n = b$, only $f(a+0)$ and $f(b-0)$, respectively, are counted. Let $a = 0$, $b = k$, and $f(x) = ((hx/k)) ((x/k))$ in (2). Observe that $f(0+0) = f(k-0) = 1/4$. On the right side of (2), let $x = ky$ in each of the integrals. An easy calculation [1, p. 23], yields

$$\int_0^k \left((h x/k) \right) \left((x/k) \right) dx = k/12 h.$$

Hence, (2) gives

$$\frac{1}{4} + s(h, k) = \frac{k}{12h} + 2k \sum_{n=1}^{\infty} I(k, h, n), \quad (3)$$

where

$$I(k, h, n) = \int_0^1 \left((h y) \right) \left((y) \right) \cos(2\pi n k y) dy.$$

We now evaluate $I(k, h, n)$. Write

$$\begin{aligned} I(k, h, n) &= \sum_{r=0}^{h-1} \int_{r/h}^{(r+1)/h} \left(h y - r - \frac{1}{2} \right) \left(y - \frac{1}{2} \right) \cos(2\pi n k y) dy \\ &= \int_0^1 \left(h y - \frac{1}{2} \right) \left(y - \frac{1}{2} \right) \cos(2\pi n k y) dy - \sum_{r=0}^{h-1} r \int_{r/h}^{(r+1)/h} \left(y - \frac{1}{2} \right) \cos(2\pi n k y) dy \\ &= \int_0^1 \left(h y - \frac{1}{2} \right) \left(y - \frac{1}{2} \right) \cos(2\pi n k y) dy - \sum_{j=1}^{h-1} \int_{j/h}^{(j+1)/h} \left(y - \frac{1}{2} \right) \cos(2\pi n k y) dy. \end{aligned}$$

Integrating by parts, we obtain

$$\begin{aligned} I(k, h, n) &= -\frac{1}{2\pi n k} \int_0^1 \left\{ 2hy - \frac{1}{2}(h+1) \right\} \sin(2\pi n k y) dy \\ &\quad + \frac{1}{2\pi n k} \sum_{j=1}^{h-1} \left\{ \left(\frac{j}{h} - \frac{1}{2} \right) \sin(2\pi n k j/h) - \frac{1}{2\pi n k} (1 - \cos(2\pi n k j/h)) \right\} \\ &= \frac{h}{(2\pi n k)^2} + \frac{1}{2\pi n k} \sum_{j=1}^{h-1} ((j/h)) \sin(2\pi n k j/h) + \frac{1}{(2\pi n k)^2} \sum_{j=0}^{h-1} \cos(2\pi n k j/h). \end{aligned}$$

The last sum on the right side above is 0 except when $n/h = \mu$, say, is a positive integer, in which case the sum is equal to h .

Using the above calculation, we find now that (3) yields

$$\begin{aligned} \frac{1}{4} + s(h, k) &= \frac{k}{12h} + \frac{h}{2\pi^2 k} \sum_{n=1}^{\infty} \frac{1}{n^2} \\ &\quad + \sum_{j=1}^{h-1} ((j/h)) \sum_{n=1}^{\infty} \frac{\sin(2\pi n k j/h)}{\pi n} + \frac{1}{2\pi^2 h k} \sum_{\mu=1}^{\infty} \frac{1}{\mu^2} \\ &= k/12h + h/12k - \sum_{j=1}^{h-1} ((j/h)) ((k j/h)) + 1/12hk, \end{aligned} \tag{4}$$

where we have used the well known Fourier series for $((x))$,

$$((x)) = -\sum_{n=1}^{\infty} \frac{\sin(2\pi n x)}{\pi n}.$$

Equation (4) is clearly equivalent to (1), and so the proof of the reciprocity theorem is complete.

The ideas employed above can be applied to many generalizations of the Dedekind sums. An account of this work will be given elsewhere.

Bruce C. Berndt, University of Illinois, Urbana, Illinois, USA

REFERENCE

- [1] H. RADEMACHER and E. GROSSWALD, *Dedekind Sums*, Carus Mathematical Monograph No. 16, Mathematical Association of America, Washington D.C., 1972.