Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 31 (1976)
Heft: 1

Artikel: On Yff's inequality for the brocard angle of a triangle
Autor: Bottema, O.
DOI: https://doi.org/10.5169/seals-31391

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 07.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

On Yff's Inequality for the Brocard Angle of a Triangle

1. Let $\alpha_1, \alpha_2, \alpha_3$ be the angles of a triangle and let ω be its Brocard angle. In 1963 Yff [1] conjectured that ω satisfies the inequality

$$8\,\omega^3 \leq \alpha_1\,\alpha_2\,\alpha_3\,. \tag{1}$$

This is a remarkable relation because it contains the angles proper and not, as is usually the case, in geometrical inequalities, their trigonometric representatives. (1) could be called a transcendental relation, while the usual ones are algebraic. There are not many statements of type (1) in elementary geometry.

Many mathematicians have tried in vain to prove (1); the present author knows about it because he was one of them. But now, quite recently, a short, elegant and ingenious proof was published in this journal by Faruk Abi-Khuzam [2]. It depends on the following lemma

$$\sin\alpha_1 \sin\alpha_2 \sin\alpha_3 \leq k\alpha_1\,\alpha_2\,\alpha_3\,, \tag{2}$$

k being the constant $(3\sqrt{3}/2\pi)^3$; for $\alpha_1 = \alpha_2 = \alpha_3 = \pi/3$ equality holds in (2).

In our opinion the proof, given for (2) is not completely satisfactory. Use is made of the infinite product

$$\sin x = x \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{\pi^2\,n^2}\right). \tag{3}$$

After substitution and re-ordering infinitely many factors the lefthand side of (2) is written as *one* infinite product. For further reduction the factors are written as the product of two linear expressions. All this seems rather light-hearted, the more so as it is well-known that the infinite product

$$\prod \left(1 - \frac{x}{\pi n}\right)\left(1 + \frac{x}{\pi n}\right) \tag{4}$$

is not absolutely convergent (see e.g. Whittaker-Watson, Modern Analysis, p. 33–34).

Here follows a more elementary proof of (2).

2. For $0 < \alpha_i < \pi$, $\alpha_1 + \alpha_2 + \alpha_3 = \pi$ we consider the function

$$F\,(\alpha_1, \alpha_2, \alpha_3) = \frac{\sin\alpha_1 \sin\alpha_2 \sin\alpha_3}{\alpha_1\,\alpha_2\,\alpha_3} \tag{5}$$

F is defined on the region G in the $(\alpha_1, \alpha_2, \alpha_3)$-space consisting of the points *inside* the triangle with the vertices $P_1\,(\pi, 0, 0)$, $P_2\,(0, \pi, 0)$, $P_3\,(0, 0, \pi)$. We define F on the perimeter of the triangle by

$$F\,(0, \alpha_2, \alpha_3) = \frac{\sin\alpha_2 \sin\alpha_3}{\alpha_2\,\alpha_3}\,,\ \alpha_2 \neq 0,\ \alpha_3 \neq 0,\ \alpha_2 + \alpha_3 = \pi$$

and analogously $F\,(\alpha_1, 0, \alpha_3)$ and $F\,(\alpha_1, \alpha_2, 0)$; furthermore $F\,(\pi, 0, 0) = F\,(0, \pi, 0) = F\,(0, 0, \pi) = 0$. F is now defined on a closed region $\bar{G}$; it is continuous and derivable

on $\bar{G}$; moreover as $0 \leq F < 1$ there is (at least) one point in $\bar{G}$ where F has its maximum value. By the usual procedure, in view of $\alpha_1 + \alpha_2 + \alpha_3 = \pi$, a maximum satisfies

$$\frac{\partial F}{\partial \alpha_1} = \frac{\partial F}{\partial \alpha_2} = \frac{\partial F}{\partial \alpha_3} (= \lambda) \ . \tag{6}$$

In G we have

$$\frac{\partial F}{\partial \alpha_1} = \frac{\sin\alpha_2 \sin\alpha_3}{\alpha_2 \alpha_3} \cdot \frac{\alpha_1 \cos\alpha_1 - \sin\alpha_1}{\alpha_1^2} = F\,(\cot \alpha_1 - \alpha_1^{-1}) \ ,$$

and, as $F \neq 0$, (6) implies

$$\cot\alpha_1 - \alpha_1^{-1} = \cot\alpha_2 - \alpha_2^{-1} = \cot\alpha_3 - \alpha_3^{-1} \ . \tag{7}$$

For $f = \cot\alpha - \alpha^{-1}$ we obtain $f' = -\sin^{-2}\alpha + \alpha^{-2} < 0$; f is therefore a decreasing function of α (we have $0 > f > -\infty$); hence (7) implies

$$\alpha_1 = \alpha_2 = \alpha_3 \ (= \pi/3) \ . \tag{8}$$

in this point we have $F = k$.

We must verify whether larger values appear on the boundary of $\bar{G}$. Between P_2 and P_3 yields

$$F = \frac{\sin\alpha_2 \sin\alpha_3}{\alpha_2 \alpha_3}, \ \alpha_2 + \alpha_3 = \pi$$

and by an argumentation analogous to the former, but now with two factors instead of three, it follows that for the maximum on P_2P_3 we have $\alpha_2 = \alpha_3 = \pi/2$ and $F = 4/\pi^2$, that is less than k. Hence $F \leq k$ on $\bar{G}$, which concludes the proof.

O. Bottema, Delft

REFERENCES

[1] P. Yff, *An analogue of the Brocard points*, Amer. Math. Monthly *70*, 500 (1963).
[2] Faruk Abi-Khuzam, *Proof of Yff's Conjecture on the Brocard Angle of a Triangle*, El. Math. *29* 141–142 (1974).

Aufgaben

Aufgabe 733. Let n be a positive integer ≥ 2. Let L be a line which intersects the $(n-1)$-dimensional hyperplanes containing the $(n-1)$-dimensional faces of a given n-dimensional simplex of vertices A_i $(i = 1, \ldots, n+1)$ in the uniquely determined points B_i. Prove that the n-dimensional volume of the convex hull of the midpoints of $\overline{A_iB_i}$ is zero. This extends the known results for $n = 2, 3$ for which the midpoints are collinear and coplanar, respectively.

M.S. Klamkin, Dearborn, Michigan, USA