

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 33 (1978)
Heft: 1

Artikel: Dissections of a Polygon
Autor: Kerr, Jeanne W. / Wetzel, John E.
DOI: <https://doi.org/10.5169/seals-32934>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ELEMENTE DER MATHEMATIK

Revue de mathématiques élémentaires – Rivista di matematica elementare

*Zeitschrift zur Pflege der Mathematik
und zur Förderung des mathematisch-physikalischen Unterrichts*

El. Math. Band 33 Heft 1 Seiten 1-24 10. Januar 1978

Dissections of a Polygon

1. Suppose that a plane oval O (a convex, open, non-empty set) is intersected by c lines forming c dissecting chords, and suppose these chords intersect in p points of O through each of which only two of the chords pass (fig. 1). We do not insist that each two of the chords intersect. Then the number R of disjoint (open) regions formed inside O is given by the pretty formula

$$R = 1 + c + p. \quad (1)$$

This formula, apparently first given in this setting by Norman Bauman [3], is analogous to a result proved in 1966 by Alfred Brousseau [2] for lines in the plane. A more extensive study of (1) and related formulas appears in [1].

For the sake of completeness, we include a proof of (1), by induction on c . The formula is correct for $c=0$. Suppose a new chord C is added to a set of c chords for which the formula holds, and suppose that C meets those chords in $k \geq 0$ points in O . Since the chord C is a segment, ray, or whole line, it is divided into $k+1$ disjoint (open) pieces by the k points of intersection. Each of these $k+1$ pieces divides a previous region into two parts. Consequently, adding C creates exactly $k+1$ new regions. The right side of (1) plainly increments by exactly $k+1$ when C is added, and it follows that (1) holds for the augmented arrangement also. This proves (1) by induction.

It is worth remarking that simple-connectivity, not convexity, is the crucial assumption; one can prove that (1) holds for any simply-connected domain in the plane dissected by cross-cuts (see [6]).

In this note we use (1) to solve some natural polygon dissection problems.

2. Let V_1, V_2, \dots, V_n be the vertices of a convex n -gon Π , named in cyclic order, and put $V_{n+1} = V_1$. For each $i = 1, 2, \dots, n$, select $x_i \geq 0$ points on the (open) edge $V_i V_{i+1}$ of Π . We divide the interior of Π (which is an oval) into regions by drawing all of the line segments determined by the vertices and the points chosen on the edges (fig. 2). If no three of these chords meet in a point inside the n -gon, how many regions are formed?

The solution is readily found using (1).




Figure 1. An oval dissected into 33 regions by a family of 10 lines.

Write σ_j for the sum of the $\binom{n}{j}$ products of the x_i 's taken j at a time, with the usual combinatorial conventions that $\sigma_j = 0$ and $\binom{n}{j} = 0$ when $j > n$.

It is easy to count the chords of various kinds. To facilitate the descriptions, we write V to refer to a vertex and E to refer to an edge point. There are clearly

$$c_1 = \binom{n}{2} - n$$

chords of the form VV , the diagonals of Π . Each point E on an edge can be joined to any of the far $n-2$ vertices, so there are

$$c_2 = (n-2)\sigma_1$$

chords of the form VE . And there evidently are

$$c_3 = \sigma_2$$




Figure 2. A dissected pentagon with data 0, 2, 1, 0, 1 divided into 116 regions.

chords of the form EE . Consequently, there are

$$c = c_1 + c_2 + c_3 = \binom{n}{2} - n + (n-2)\sigma_1 + \sigma_2$$

chords in this dissected oval.

To count the points of intersection, we rely on the observation that four points on Π , no three of which are collinear, determine six chords exactly two of which meet inside Π to make one point of intersection. There are, in all, $n + \sigma_1$ vertices and edge points on the n -gon, so there would be

$$\binom{n + \sigma_1}{4}$$

points determined inside Π but for the collinearities. We count the points lost due to collinearity and subtract them off.

Four of the $n + \sigma_1$ points may be selected on the same (closed) side in

$$\sum_{i=1}^n \binom{x_i + 2}{4}$$

ways, and three of the points may be chosen on one (closed) side and the fourth elsewhere in

$$\sum_{i=1}^n \binom{x_i+2}{3} (n-2+\sigma_1-x_i)$$

ways. Consequently, the c chords meet to form

$$p = \binom{n+\sigma_1}{4} - \sum_{i=1}^n \left[\binom{x_i+2}{4} + \binom{x_i+2}{3} (n-2+\sigma_1-x_i) \right]$$

points inside Π .

It follows from (1) that there are

$$R = 1 - n + \binom{n}{2} + (n-2)\sigma_1 + \sigma_2 + \binom{n+\sigma_1}{4} - \sum_{i=1}^n \left[\binom{x_i+2}{4} + \binom{x_i+2}{3} (n-2+\sigma_1-x_i) \right]$$

regions formed inside the partitioned polygon.

3. Other natural polygon dissection problems can be posed in this setting and solved using (1). Instead of partitioning the interior of Π by drawing all possible segments determined by the vertices and edge points, we could use only certain of these segments.

There are seven natural sets of dissecting chords to consider, indicated symbolically by (VV) , (VE) , (EE) , (VV, VE) , (VV, EE) , (VE, EE) , and (VV, VE, EE) ; and there are seven associated partition problems. In the (VE, EE) -problem, for example, all the line segments that join a vertex to an edge point or two edge points are drawn. The (VV) -problem is the classical case of an n -gon dissected by its diagonals. The situation studied in section 2 above is the (VV, VE, EE) -problem.

We use (1) to solve all of these problems. The chords of various kinds have already been counted. It is a little more complicated to count the points.

We say that a point P determined inside Π is of *type* $WX-YZ$ provided the two chords that meet at P are of types WX and YZ . Thus every interior point of intersection is of one of the six types $VV-VV$, $VV-VE$, $VV-EE$, $VE-VE$, $VE-EE$, and $EE-EE$. We count the points of each type separately.

There evidently are

$$p_1 = \binom{n}{4}$$

points of type $VV-VV$.

A point E can be selected in σ_1 ways, and then three vertices V (not including both endpoints of the edge on which E was chosen) can be picked in

$$\binom{n-2}{3} + 2\binom{n-2}{2} = \binom{n}{3} - \binom{n-2}{1}$$

ways. So there are

$$p_2 = \left[\binom{n}{3} - n + 2 \right] \sigma_1$$

points of type *VV-VE*.

Points of type *VV-EE* are obtained by choosing two edge points on different edges and then choosing two vertices, one on each side of the line joining the two chosen edge points (but choosing both vertices of either of the two edges that contain the chosen edge points is not permitted). So there clearly are

$$p_3 = \sum_{i < j} [(j-i)(n-j+i)-2] x_i x_j$$

$$= \sum_{i < j} (j-i)(n-j+i) x_i x_j - 2 \sigma_2$$

points of type *VV-VE*.

Points of type *VE-VE* arise in two different ways. The two edge points can be chosen on the same edge and any two of the $n-2$ far vertices chosen, which can be done in

$$\binom{n-2}{2} \sum_{i=1}^n \binom{x_i}{2}$$

ways; or the two edge points can be chosen on different edges and the two vertices chosen on the same side of the line through them, which can be done in

$$\sum_{i < j} \left[\binom{j-i}{2} + \binom{n-j+i}{2} \right] x_i x_j$$

$$= \binom{n}{2} \sigma_2 - \sum_{i < j} (j-i)(n-j+i) x_i x_j$$

ways. So in all there are

$$p_4 = \binom{n-2}{2} \sum_{i=1}^n \binom{x_i}{2} + \binom{n}{2} \sigma_2 - \sum_{i < j} (j-i)(n-j+i) x_i x_j$$

points of type *VE-VE*.

Points of type *VE-EE* also arise in two ways: the three edge points can come from three different edges, or two edge points can come from the same edge and the third from another edge. There are clearly $n\sigma_3$ points of the first sort and

$$(n-2) \sum_{i=1}^n \binom{x_i}{2} (\sigma_1 - x_i)$$

points of the second. So there are

$$p_5 = n \sigma_3 + (n-2) \sum_{i=1}^n \binom{x_i}{2} (\sigma_1 - x_i)$$

points of type *VE-EE*.

There are three ways that points of the type *EE-EE* arise, depending on whether the four edge points are chosen on different sides, which can be done in σ_4 ways; or two are chosen on the same side and the other two on different sides, which can be accomplished in

$$\sum_{i=1}^n \binom{x_i}{2} \sum_{\substack{j < k \\ j, k \neq i}} x_j x_k = \sum_{i=1}^n \binom{x_i}{2} (\sigma_2 - \sigma_1 x_i + x_i^2)$$

ways; or two points are chosen on each of two sides, which can be accomplished in

$$\sum_{i < j} \binom{x_i}{2} \binom{x_j}{2}$$

ways. So in all there are

$$p_6 = \sigma_4 + \sum_{i=1}^n \binom{x_i}{2} (\sigma_2 - \sigma_1 x_i + x_i^2) + \sum_{i < j} \binom{x_i}{2} \binom{x_j}{2}$$

points of type *EE-EE*.

It is an easy matter now to use (1) to write formulas for the number R of regions formed in each of the various cases.

(*VV*). Here $c = c_1$ and $p = p_1$, and so

$$R = 1 + \binom{n}{2} - n + \binom{n}{4} = \binom{n-1}{2} + \binom{n-1}{3} + \binom{n-1}{4},$$

a familiar formula (see [4], [5] p. 99–107, and [7] p. 13, 108–112).

(*VE*). Here $c = c_2$ and $p = p_4$, and so $R = 1 + c_2 + p_4$.

(*EE*). Here $c = c_3$ and $p = p_6$, and so $R = 1 + c_3 + p_6$.

(*VV, VE*). In this case we have $c = c_1 + c_2$ chords, which meet to form all points of the types *VV-VV*, *VV-VE*, and *VE-VE*. So $p = p_1 + p_2 + p_4$, and $R = 1 + c_1 + c_2 + p_1 + p_2 + p_4$.

(*VV, EE*). Here we have $c = c_1 + c_3$ chords, which meet to form all points of the types *VV-VV*, *VV-EE*, and *EE-EE*. So $p = p_1 + p_3 + p_6$, and $R = 1 + c_1 + c_3 + p_1 + p_3 + p_6$.

(*VE, EE*). There are $c = c_2 + c_3$ chords, which meet to form all points of the types *VE-VE*, *VE-EE*, and *EE-EE*. So $p = p_4 + p_5 + p_6$, and $R = 1 + c_2 + c_3 + p_4 + p_5 + p_6$.

(*VV, VE, EE*). In this problem there are $c = c_1 + c_2 + c_3$ chords, which meet to form all points of all six types. So $p = p_1 + p_2 + p_3 + p_4 + p_5 + p_6$, and $R = 1 + c_1 + c_2 + c_3 + p_1 + p_2$

$+p_3+p_4+p_5+p_6$. It is not entirely obvious that this formula is the same as the formula obtained in section 2 above for this problem.

Jeanne W. Kerr, University of California, San Diego, La Jolla, Cal.,
and John E. Wetzel, University of Illinois, Urbana, Ill., USA

ACKNOWLEDGMENT

The authors would like to thank Marianne Wojciechowski Jankowski for preparing the figures.

REFERENCES

- 1 G. L. Alexanderson and John E. Wetzel: Dissections of a plane oval, to appear in Amer. Math. Monthly.
- 2 Bro. U. Alfred «Brousseau»: A mathematician's progress. Math. Teacher 59, 722–727 (1966).
- 3 Norman Bauman: Solution to E 2359. Amer. Math. Monthly 80, 561–562 (1973).
- 4 J. W. Freeman: The number of regions determined by a convex polygon. Math. Mag. 49, 23–25 (1976).
- 5 Ross Honsberger: Mathematical Gems, Dolciani Mathematical Expositions, No. 1. Math. Assoc. of America, Washington, D.C., 1973.
- 6 John E. Wetzel: Dissections of a simply-connected plane domain, to appear in Amer. Math. Monthly.
- 7 A. M. Yaglom and I. M. Yaglom: Challenging Mathematical Problems with Elementary Solutions, Vol. 1. Holden-Day, San Francisco 1964.

Extremaleigenschaften rotationssymmetrischer Kegelstümpfe im gewöhnlichen Raum (1. Teil)

In dieser Note behandeln wir ein Extremalproblem, Teilproblem eines viel allgemeineren, unter III zu erläuternden Extremalproblems über konvexe Rotationskörper, und letzteres ordnet sich einem Hauptproblem über allgemeine konvexe Körper unter, das unter II kurz dargelegt wird. Die gesonderte Behandlung des Kegelstumpfproblems rechtfertigt sich durch die mit gesicherten Teilresultaten untermauerte Vermutung, dass die Extremalkörper unseres Spezialproblems mit Minimumeigenschaften dieselben im Problem III wenigstens teilweise beibehalten. Weitergehende Untersuchungen bezüglich der notwendigen Bedingungen für Extrema im Problem III lassen sogar den Schluss zu, dass die Aussonderung der Kegelstümpfe nicht zu umgehen ist.

Unsere Ausführungen beziehen sich ausschliesslich auf den gewöhnlichen Raum und die gewöhnliche Ebene.