**Zeitschrift:** Elemente der Mathematik

**Herausgeber:** Schweizerische Mathematische Gesellschaft

**Band:** 37 (1982)

Heft: 2

**Artikel:** The forcibly-tree and forcibly-unicyclic degree sequences

**Autor:** Exoo, Geoffrey / Harary, Frank

**DOI:** https://doi.org/10.5169/seals-36388

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 29.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# The forcibly-tree and forcibly-unicyclic degree sequences

Dedicated to Frank Boesch

### 1. Introduction

We follow the notation and terminology of [3]. Thus the degree sequence  $\pi$  of a graph G is written  $d_1 \ge d_2 \ge \cdots \ge d_p$ . A given nonincreasing sequence  $\sigma$  of positive integers is called *graphical* if  $\sigma$  is the degree sequence of some graph. Criteria for  $\sigma$  to be graphical are well known; see [3], p. 58, or Hakimi and Schmeikel [2] which presents a recent survey of the literature on graphical degree sequences.

Let P be a property of graphs, such as hamiltonian, tree, unicyclic, etc. A graphical sequence  $\pi$  is called *potentially-P* if there exists a graph G which realizes  $\pi$  having property P. Similarly  $\pi$  is *forcibly-P* if it is not only potentially-P but every graph which realizes  $\pi$  satisfies property P. The latter concept was introduced by Nash-Williams [4] in his discussion of forcibly-hamiltonian degree sequences.

It is easy to see [3], p. 62, that a degree sequence  $(d_1, d_2, ..., d_p)$  is potentially-tree if and only if  $2q = \sum d_i$ , every  $d_i$  is positive and q = p - 1.

Potentially-unicyclic degree sequences were characterized by Boesch and Harary [1] as precisely those graphical sequences  $d_1, ..., d_p$  such that  $d_3 > 1, d_p > 0$  and  $\sum d_i = 2p$ . Our present object is to characterize forcibly-tree and forcibly-unicyclic degree sequences.

**Theorem 1.** A graphical degree sequence  $d_1 \ge d_2 \ge \cdots d_p \ge 1$  is forcibly-tree if and only if either p = 2 and  $d_1 = d_2 = 1$  or  $d_3 = 1$  and  $d_1 + d_2 = p$ .

Proof: If a tree T contains a path of length 4, let  $v_1v_2v_3v_4v_5$  be its points. Let G(T) be the graph with point set V(T), whose line set is  $E(G) = E(T) \cup \{v_1v_5, v_2v_4\} - \{v_1v_2, v_4v_5\}$ . Then G(T) is not a tree but has the same degree sequence as T. Hence no forcibly-tree degree sequence can be realized by a tree containing  $P_5$  and so is only realized by  $K_2$ , a star or a double star. The latter two families of trees satisfy  $d_3 = 1$  and  $d_1 + d_2 = p$ .

For the converse it is clear that a degree sequence of the given type can only be the degree sequence of a double star, a star or  $K_2$ . These are precisely the trees with diameter at most three.  $\Box$ 

**Theorem 2.** A graphical degree sequence  $d_1 \ge d_2 \ge \cdots \ge d_p \ge 1$  is forcibly-unicyclic if and only if it is either one of the sequences  $2^3$ ,  $2^4$ ,  $2^5$ ,  $d_2^4$   $d_1^{d-2}$ ,  $d_2^3$   $d_1^{d-2}$ , where  $d_1 = d \ge 3$ , or satisfies  $d_3 \ge 2$ ,  $d_4 = 1$  and  $d_1 + d_2 + d_3 = p + 3$ .

Proof: The first step is to list a family of unicyclic graphs with the property that any unicyclic supergraph of one of these does not have a forcibly-unicyclic degree sequence. These minimal unicyclic graphs are the cycles  $C_n, n \ge 6$ , and the graphs in figure 1.




Figure 1. Minimal unicyclic graphs with nonforcible degree sequences.

We also show in figure 1 how to transform each minimal graph into a nonunicyclic graph with the same degree sequence. In each case it is immediate that any unicyclic graph which contains the given graph as a subgraph can be similarly transformed. Of course each cycle  $C_n$ ,  $n \ge 6$ , is sequentially equivalent to  $C_{n-3} \cup K_3$ .




Figure 2. The graphs with forcibly-unicyclic degree sequences.

Next we determine and display in figure 2 the unicyclic graphs which contain no subgraph from the forbidden family, i.e., no long cycle and none of the graphs in figure 1.

Evidently the degree sequences of the graphs in figure 2 are precisely those given in the statement of theorem 2. It is easy to verify that each of these is forcibly-unicyclic. In later communications we plan to study forcibly-P degree sequences for other properties P.

Geoffrey Exoo, Bell Laboratories, Holmdel, N.J. 07733 Frank Harary, University of Michigan, Ann Arbor, Mich. 48109

### REFERENCES

- 1 F.T. Boesch and F. Harary: Unicyclic realizability of a degree list. Networks 8, 93-96 (1978).
- 2 S.L. Hakimi and E.P. Schmeikel: Graphs and their degree sequences: A survey. Theory and Applications of Graphs. Springer Lecture Notes 642, 225-235 (1978).
- 3 F. Harary: Graph Theory. Addison-Wesley, Reading 1969.
- 4 C. St. J.A. Nash-Williams: On hamiltonian circuits in finite graphs. Proc. Am. Math. Soc. 17, 466-467 (1966).