Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 37 (1982)
Heft: 2

Artikel: On lattice polytopes having interior lattice points
Autor: Zaks, J. / Perles, M.A. / Wills, J.M.
DOI: https://doi.org/10.5169/seals-36389

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 09.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

On lattice polytopes having interior lattice points

Let $V(K)$ denote the volume of a convex body K in the Euclidean d-space E^d, and let $G(K)$ and $G^0(K)$ be defined by $G(K)=\operatorname{card}(K\cap Z^d)$ and $G^0(K)=\operatorname{card}(\operatorname{int} K\cap Z^d)$, where Z^d denotes the set of all the lattice points in E^d, i.e., points having only integral coordinates.
The following theorems of Minkowski are well known.

Theorem A (see [2], p. 76). *If K is a centrally symmetric convex body in E^d and $G^0(K)=1$, then $V(K)\leq 2^d$.*

Theorem B (see [2], p. 79 and 96). *If K is a centrally symmetric convex body in E^d and $G^0(K)=1$, then $G(K)\leq 3^d$; if $G^0(K)=1$ and $V(K)=2^d$, then K is a convex polytope and it has at most $2^{d+1}-2$ facets.*

There are no analogues of theorems A and B for convex bodies which are not centrally symmetric. The purpose of this paper is to give some information on $V(P)$ and $G(P)$ in the case where P is a lattice polytope (i.e., P has all of its vertices in Z^d) satisfying $G^0(P)=n, n\geq 1$. We will see that V nad G behave very different from the centrally symmetric case.
Let $g(d,n)$ and $v(d,n)$ be defined for $n\geq 1$ and $d\geq 2$ as follows: $g(d,n)=\sup\{G(P)\mid P\subset E^d, G^0(P)=n\}$ and $v(d,n)=\sup\{V(P)\mid P\subset E^d, G^0(P)=n\}$.
Scott, [3] proved that $g(2,1)=10$ and that $g(2,n)=3n+6$ for all $n\geq 2$. Our main result is the following

Theorem. *For all $d, d\geq 4$, and for all $n, n\geq 1$,*

$$v(d,n)\geq \frac{n+1}{d!}\,2^{2^{d-a}} \quad \text{and} \quad g(d,n)\geq \frac{n+1}{6(d-2)!}\,2^{2^{d-a}},$$

where $a=0.5856\ldots$; $v(3,n)\geq 6(n+1)$, $g(3,n)\geq 16n+23$, $v(4,1)\geq 147$ and $g(4,1)\geq 680$.

We need the following

Lemma. *If $(a_n)_{n=1}^{\infty}$ is defined by $a_1=2$ and $a_d=\prod_{i=1}^{d-1} a_i+1$ for $d\geq 2$, then*

(i) $a_1=2, a_2=3, a_3=7, a_4=43$ and $a_5=1807$;

(ii) for all d, $d\geq 1$, $1-\sum_{i=1}^{d}\frac{1}{a_i}=\left(\prod_{i=1}^{d} a_i\right)^{-1}$;

(iii) for all d, $d\geq 4$, $\prod_{i=1}^{d} a_i\geq 2^{2^{d-a}}$ where $a=0.5856\ldots$

Proof of the lemma: (i) is trivial, (ii) is true for $d=1$, and if it is true for d, then by induction

$$1-\sum_{i=1}^{d+1}\frac{1}{a_i}=\left(\prod_{i=1}^{d}a_i\right)^{-1}-\frac{1}{a_{d+1}}=\left(\prod_{i=1}^{d+1}a_i\right)^{-1}.$$

Let the real a be defined by $2^{2^{d-a}}=a_1a_2a_3a_4=1806$, with $d=4$; thus $a=0.5856\ldots$ Suppose (iii) holds for d, then

$$\prod_{i=1}^{d+1}a_i>\left(\prod_{i=1}^{d}a_i\right)^2\geq 2^{2\cdot 2^{d-a}}=2^{2^{d+1-a}},$$

hence it holds for $d+1$.

Proof of the theorem: Let $k_i=a_i$ for $i=1,\ldots,d-1, k_d=2(a_d-1)$, and let S_1^d be the d-simplex defined by

$$S_1^d=\left\{(x_1,\ldots,x_d)\in E^d \mid x_i\geq 0,\quad \sum_{i=1}^{d}\frac{x_i}{k_i}\leq 1\right\}.$$

By (ii) we have

$$\sum_{i=1}^{d}\frac{1}{k_i}<1 \quad\text{and}\quad \frac{1}{k_d}+\sum_{i=1}^{d}\frac{1}{k_i}=1,$$

hence $(1,\ldots,1)\in \operatorname{int} S_1^d$ and $(1,\ldots,1,2)\notin \operatorname{int} S_1^d$; $k_1<k_2<\cdots<k_d$ imply that $G^0(S_1^d)=1$. For general $n, n\geq 1$, replace $k_d=2(a_d-1)$ by $k_d=(n+1)(a_d-1)$, and obtain the d-simplex S_n^d. It follows that the only lattice points of $\operatorname{int} S_n^d$ are $(1,\ldots,1,j)$ for $1\leq j\leq n$, hence $G^0(S_n^d)=n$.
As for the volume of S_n^d, $V(S_n^d)=(1/d!)\prod_{i=1}^{d}k_i=((n+1)/d!)\,(a_d-1)^2$ and by (iii):

$$a_d-1\geq 2^{2^{d-1-a}},\qquad \text{therefore}\quad v(d,n)\geq\frac{n+1}{d!}\,2^{2^{d-a}}.$$

To compute $G(S_d^n)$ let $d\geq 3$ and $S=S_n^d\cap\{x_1=0\}\cap\{x_2=0\}$.
It is easy to see that for each $t\in \operatorname{aff} S$
$G(S+t)\leq G(S)$ and equality iff x is a lattice point.
If C_{d-2} denotes the unit cube in $\operatorname{aff} S$ and V_{d-2} the volume in $\operatorname{aff} S$, we have

$$V_{d-2}(S)=\int_{C_{d-2}}G(S+t)\,dt<G(S)<G(S_n^d).$$

With 6 $V_{d-2}(S)=d(d-1)\,V(S_n^d)$ we have

$$G(S_n^d)\leq\frac{d(d-1)}{6}\,V(S_n^d)=\frac{n+1}{6(d-2)!}\,2^{2^{d-a}}.$$

The case $d=3$: Clearly: $V(S_n^3)=6(n+1)$.
To compute $G(S_n^3)$ we count the integers $x_3 \geq 0$ satisfying

$$3(n+1)x_1+2(n+1)x_2+x_3 \leq 6(n+1)$$

for $(x_1,x_2)=(0,0), (0,1), (0,2), (1,0), (1,1)$ and $(2,0)$.
Easy computation gives $G(S_n^3)=16n+23$.
The case $d=4$: $V(S_1^4)=147$.
To compute $G(S_1^4)$ we count the pairs (x_3,x_4) of integers $x_3 \geq 0, x_4 \geq 0$ satisfying

$$42x_1+28x_2+7x_3+x_4 \leq 84$$

for $(x_1,x_2)=(0,0), (0,1), (0,2), (0,3), (1,0), (1,1), (2,0)$.
Counting yields $G(S_1^4)=680$.
This completes the proof of the theorem.
We raise the following *conjecture*:

$$v(d,n)<\infty \quad \text{for all} \quad d \geq 3 \quad \text{and} \quad n \geq 1 .$$

We remark that the conjecture implies $g(d,n)<\infty$ by Blichfeldt ([1], p.55): $G(P) \leq d!\,V(P)+d$ for nondegenerate lattice polytopes (compare [4]).
Similar problems may be asked for the number of i-dimensional faces of convex lattice polytopes P satisfying $G^0(P)=n$, where $0 \leq i \leq d-1$.

J. Zaks, University of Haifa and C.R.M.A., University of Montreal,
M. A. Perles, Hebrew University, Jerusalem,
J. M. Wills, University of Siegen, West Germany

REFERENCES

1 C.G. Lekkerkerker: Geometry of numbers. Wolters-Noordhoff, Groningen 1969.
2 H. Minkowski: Geometrie der Zahlen. Teubner, Leipzig 1910.
3 P.R. Scott: On convex lattice polygons. Bull. Austr. Math. Soc. *15*, 395-399 (1976).
4 J.M. Wills: Gitterzahlen und innere Volumina. Comm. Math. Helv. *53*, 508-524 (1978).

© 1982 Birkhäuser Verlag, Basel 0013-6016/82/020044-03$1.50+0.20/0

Kleine Mitteilungen

A note on the successive remainders of the exponential series

1. For real $x \neq 0$, we have by Taylor's theorem

$$e^x=1+\frac{x}{1!}+\frac{x^2}{2!}+\cdots+\frac{x^n}{n!}+\frac{x^{n+1}}{(n+1)!}\phi_n(x), \tag{1.1}$$

where

$$\phi_n(x)=e^{x\theta_n(x)}, \qquad 0<\theta_n(x)<1 . \tag{1.2}$$