**Zeitschrift:** Elemente der Mathematik

**Herausgeber:** Schweizerische Mathematische Gesellschaft

**Band:** 38 (1983)

Heft: 1

**Artikel:** The impossibility of a tesselation of the plane into equilateral triangles

whose sidelengths are mutually different, one of them being minimal

Autor: Scherer, Karl

**DOI:** https://doi.org/10.5169/seals-37174

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 01.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

### ELEMENTE DER MATHEMATIK

Revue de mathématiques élémentaires - Rivista di matematica elementare

Zeitschrift zur Pflege der Mathematik und zur Förderung des mathematisch-physikalischen Unterrichts

El. Math. Band 38 Nr. 1 Seiten 1-32 Basel, den 10. Januar 1983

# The impossibility of a tesselation of the plane into equilateral triangles whose sidelengths are mutually different, one of them being minimal

**Theorem.** There is no tesselation of the euclidean plane  $\mathbb{R}^2$  into equilateral triangles whose sidelengths are mutually different, one of them being minimal.

Proof: Assume that there is such a tesselation of  $\mathbb{R}^2$  into equilateral triangles  $t_i$ ,  $i \in I$ , where I is an arbitrary set of indices. We shall eventually see that this assumption leads to a contradiction.

For  $i \in I$ , let  $l_i$  denote the sidelength of  $t_i$ . Let l be the minimum of the sidelengths. By scaling we can attain l = 1. Then the area of each triangle is at least

$$\frac{1}{4} \cdot \sqrt{3}$$
.

Therefore the triangle can be enumerated and we can assume  $I = \mathbb{N}$  and  $l_1 = 1$ . For each  $i \in \mathbb{N}$  led  $d_i$  denote the boundary of  $t_i$ . Further we define D to be the 'grid' of the tesselation:  $D = \bigcup \{d_i, i \in \mathbb{N}\}$ .


If n and  $\varepsilon$  are positive real numbers, the points

$$P_1 = \left(-\frac{n}{2} - \frac{\varepsilon}{2}, \frac{\varepsilon}{2}\sqrt{3}\right), \qquad P_2 = \left(-\frac{n}{2}, 0\right), \quad P_3 = \left(\frac{n}{2} - \frac{\varepsilon}{2}, \frac{\varepsilon}{2}\sqrt{3}\right)$$
and 
$$P_4 = \left(\frac{n}{2}, 0\right)$$

define a parallelogram in  $\mathbb{R}^2$  with the sidelengths  $\varepsilon$  and n and with angles of 60 and 120 degrees. Let  $S(n,\varepsilon)$  denote the union of the three sides  $\overline{P_1P_2}$ ,  $\overline{P_2P_3}$  and  $\overline{P_3P_4}$  (fig. 1).

More generally we denote by  $S(n,\varepsilon)$  any subset of the plane that can be obtained from this special  $S(n,\varepsilon)$  by translations, rotations and reflections. If any such  $S(n,\varepsilon)$  is given, let  $R_n$  denote the interior of the associated parallelogram.

We show that there are a strongly decreasing sequence of positive numbers  $n_j, j \in \mathbb{N}$ , with  $n_j \le n_{j-1} - 1$  ( $j \ge 2$ ) and for each j a number  $\varepsilon_j \in (0, \infty)$  as well as a set  $S_{n_j} = S(n_j, \varepsilon_j)$ , contained in D, such that  $S_{n_j}$  suffices one of the following two properties (or both):






Figure 2

- a) There is a natural number k depending on j such that  $n_i = l_k$  and  $R_{l_k} \cap t_k = \phi$ .
- b) The 'base' of  $S_{n_i}(\overline{P_2P_3})$  in fig. 1) is not a side of a triangle of the tesselation.

The numbers  $n_j$  and the associated sets  $S_{n_j}$  will be constructed by induction on j. Since  $n_j \le n_{j-1} - 1$  for  $j \ge 2$ , some of the numbers  $n_j$  must be negative. But we assumed that each  $n_j$  is positive. Thus we get a contradiction and the theorem is proved.

The induction is performed in two steps.


### First step

We show that there is a subset  $S_{n_1}$  of D for some positive real number  $n = n_1$  which has property a) or property b).

Look at figure 2. Let  $P_1$ ,  $P_2$  and  $P_3$  be the vertices of  $t_1$ . Since  $t_1$  is the smallest triangle of the tesselation, each neighbor of  $t_1$  (i.e. each triangle that shares a boundary segment of positive measure with  $t_1$ ) must be larger than  $t_1$ . It is obvious that this is only possible if there are only three neighbors of  $t_1$ , say  $t_2$ ,  $t_4$  and  $t_6$ , and if (up to symmetry) they are arranged like in figure 2.

Hence there exist three triangles in the tesselation that have only a vertex in common with  $t_1$ . We can assume that these are the triangle  $t_3$ ,  $t_5$  and  $t_7$  and that the triangles  $t_2$ ,  $t_3$ ,  $t_4$ ,  $t_5$ ,  $t_6$  and  $t_7$  are arranged around  $t_1$  in a clockwise manner.

Let us call  $t_i$ , i=2,...,7, the surrounding triangles. The vertices of the surrounding triangles that are not points of  $t_1$  may be called the outer corners. Clearly each surrounding triangle has exactly two outer corners. For each surrounding triangle  $t_k$ , let  $P_{2k}$  and  $P_{2k+1}$  be the outer corners written clockwise (see fig. 3). Since the triangles of the tesselation have pairwise different sizes, we get the inequalities  $P_5 \neq P_6$ ,



El. Math., Vol. 38, 1983

 $P_9 \neq P_{10}$  and  $P_{13} \neq P_{14}$ .

We now have to distinguish the two cases wether  $t_2$  is larger or smaller than  $t_3$ .

Case 1. We assume that  $l_2 > l_3$ .

Consider figure 3. If D would contain a horizontal straight line whose right endpoint is  $P_7$ , the points  $P_6$  and  $P_7$  would define a set  $S_{l_3} \subset D$  with  $R_{l_3} \cap t_3 = \phi$ .

If this is not the case, D contains a straight line which elongates  $\overline{P_6P_7}$  over  $P_7$  and therefore a straight line whose left endpoint is  $P_7$ , too.

If  $P_7 \neq P_8$ , the points  $P_2$  and  $P_7$  define a set  $S_{n_1} \subset D$  that has property b). So we may assume  $P_7 = P_8$ .

Now consider all systems of straight lines that may start at  $P_9$  (fig. 4). In the cases i) and ii) of figure 4 there is a set  $S_{l_4} \subset D$  with  $R_{l_4} \cap t_4 = \phi$ , defined by  $P_7$  and  $P_9$ . In case iii) there is a set  $S_{l_4} \subset D$  with  $R_{l_4} \cap t_4 = \phi$ , defined by  $\overline{P_2 P_{11}}$ ,  $\overline{P_2 P_9}$  and the elongation of  $P_8 P_9$  over  $P_9$ . Therefore we may assume that the lines starting from  $P_9$  look as in figure 4 iv).

This implies  $l_4 > l_5$ .

Two further quite similar argumentations – we now have  $l_4 > l_5$  like we had  $l_2 > l_3$ , and we will get  $l_6 > l_7$  – show that if there is not a set  $S_{n_1} \subset D$  with property a) or b), the conditions  $P_{11} = P_{12}$  and  $P_4 = P_{15}$  must be fulfilled. Moreover, the line systems starting at  $P_5$ ,  $P_9$  and  $P_{13}$  are of type iv) of figure 4 (see fig. 5).

Let  $d_1$ ,  $d_2$  and  $d_3$  denote the (positive) lengths of  $\overline{P_5P_6}$ ,  $\overline{P_9P_{10}}$  and  $\overline{P_{13}P_{14}}$ , respectively. Then each  $d_i$ , i=1,2,3 is a sum of sidelengths of some triangles of the tesselation. Therefore  $d_i > l_1$  for i=1,2,3. Hence  $d_1 + d_2 + d_3 > 3$   $l_1$ .

Comparing the sidelength of the seven triangles  $t_1$ ,  $t_2$ ,  $t_3$ ,  $t_4$ ,  $t_5$ ,  $t_6$  and  $t_7$  yields the equations  $l_2 = l_1 + l_7$ ,  $l_2 = l_3 + d_1$ ,  $l_4 = l_1 + l_3$ ,  $l_4 = l_5 + d_2$ ,  $l_6 = l_1 + l_5$ ,  $l_6 = l_7 + d_3$ . They tell us that  $3 l_1 = d_1 + d_2 + d_3$  in contradiction to the last inequality.


Therefore in case 1 there must be a set  $S_{n_1} \subset D$  with property a) or b).


Case 2. We assume that  $l_2 < l_3$ .


If there is not a set  $S_{l_2} \subset D$  with  $R_{l_2} \cap t_2 = \phi$  defined by  $P_4$  and  $P_5$ , the situation at  $P_4$  can only be as indicated in figure 6. Then the points  $P_1$  and  $P_4$  define a set  $S_{l_2} \subset D$  with  $P_{l_2} \cap t_2 = \phi$ .

### **Second step**

A set  $S_n (= S_{n_i}) \subset D$  may be given that has property a) or b). It will be shown that there is a set  $S_m \subset D$  with  $m \le n-1$  which has property a). We then put  $n_{i+1} := m$ .










Figure 7b

Figure 7c

After a suitable translation, rotation and reflection if necessary we can assume that the given  $S_n$  has the position shown in figure 1. With the notations of figure 1, let  $t_r$  be the triangle of the tesselation which 'stands on the base of  $S_n$  at the left corner', which means that  $t_r$  is defined by the three conditions  $P_2 \in t_r$ ,  $R_n \cap t_r \neq \phi$ ,  $\overline{P_2 P_3}$  contains a side of  $t_r$  (see fig. 7a). The vertices of  $t_r$  may be  $P_2$ ,  $P_5$  and  $P_6$ , where  $P_5 \in \overline{P_2 P_3}$ . Since  $S_n$  has property a) or b),  $P_3$  is different from  $P_5$ .

Let  $t_s$  denote the neighbor of  $t_r$  at the side  $\overline{P_5P_6}$  such that  $P_5 \in t_s$ . We now have to distinguish whether  $l_s$  is greater or less than  $l_r$ .

Case 1. If  $l_s > l_r$ , the points  $P_2$  and  $P_6$  define a set  $S_{l_r} \subset D$  with  $R_{l_r} \cap t_r = \phi$  (see fig. 7 b). Moreover,  $l_r \le n-1$  because  $S_n \subset D$  and  $l_1 = 1$ . So we can define  $m = l_r$ .

Case 2. If  $l_s < l_r$ , let  $P_7$  denote the vertex of  $t_s$  that is not a point of  $t_r$  and let  $P_8$  denote its third vertex (fig. 7c). An examination of all possible line systems starting at  $P_7$  (they are listed in fig. 4) shows that in each case there is a set  $S_{l_s} \subset D$  with  $R_{l_s} \cap t_s = \phi$ , either defined by the points  $P_5$  and  $P_7$  or by  $P_7$  and  $P_8$ . Moreover,  $l_s < l_r \le n-1$ . So we can define  $m = l_s$ .

Hence the induction step is completed, q.e.d.

Karl Scherer, Universität Kaiserslautern

#### REFERENCES

- 1 W. Tutte: Squaring the Square. Can. J. Math. 1950, 197-209.
- 2 Ross Honsberger: Ingenuity in Mathematics. Mathematical Association of America 23, Yale University, 1970.

© 1983 Birkhäuser Verlag, Basel

0013-6018/83/010001-04\$1.50+0.20/0

## Quelques considérations concernant le problème de l'aiguille de Buffon dans l'espace euclidien $E_n$

0. Soit  $E_n$  l'espace euclidien à n dimensions de coordonnées  $x_1, ..., x_n$ . La mesure élémentaire cinématique dans  $E_n$ , invariante par rapport au groupe de mouvements euclidiens, est [1]:

$$dK = dP \wedge dO_{n-1} \wedge \cdots \wedge dO_1, \tag{1}$$