

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 38 (1983)
Heft: 1

Artikel: A remark on a paper by A. Grytczuk
Autor: Redmond, Don
DOI: <https://doi.org/10.5169/seals-37178>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

As the result of this procedure, all the points of H' other than w have degree d and the degree r of the point w is odd, since the total deficiency s is odd. Finally we remove any $(d-r)/2$ lines of the 1-factor F_{r+1} from H' and join w with the resulting $d-r$ points of degree $d-1$. The graph so constructed is d -regular, contains G , and has order $p+d+2$.

We now show by a family of examples that $d+2$ is best possible. Let G be obtained from the complete graph K_{p-1} with $p \geq 5$ by subdividing just one line by the insertion of a new point of degree 2; the graph G_2 in figure 1 illustrates $p=5$. Then we can readily see that if p is odd, at least $d+2$ new points are needed to construct a d -regular supergraph of G . \square

Remark 1. The smallest d -regular supergraph H will of course depend on the structure of G and its order can range between p and $p+d+2$.

Remark 2. When pr is even, the minimum order of an r -regular supergraph H will range between p and $p+d+1$. Thus the bound in the theorem is decreased by 1 in this case.

The strengthening of the theorem in the following statement is easily accomplished by a proof which we omit as it is entirely analogous.

Corollary. *Let G be a graph of order p with maximum degree d , and r be an integer such that $d \leq r \leq p-2$. Then G has an r -regular supergraph of order at most $p+r+1$ or $p+r+2$ if pr is even or odd, respectively. \square*

Jin Akiyama, Hiroshi Era and Frank Harary,
Nippon Ika University, Kawasaki, Japan; University of Michigan, Ann Arbor, USA

REFERENCES

- 1 P. Erdős and P. Kelly: The minimal regular graph containing a given graph. Am. Math. Monthly 70, 1074-1075 (1963).
- 2 P. Erdős and P. Kelly: The minimal regular graph containing a given graph. In: A Seminar on Graph Theory, p.65-69. F. Harary, ed. Holt, New York 1967.
- 3 F. Harary: Graph Theory. Addison-Wesley, Reading 1969.
- 4 D. König: Theorie der endlichen und unendlichen Graphen. Leipzig 1936, reprinted, Chelsea, New York 1950.

A remark on a paper by A. Grytczuk

In [3] Grytczuk showed that if $c_k(n)$ denotes the Ramanujan trigonometric sum, then

$$\sum_{d|k} |c_d(n)| = 2^{\omega(k/(k,n))} (k, n), \tag{1}$$

where $\omega(m)$ denotes the number of distinct prime divisors of m . In this note we prove a generalization of (1).

Let h be an arithmetic function and if m is a natural number, let

$$h_m(n) = \begin{cases} h(n) & \text{if } n|m \\ 0 & \text{else} \end{cases}$$

Define

$$H_m(n) = \sum_{d|(m,n)} \mu(m/d) h(d) = \sum_{d|m} \mu(m/d) h_n(d). \quad (2)$$

Note that $H_1(n) = h(1)$ and if $a \geq 1$, then

$$H_{p^a}(n) = \begin{cases} h_n(p^a) - h_n(p^{a-1}) & \text{if } p^a | n \\ h(p^a) - h(p^{a-1}) & \text{if } p^{a-1} || n \\ 0 & \text{if } p^{a-1} \nmid n \end{cases}$$

We now show that $H_m(n)$ is a multiplicative function of m if $h(m)$ is also. To do this it will suffice to show that $h_n(d)$ is a multiplicative function of d , since, by (2), $H_m(n)$ is then a convolution of two multiplicative functions. If $(m, n) = 1$ and k is a natural number, then $mn|k$ if and only if $m|k$ and $n|k$. Thus, by the definition of $h_k(mn)$, we have

$$\begin{aligned} h_k(mn) &= \begin{cases} h(mn) & \text{if } mn|k \\ 0 & \text{else} \end{cases} = \begin{cases} h(m)h(n) & \text{if } m|k \text{ and } n|k \\ 0 & \text{else} \end{cases} \\ &= \begin{cases} h_k(m)h_k(n) & . \end{cases} \end{aligned}$$

Thus $h_k(d)$ is a multiplicative function of d if h is multiplicative.

Theorem. *If h is a multiplicative function such that*

$$h(p^a) - h(p^{a-1}) \geq 0 \quad (3)$$

for all $a \geq 1$ and primes p , then for all positive integers k and m , we have

$$\sum_{d|k} |H_d(n)| = 2^{\omega(k/(k,n))} h((k, n)). \quad (4)$$

Proof: Since $H_d(n)$ is multiplicative so is $|H_d(n)|$ and thus so is the left hand side of (4) ([4], theorem 265). Since, for n fixed and $(k, l) = 1$, we have $(kl, n) = (k, n)(l, n)$ (see [5], p. 17) and since h is multiplicative we see that the right hand side of (4) is also a multiplicative function of k . To prove the theorem it therefore suffices to show that (4) holds when k is a prime power, $k = p^a$.

Note that, by (3), $h(p^a) \geq h(p^{a-1}) \geq \dots \geq h(1) \geq 0$, since h multiplicative implies $h(1) = 1$ or $h(1) = 0$.

Suppose $p^b \parallel n$. If $0 \leq b < a$, then

$$\begin{aligned} \sum_{j=0}^a |H_{p^j}(n)| &= h(1) + \sum_{j=1}^b (h(p^j) - h(p^{j-1})) + |H_{p^{b+1}}(n)| \\ &= h(1) + h(p^b) - h(1) + h(p^b) = 2h(p^b). \end{aligned} \tag{5}$$

If $b \geq a$, then

$$\sum_{j=0}^a |H_{p^j}(n)| = h(1) + \sum_{j=1}^a (h(p^j) - h(p^{j-1})) = h(p^a). \tag{6}$$

If we compare (5) and (6) with the right hand side of (4), in each of the two cases we see that they agree. This proves our theorem.

Examples:

1. Let r be a positive integer and

$$C_k^{(r)}(n) = \sum_{\substack{x_1, \dots, x_r \pmod k \\ (x_1, \dots, x_r, k) = 1}} \exp(2\pi i n (x_1 + \dots + x_r)/k).$$

In [2], theorem 1, Cohen proves that

$$C_k^{(r)}(n) = \sum_{d|(k,n)} d^r \mu(k/d).$$

Thus, if we take $h(n) = n^r$, we have, by the theorem,

$$\sum_{d|k} |C_d^{(r)}(n)| = 2^{\omega(k/(k,n))} (k,n)^r.$$

The case $r = 1$ is (1), the result of Grytczuk.

2. Let $h(n) = d(n)$, the divisor function. Then d satisfies the hypotheses of the theorem and in this case

$$H_{p^a}(n) = \begin{cases} 1 & \text{if } p^a | n \\ -a & \text{if } p^{a-1} \parallel n. \\ 0 & \text{if } p^{a-1} \nmid n \end{cases}$$

The theorem states that in this case

$$\sum_{d|k} |H_d(n)| = 2^{\omega(k/(k,n))} d((k,n)).$$

It might be interesting to note that the evaluation of the sum $\sum_{d|k} H_d(n)$ is much easier even if h is not known to be multiplicative. Indeed it follows immediately from the Möbius inversion formula ([4], theorem 266) that

$$H_m(n) = \sum_{d|(m,n)} \mu(m/d) h(d) = \sum_{d|m} \mu(m/d) h_n(d)$$

if and only if

$$\sum_{d|k} H_d(n) = h_n(k) = \begin{cases} h(k) & \text{if } k|n \\ 0 & \text{else} \end{cases}.$$

Finally we remark that it may be possible to generalize our result further to the class of functions considered by Anderson and Apostol in [1]. We hope to return to this in a later paper.

ACKNOWLEDGMENT

I would like to thank John Steinig for his suggestions for the improvement of the exposition of this paper.

Don Redmond, Southern Illinois University, Carbondale, USA

REFERENCES

- 1 D.R. Anderson and T.M. Apostol: The Evaluation of Ramanujan's Sums and Generalizations. *Duke Math. J.* 20, 211–216 (1953).
- 2 E. Cohen: Trigonometric Sums in Elementary Number Theory. *Am. Math. Monthly* 66, 105–117 (1959).
- 3 A. Grytczuk: An Identity Involving Ramanujan's Sum. *El. Math.* 36, 16–17 (1981).
- 4 G.H. Hardy and E.M. Wright: *An Introduction to the Theory of Numbers*, 5th ed. Oxford University Press, 1979.
- 5 W.J. LeVeque: *Topics in Number Theory*, vol. 1. Addison-Wesley, 1956.

© 1983 Birkhäuser Verlag, Basel

0013-6018/83/010017-04\$1.50 + 0.20/0

Nachtrag

P. Läuchli: Das Gitterspiel, *El. Math.* 37, 109–113 (1982).

Nach Drucklegung des Aufsatzes stellte ich aufgrund eines neuen Buches [2], dessen Manuskript mir freundlicherweise von den Autoren zur Verfügung gestellt wurde, fest, dass das «Gitterspiel» von C. Berge schon 1907 von Wythoff beschrieben wurde [3]. Ferner hat Coxeter [1] eine sehr elegante Verbindung zum goldenen Schnitt gezogen.

P. Läuchli

- 1 H.S.M. Coxeter: The golden section, phyllotaxis, and Wythoff's game. *Scr. Math.* 19, 135–143 (1953).
- 2 G. Schmidt und T. Ströhlein: *Relationen und Graphen*. Erscheint 1983.
- 3 W.A. Wythoff: A modification of the game of Nim, *Nieuw. Archf. Wisk.* (2), 7, 199–202 (1907).

© 1983 Birkhäuser Verlag, Basel

0013-6018/83/010020-01\$1.50 + 0.20/0