

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 39 (1984)
Heft: 5

Rubrik: Berichtigung

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

that f^2 and g have the same leading term. One thereby obtains distinct K -reciprocal representations, $1/h_1 + \dots = 1/f + \dots$; adding $1/X$ to (the left of) each produces distinct such representations with the same initial term.

The two representations given by the above recipe may, however, have unequal lengths. Consider, for instance, the choices $f = X^2 + 1$, $h_1 = X^2 - 1$, $r = X$. The recipe produces

$$\begin{aligned} \frac{X^4 + X^3 + 2X - 1}{X^5 + X^2 - X} &= \frac{1}{X} + \frac{1}{X^2 - 1} + \frac{1}{-X^5 + X^3 - X^2 + X + 1} + \frac{1}{p_{11}} \\ &= \frac{1}{X} + \frac{1}{X^2 + 1} + \text{sum of three additional terms.} \end{aligned} \tag{4}$$

Let w denote the rational function in (4), we compute via the first equation in (4) that

$$\begin{aligned} v = 1/X + w/X &= \frac{X^5 + X^4 + X^3 + X^2 + X - 1}{X^6 + X^3 - X^2} \\ &= \frac{1}{X} + \frac{1}{X^2} + \frac{1}{X^3 - X} + \frac{1}{-X^6 + X^4 - X^3 + X^2 + X} + \frac{1}{p_{12}}, \end{aligned}$$

which explains the origin of the first equation in (3). How does one obtain the second equation in (3)? Simply apply the Theorem's algorithm to w , multiply the result through termwise by $1/X$, and then add $1/X$ to (the left of) the ensuing expression. Finally, it is amusing to note that an application of the Theorem's algorithm directly to v produces yet another distinct \mathbf{Q} -reciprocal representation of v .

David E. Dobbs and Robert M. McConnel
University of Tennessee, Knoxville (USA)

REFERENCES

- 1 W.J. LeVeque: Fundamentals of number theory. Addison-Wesley, Reading 1977.
- 2 I. Niven and H.S. Zuckerman: An introduction to the theory of numbers, 3rd ed. Wiley, New York 1972.

© 1984 Birkhäuser Verlag, Basel

0013-6018/84/050126-04\$1.50 + 0.20/0

Berichtigung

**Correction to paper: Packing of 180 equal circles on a sphere.
Elemente der Mathematik. Vol.38, 1983, 119–122**

Professor H.S.M. Coxeter kindly drew my attention to the fact that figures of packings of 72 and 180 circles are chiral and not centro-symmetric. Namely, central symmetry can occur in tessellation $\{3, q + \}_{b,c}$ ($q = 4$ or 5) if $bc(b - c) = 0$, that is, the tessellation has a plane of symmetry. Thus, the statement in the last sentence of the paper is not valid.

Tibor Tarnai
Hungarian Institute for Building Science Budapest