Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 39 (1984)
Heft: 2

Artikel: A remark on buffon's needle problem
Autor: Abel, Ulrich
DOI: https://doi.org/10.5169/seals-38015

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A Remark on Buffon's Needle Problem

Let G_n be a grid in the Euclidean space $\mathbb{R}^n$ with coordinates $x_1, x_2, \ldots, x_n$ determined by hyperplanes parallel to the hyperplanes with the equation $x_i = 0$ separated by a distance of $2L$.
In his generalization of Buffon's Needle Problem to n dimensions Stoka [2] gives an expression for the probability that a segment ω of length L which will be "thrown" in a random fashion into the $\mathbb{R}^n$ cuts the grid G_n.
Let A_j be the event: the segment ω cuts a hyperplane of G_n parallel to the hyperplane with the equation $x_j = 0$. In [2] and [3] Stoka shows that the probability for the event A_j is

$$P(A_j) = \frac{\Gamma\left(\frac{n}{2}\right)}{(n-1)\,\Gamma\left(\frac{n-1}{2}\right)\sqrt{\pi}} \qquad (j = 1, 2, \ldots, n) \tag{1}$$

(formula (9) in [3] is a misprint) and finds for the estimator

$$\hat{P}_n = \frac{1}{nN} \sum_{j=1}^{n} (\text{number of times } A_j \text{ occurs in } N \text{ independent trials})$$

of $P(A_j)$ the variance

$$D^2(\hat{P}_n) = \frac{1}{nN}\,\theta_n \tag{2}$$

with

$$\theta_n = \frac{(2n-1)\,\Gamma\left(\frac{n}{2}\right)}{(n-1)\,\Gamma\left(\frac{n-1}{2}\right)\sqrt{\pi}} - \frac{n\left[\Gamma\left(\frac{n}{2}\right)\right]^2}{(n-1)^2\left[\Gamma\left(\frac{n-1}{2}\right)\right]^2\pi} - n + 1$$

$$+ \frac{2^{n-2}(n-1)\,\Gamma\left(\frac{n}{2}\right)}{\pi^{3/2}(n-2)!}\left[\left(\pi + \frac{1}{2n}\right)\Gamma\left(\frac{n-1}{2}\right) - \frac{2\sqrt{\pi}}{n-1}\,\Gamma\left(\frac{n}{2}\right)\right]$$

([3], formula (15)). On the other hand he calculates for the estimator $\hat{P}_1 = M^{-1}$ (number of times A_1 occurs in M independent trials) of $P(A_j)$ the variance

$$D^2(\hat{P}_1) = \frac{1}{M}\,\frac{\Gamma\left(\frac{n}{2}\right)}{(n-1)\,\Gamma\left(\frac{n-1}{2}\right)\sqrt{\pi}}\left[1 - \frac{\Gamma\left(\frac{n}{2}\right)}{(n-1)\,\Gamma\left(\frac{n-1}{2}\right)\sqrt{\pi}}\right] \tag{3}$$

([3], formula (16)). For the case $n=2$ compare Schuster [1]. If we set both variances (2) and (3) equal we receive

$$M=\sigma(n)\, n N \tag{4}$$

with

$$\sigma(n)=\frac{\Gamma\left(\frac{n}{2}\right)\left[(n-1)\,\Gamma\left(\frac{n-1}{2}\right)\sqrt{\pi}-\Gamma\left(\frac{n}{2}\right)\right]}{(n-1)^2\left[\Gamma\left(\frac{n-1}{2}\right)\right]^2\pi\,\theta_n}. \tag{5}$$

Thus, N independent trials in G_n give the same information about the probability (1) as $\sigma(n)\, n N$ independent trials in a grid determined by hyperplanes parallel to the hyperplane with the equation $x_1=0$. Schuster [1] finds $\sigma(2)\approx 1.1114$. Stoka [3] calculates $\sigma(3)\approx 1.1121$ and $\sigma(4)\approx 1.1039$ and conjectures that $\lim\limits_{n\to\infty}\sigma(n)=1$. In this note we give an asymptotic expression for $\sigma(n)$ and show that Stoka's conjecture is true.

Theorem. *It holds*

$$\sigma(n)=\frac{\pi\, n!-2^{n-2}\, n\left[\Gamma\left(\frac{n}{2}\right)\right]^2}{\pi\, n!-2^{n-2}\, n^2\left[\Gamma\left(\frac{n}{2}\right)\right]^2+2^{n-3}(n-1)^3\left[\Gamma\left(\frac{n-1}{2}\right)\right]^2}, \tag{6}$$

$$\sigma(n)\cong\frac{\sqrt{2\pi}-n^{-1/2}\left(\frac{n-2}{n}\right)^{n-1}e^2}{\sqrt{2\pi}-n^{1/2}\left(\frac{n-2}{n}\right)^{n-1}e^2+n^{1/2}\left(\frac{n-1}{n}\right)^3\left(\frac{n-3}{n}\right)^{n-2}e^3}\quad(n\to\infty) \tag{7}$$

and

$$\lim_{n\to\infty}\sigma(n)=1\,. \tag{8}$$

Proof: First we simplify the expression (5). For abbreviation we set $a=a_n=\Gamma\left(\frac{n-1}{2}\right)$ and $b=b_n=\Gamma\left(\frac{n}{2}\right)$. Then by (5)

$$\sigma(n)=\frac{(n-1)\sqrt{\pi}\,a\,b-b^2}{(n-1)^2\pi\left\{\frac{(2n-1)\,a\,b}{(n-1)\sqrt{\pi}}-\frac{n\,b^2}{(n-1)^2\pi}-(n-1)\,a^2+\right.}$$

$$\left.+\frac{2^{n-2}(n-1)\left(\pi+\frac{1}{2n}\right)a^3\,b}{\pi^{3/2}(n-2)!}-\frac{2^{n-1}\,a^2\,b^2}{\pi\,(n-2)!}\right\}.$$

The formula

$$\Gamma(x)\,\Gamma\left(x+\frac{1}{2}\right)=\frac{\sqrt{\pi}}{2^{2x-1}}\,\Gamma(2x)$$

gives us

$$a\,b=\frac{\sqrt{\pi}}{2^{n-2}}\,(n-2)!$$

and therefore

$$\sigma(n)=\frac{2^{-(n-2)}\,\pi\,(n-1)!-b^2}{2^{-(n-2)}\,\pi\,(n-1)!\,(2n-1)-n\,b^2+\dfrac{(n-1)^3}{2n}\,a^2-2^{-(n-3)}\,\pi\,(n-1)!\,(n-1)}$$

$$=\frac{\pi\,n!-2^{n-2}\,n\,b^2}{\pi\,n!-2^{n-2}\,n^2\,b^2+2^{n-3}\,(n-1)^3\,a^2}$$

which is (6).

By Stirling's formula for the Gamma-function it holds

$$\log\Gamma(x+1)=\left(x+\frac{1}{2}\right)\log x-x+c+0\left(\frac{1}{x}\right)\qquad(x\to+\infty)$$

with

$$c=\frac{1}{2}\log(2\pi)\,.$$

This implies

$$\sigma(n)\cong\frac{\pi\,n^{n+1/2}\,e^{-n+c}-2^{n-2}\,n\left(\dfrac{n-2}{2}\right)^{n-1}e^{-(n-2)+2c}}{\pi\,n^{n+1/2}\,e^{-n+c}-2^{n-2}\,n^2\left(\dfrac{n-2}{2}\right)^{n-1}e^{-(n-2)+2c}+2^{n-3}\,(n-1)^3\left(\dfrac{n-3}{2}\right)^{n-2}e^{-(n-3)+2c}}$$

$$=\frac{\sqrt{2\pi}-n^{-1/2}\left(\dfrac{n-2}{n}\right)^{n-1}e^2}{\sqrt{2\pi}-n^{1/2}\left(\dfrac{n-2}{n}\right)^{n-1}e^2+n^{1/2}\left(\dfrac{n-1}{n}\right)^3\left(\dfrac{n-3}{n}\right)^{n-2}e^3}$$

which is (7).

Evidently the numerator of (7) converges to $\sqrt{2\pi}$ for $n \to \infty$. It is sufficient to show that

$$d = \lim_{n \to \infty} n^{1/2} \left[\left(\frac{n-2}{n}\right)^{n-1} - \left(\frac{n-1}{n}\right)^3 \left(\frac{n-3}{n}\right)^{n-2} e \right] = 0 .$$

Now

$$|d| \leqq \lim_{n \to \infty} n^{1/2} \left(\frac{n-1}{n}\right)^3 \left(\frac{n-3}{n}\right)^{n-2}$$
$$\cdot \left\{ \left| \left(\frac{n}{n-1}\right)^3 \left(\frac{n-2}{n}\right) - 1 \right| \left(\frac{n-2}{n-3}\right)^{n-2} + \left| \left(\frac{n-2}{n-3}\right)^{n-2} - e \right| \right\} .$$

Because of

$$\left(\frac{n}{n-1}\right)^3 \frac{n-2}{n} - 1 = 0\left(\frac{1}{n}\right) \qquad (n \to \infty)$$

it remains to show that

$$\left(1 + \frac{1}{n}\right)^{n+1} = e + 0\left(\frac{1}{n}\right) \qquad (n \to \infty) .$$

To see this we set $x_i = i/n$ $(i = 0, 1, \ldots, n)$ and Taylor's formula gives us

$$e^{x_i} = \left(1 + \frac{1}{n}\right) e^{x_{i-1}} + \frac{1}{2n^2} e^{\xi_i} \qquad \text{with} \qquad x_{i-1} < \xi_i < x_i .$$

This implies

$$e = e^{x_n} = \left(1 + \frac{1}{n}\right)^n e^{x_0} + \frac{1}{2n^2} \sum_{i=1}^{n} \left(1 + \frac{1}{n}\right)^{n-i} e^{\xi_i}$$
$$= \left(1 + \frac{1}{n}\right)^n + 0\left(\frac{1}{n}\right) \qquad (n \to \infty) .$$

Thus the Theorem is proved.

Ulrich Abel
Mathematisches Institut der Justus-Liebig-Universität, Giessen

REFERENCES

1 E. F. Schuster: Buffon's needle experiment. Am. Math. Monthly *81*, 26–29 (1974).
2 M. I. Stoka: Une extension du problème de l'aiguille de Buffon dans l'espace euclidien R^n. Boll. U.M.I. (IV) *10*, 386–389 (1974).
3 M. I. Stoka: Quelques considérations concernant le problème de l'aiguille de Buffon dans l'espace euclidien E_n. El. Math. *38*, 4–11 (1983).

© 1984 Birkhäuser Verlag, Basel 0013-6018/84/020034-04$1.50 + 0.20/0