

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 42 (1987)
Heft: 1

Artikel: On a diophantine equation $(x^2 - 1)(y^2 - 1) = (z^2 - 1)^2$
Autor: Hirose, Shoichi
DOI: <https://doi.org/10.5169/seals-40025>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On a diophantine equation $(x^2 - 1)(y^2 - 1) = (z^2 - 1)^2$

1. Schinzel and Sierpiński [4] gave the complete solution of the equation

$$(x^2 - 1)(y^2 - 1) = \left(\left(\frac{y-x}{2} \right)^2 - 1 \right)^2 \quad (1)$$

in positive integers $x, y, x < y$. Equation (1) is a special case of the equation

$$(x^2 - 1)(y^2 - 1) = (z^2 - 1)^2. \quad (2)$$

The only known solution of (2) which is not a solution of (1) is $(x, y, z) = (4, 31, 11)$ found by Szymiczek [5].

In this paper we give a method for finding positive integer solutions x, y, z of (2), and show two new solutions.

2. Let $1 < x < y$, $x^2 - 1 = D v^2$, $y^2 - 1 = D' v'^2$, where D and D' have no square factors. Then we get $D = D'$, because

$$(x^2 - 1)(y^2 - 1) = D D' (v v')^2 = (z^2 - 1)^2.$$

If (u_1, v_1) is the minimal positive solution of the equation

$$u^2 - D v^2 = 1,$$

then all positive solutions are given by (u_k, v_k) for $k = 1, 2, 3, \dots$ where u_k and v_k are the integers defined by

$$u_k + v_k \sqrt{D} = (u_1 + v_1 \sqrt{D})^k. \quad (3)$$

Since we can put $x = u_i, v = v_i, y = u_j, v' = v_j, i < j$, we have

$$D v_i v_j + 1 = z^2. \quad (4)$$

From (3), it follows that

$$u_{k+1} = u_1 u_k + D v_1 v_k,$$

$$v_{k+1} = u_1 v_k + v_1 u_k.$$

Hence, by mathematical induction, $u_k = f_k(u_1)$ is a polynomial of degree k in u_1 , and $v_k/v_1 = g_k(u_1)$ is a polynomial of degree $k-1$ in u_1 . Indeed,

$$g_k(u_1) = \sum_{n=0}^{\left[\frac{k-1}{2}\right]} (-1)^n \binom{k-n-1}{n} (2u_1)^{k-2n-1}.$$

Thus, the left side of (4)

$$\begin{aligned} D v_i v_j + 1 &= D v_1 g_i(u_1) \cdot v_1 g_j(u_1) + 1 \\ &= (u_1^2 - 1) g_i(u_1) g_j(u_1) + 1 = F_{ij}(u_1) \end{aligned}$$

is a polynomial of degree $i+j$ in u_1 . If there is a solution (u_1, z) , $u_1 > 1$ of the equation

$$F_{ij}(u_1) = z^2,$$

then $(x, y, z) = (u_i, u_j, z)$ is a solution of (2).

(i) $i = 1, j = 2$. Since $v_2 = 2u_1 v_1$, we get $g_2(u_1) = 2u_1$, and so $F_{12}(u_1) = (u_1^2 - 1) \cdot 2u_1 + 1 = 2u_1^3 - 2u_1 + 1$. $F_{12}(u_1) = z^2$ has the following three solutions (u_1, z) , $1 < u_1 < 10^5$:

$$(u_1, z) = (3, 7), (4, 11), (155, 2729).$$

Therefore, we obtain the following three solutions of (2), the first of which is a solution of (1), and the last of which is new.

$$(x, y, z) = (3, 17, 7), (4, 31, 11), (155, 48049, 2729).$$

(ii) $i = 1, j = 4$. Since $v_4 = v_1(8u_1^3 - 4u_1)$, we get

$$F_{14}(u_1) = (u_1^2 - 1)(8u_1^3 - 4u_1) + 1 = 8u_1^5 - 12u_1^3 + 4u_1 + 1.$$

$F_{14}(u_1) = z^2$ has the following solution (u_1, z) , $1 < u_1 < 10^3$:

$$(u_1, z) = (2, 13).$$

Therefore, we obtain the following new solution of (2), where x is even, y and z are odd:

$$(x, y, z) = (2, 97, 13).$$

(iii) In the case of $3 \leq i+j \leq 9$, except above, we get only the following three solutions, all of which are solutions of (1).

i	j		u_1	x	y	z
2	3	$1 < u_1 < 10^3$	3	17	99	41
3	4	$1 < u_1 < 10^2$	3	99	577	239
4	5	$1 < u_1 < 35$	3	577	3363	1393

The equation (2) has no solutions other than above for all integers x and y , where $1 < x < y < 10^3$.

Though we cannot solve (2) completely, we conjecture that there are only three solutions of (2) which are not solutions of (1).

3. We applied our method to the equation

$$(x^2 + 1)(y^2 + 1) = (z^2 + 1)^2,$$

but we could not obtain other solutions than those Williams [6] found.

Our method cannot be applied to the equation

$$(x^2 - e)(y^2 - e) = (z^2 - e)^2$$

where $e \neq \pm 1, \pm 4$. However, our method may be applied to the equation

$$(x^2 - e)(y^2 - e) = \{h(z)\}^2$$

where $e = \pm 1, \pm 4$ and $h(z)$ is a linear or quadratic expression in z .

For example, the equations

$$(x^2 + 1)(y^2 + 1) = (z^2 + t^2)^2,$$

$$(x^2 - 1)(y^2 - 1) = (z^2 - t^2)^2$$

have always integer solutions for all integers t , given by

$$(x, y, z) = (t, 4t^3 + 3t, 2t^2 + 1)$$

$$(x, y, z) = (2t^2 \pm 1, 8t^4 \pm 8t^2 + 1, 4t^3 \pm 3t),$$

respectively.

Shoichi Hirose, Mita High School, Tokyo, Japan

REFERENCES

- 1 R. K. Guy: Unsolved problems in number theory. New York: Springer-Verlag, 1981, 105.
- 2 J. A. H. Hunter: Problem 5020. Amer. Math. Monthly 69, 316–317 (1962).
- 3 R. V. Iyer: Solution 5020. Amer. Math. Monthly 70, 574 (1963).
- 4 A. Schinzel and W. Sierpiński: Sur l'équation diophantienne $(x^2 - 1)(y^2 - 1) = [(y - x)/2]^2 - 1$.
El. Math. 18, 132–133 (1963).
- 5 K. Szymiczek: On a diophantine equation. El. Math. 22, 37–38 (1967).
- 6 H. C. Williams: Note on a diophantine equation. El. Math. 25, 123–125 (1970).