

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 42 (1987)
Heft: 1

Artikel: An elementary proof of the Theorem of Beckman and Quarles
Autor: Benz, Walter
DOI: <https://doi.org/10.5169/seals-40026>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

An elementary proof of the Theorem of Beckman and Quarles

1. I have been asked by colleagues to write down that proof of the fundamental and classical Theorem of Beckman, Quarles [1] that I have presented in a beginners course on Geometric Transformations for students already familiar with the basic methods of Linear Algebra. The proof in question, which is already sketched in a more general context in [2], is a mixture of ideas of Beckman, Quarles [1], Schröder [5], Benz [2] up to some new details. In this connection we also refer to Parhomenko and Modenov [4] and to their proof of the Theorem in question.

Let \mathbb{R}^n ($1 < n < \infty$) be equipped with the usual scalar product

$$a \cdot b := \sum_{i=1}^n \alpha_i \beta_i$$

for $a = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$ and $b = (\beta_1, \dots, \beta_n) \in \mathbb{R}^n$.
Then

$$\|a - b\| := \sqrt{(a - b)^2}$$

is called the distance of $a, b \in \mathbb{R}^n$.

Theorem of Beckman and Quarles: Suppose $k > 0$ to be a fixed real number and suppose f to be a mapping of \mathbb{R}^n ($1 < n < \infty$) into itself such that

$$\|p - q\| = k \text{ implies } \|f(p) - f(q)\| = k$$

for all $p, q \in \mathbb{R}^n$. Then f is an isometry of \mathbb{R}^n and hence a bijective linear mapping up to a translation.

In section 2 we shall collect some simple facts which are useful later on. Those elementary facts could be presented in a course far ahead the proof of the theorem in question, possibly in the form of exercises for the students.

The proof itself will be given in sections 3 and 4. It might be noticed that the original theorem in [1] was formulated for multivalued transformations f . This is however no substantial generalization as was pointed out in [3] in the case of Lorentz transformations of \mathbb{R}^n .

2. Throughout this note exactly the elements of \mathbb{R}^n ($1 < n < \infty$) are called points.

1) Suppose that a, m, b are points such that

$$\|m - a\| = \|b - m\| = \frac{1}{2} \|b - a\|.$$

Then $m = \frac{1}{2}(a + b)$.

Proof: Putting $q := \|m - a\|$, $a' := m - a$, $b' := b - m$ we have $(b - a)^2 + (a' - b')^2 = (a' + b')^2 + (a' - b')^2 = 4q^2$ and hence $(a' - b')^2 = 0$.

2) A set of n distinct points of \mathbb{R}^n which are pairwise of distance $\beta > 0$ will be called a β -set. Suppose that α, β are positive real numbers with

$$\gamma(\alpha, \beta) := 4\alpha^2 - 2\beta^2 \left(1 - \frac{1}{n}\right) > 0$$

and suppose that P is a β -set. Then there exist exactly two distinct points in \mathbb{R}^n which have distance α from all $p \in P$. Those two points will be called the α -associated points of P . Their distance is $\sqrt{\gamma(\alpha, \beta)}$.

Proof: a) Let $P = \{p_1, \dots, p_n\}$ be a β -set. Then for $i, j \in \{1, 2, \dots, n-1\}$ with $i \neq j$ we have

$$(p_i - p_n)(p_j - p_n) = \frac{1}{2}\beta^2,$$

because of $\beta^2 = (p_i - p_j)^2 = ((p_i - p_n) - (p_j - p_n))^2$. Define $\lambda_r := \frac{\beta}{\sqrt{2r(r+1)}}$ for $r = 1, 2, \dots$ and e_1, \dots, e_{n-1} by $(1+s)\lambda_s e_s := (p_s - p_n) - \sum_{r=1}^{s-1} \lambda_r e_r$ for $s = 1, \dots, n-1$. Obviously, $e_i^2 = 1$. We now prove

$$e_i e_j = \begin{cases} 1 & \text{for } i = j \leq n-1 \\ 0 & \text{for } i < j \leq n-1 \end{cases}$$

by induction along the sequence

$$(1, 1), (1, 2), (2, 2), (1, 3), (2, 3), (3, 3), \dots, (n-1, n-1) \quad \text{for } (i, j):$$

Step $(i, i) \rightarrow (1, i+1)$: Here we have

$$\begin{aligned} \frac{1}{2}\beta^2 &= (p_1 - p_n)(p_{i+1} - p_n) = 2\lambda_1 e_1 \left(\sum_{r=1}^i \lambda_r e_r + (2+i)\lambda_{i+1} e_{i+1} \right) \\ &= 2\lambda_1^2 + 2(2+i)\lambda_1 \lambda_{i+1} e_1 e_{i+1}, \end{aligned}$$

and hence $e_1 e_{i+1} = 0$, because of $\frac{1}{2}\beta^2 = 2\lambda_1^2$.

Step $(i-1, j) \rightarrow (i, j)$ in case $i < j$: Here we have

$$\begin{aligned} \frac{1}{2}\beta^2 &= (p_i - p_n)(p_j - p_n) = \left(\sum_{r=1}^{i-1} \lambda_r e_r + (1+i)\lambda_i e_i \right) \left(\sum_{r=1}^{j-1} \lambda_r e_r + (1+j)\lambda_j e_j \right) \\ &= \sum_{r=1}^{i-1} \lambda_r^2 + (1+i)\lambda_i^2 + (1+i)(1+j)\lambda_i \lambda_j e_i e_j, \end{aligned}$$

and hence $e_i e_j = 0$, because of $\frac{1}{2}\beta^2 = \sum_{r=1}^{i-1} \lambda_r^2 + (1+i)\lambda_i^2$ by observing

$$\lambda_r^2 = \frac{\beta^2}{2} \left(\frac{1}{r} - \frac{1}{r+1} \right).$$

Step $(i-1, i) \rightarrow (i, i)$: We finally have

$$\beta^2 = (p_i - p_n)^2 = \left(\sum_{r=1}^{i-1} \lambda_r e_r + (1+i) \lambda_i e_i \right)^2 = \sum_{r=1}^{i-1} \lambda_r^2 + (1+i)^2 \lambda_i^2 e_i^2,$$

and hence $e_i^2 = 1$.

b) Suppose now that $q \in \mathbb{R}^n$ has distance α from all $p_s \in P$. This implies

$$(q - p_n)(p_s - p_n) = \frac{1}{2} \beta^2 \quad \text{for all } s = 1, \dots, n-1,$$

because of $\alpha^2 = (q - p_s)^2 = ((q - p_n) - (p_s - p_n))^2$.

Put $q - p_n := \sum_{r=1}^n \mu_r e_r$, $\mu_r \in \mathbb{R}$, by extending $\{e_1, \dots, e_{n-1}\}$ of part a) to an orthonormal basis $\{e_1, \dots, e_n\}$ of \mathbb{R}^n . We get the equation

$$\frac{1}{2} \beta^2 = (q - p_n)(p_s - p_n) = \sum_{r=1}^{s-1} \mu_r \lambda_r + (1+s) \mu_s \lambda_s \quad \text{for } s = 1, \dots, n-1.$$

The case $s = 1$ leads to $\mu_1 = \lambda_1$, and having already $\mu_i = \lambda_i$ for $i \in \{1, \dots, s-1\}$, $s < n$, we also get $\mu_s = \lambda_s$ by comparing the equation above with

$$\frac{1}{2} \beta^2 = \sum_{r=1}^{s-1} \lambda_r^2 + (1+s) \lambda_s^2.$$

Hence $q - p_n = \sum_{r=1}^{n-1} \lambda_r e_r + \mu_n e_n$. Now $(q - p_n)^2 = \alpha^2$ leads to

$$\mu_n^2 = \alpha^2 - \sum_{r=1}^{n-1} \lambda_r^2 = \alpha^2 - \frac{\beta^2}{2} \left(1 - \frac{1}{n} \right) = \frac{1}{4} \gamma(\alpha, \beta).$$

There are exactly two solutions q , namely the points

$$q_i = p_n + \sum_{r=1}^{n-1} \lambda_r e_r \pm \frac{1}{2} \sqrt{\gamma(\alpha, \beta)} \cdot e_n, \quad i = 1, 2,$$

which are in fact of distance α from all $p \in P$. Obviously, $(q_1 - q_2)^2 = \gamma(\alpha, \beta)$.

3) Again suppose that α, β are positive real numbers with $\gamma(\alpha, \beta) > 0$. Let x, y be points of distance $\sqrt{\gamma(\alpha, \beta)}$. Then there exists a β -set P such that x, y are the α -associated points of P .

Proof: Define $e_n := \frac{y-x}{\sqrt{\gamma(\alpha, \beta)}}$ and extend $\{e_n\}$ to an orthonormal basis $\{e_1, \dots, e_n\}$ of \mathbb{R}^n . If p_n is an arbitrary point of \mathbb{R}^n , then $P = \{p_1, \dots, p_n\}$ with

$$p_s - p_n := \sum_{r=1}^{s-1} \lambda_r e_r + (1+s) \lambda_s e_s \quad \text{for } s = 1, \dots, n-1$$

is a β -set by using the earlier defined λ_r . If we now take the special point

$$p_n := \frac{x+y}{2} - \sum_{r=1}^{n-1} \lambda_r e_r,$$

then the α -associated points of P are given by (see part b) of 2))

$$q_i = p_n + \sum_{r=1}^{n-1} \lambda_r e_r + \frac{1}{2} \sqrt{\gamma(\alpha, \beta)} e_n = \frac{x+y}{2} \pm \frac{y-x}{2} = \begin{cases} y \\ x \end{cases}.$$

3. Proposition: Let $\varrho > 0$ be a fixed real number and let $N > 2$ be a fixed integer. Suppose that $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($1 < n < \infty$) is a mapping such that

$$\alpha) \quad \|x-y\| = \varrho \text{ implies } \|f(x)-f(y)\| \leq \varrho,$$

$$\beta) \quad \|x-y\| = N\varrho \text{ implies } \|f(x)-f(y)\| = N\varrho$$

for all $x, y \in \mathbb{R}^n$. Then $\|x-y\| = \|f(x)-f(y)\|$ holds true for all $x, y \in \mathbb{R}^n$.

Proof: a) Distances ϱ and 2ϱ are preserved under f : Having points x, y with $\|x-y\| = \varrho$ define $z := 2y-x$ and having points x, z with $\|x-z\| = 2\varrho$ define $y := \frac{1}{2}(x+z)$. Put $p_\lambda := x + \frac{\lambda}{2}(z-x)$ for $\lambda = 0, 1, \dots, N$. Observe $\|f(p_0)-f(p_N)\| = N\varrho$ and $\|f(p_\lambda)-f(p_{\lambda+1})\| \leq \varrho$ for $\lambda = 0, 1, \dots, N-1$ because of $\|p_0-p_N\| = N\varrho$ and $\|p_\lambda-p_{\lambda+1}\| = \varrho$. The triangle inequality yields

$$\begin{aligned} N\varrho = \|f(p_0)-f(p_N)\| &\leq \|f(p_0)-f(p_2)\| + \sum_{\lambda=2}^{N-1} \|f(p_\lambda)-f(p_{\lambda+1})\| \leq \\ &\leq \sum_{\lambda=0}^{N-1} \|f(p_\lambda)-f(p_{\lambda+1})\| \leq N\varrho \end{aligned}$$

and hence $\|f(p_\lambda)-f(p_{\lambda+1})\| = \varrho$ ($\lambda = 0, 1, \dots, N-1$) and

$$\|f(p_0)-f(p_2)\| = \|f(p_0)-f(p_1)\| + \|f(p_1)-f(p_2)\|.$$

Because of $p_0 = x, p_1 = y, p_2 = z$ we thus have

$$\|f(x)-f(z)\| = 2\varrho \quad \text{and} \quad \|f(x)-f(y)\| = \varrho.$$

b) Suppose that $\|x-y\| = \varrho$ for $x, y \in \mathbb{R}^n$. Then

$$f(x + \lambda(y-x)) = f(x) + \lambda(f(y)-f(x)) \tag{1}$$

holds true for all $\lambda = 0, 1, 2, \dots$: Put $p_\lambda := x + \lambda(y-x)$ for $\lambda = 0, 1, 2, \dots$ and observe

$$\|p_\lambda - p_{\lambda-1}\| = \varrho = \|p_{\lambda+1} - p_\lambda\| \quad \text{and} \quad \|p_{\lambda+1} - p_{\lambda-1}\| = 2\varrho$$

for $\lambda = 1, 2, \dots$. Since distances ϱ and 2ϱ are preserved we get

$$\varrho = \|f(p_\lambda) - f(p_{\lambda-1})\| = \|f(p_{\lambda+1}) - f(p_\lambda)\| = \frac{1}{2} \|f(p_{\lambda+1}) - f(p_{\lambda-1})\|$$

and hence (compare 1) in section 2) $f(p_\lambda) = \frac{1}{2}[f(p_{\lambda-1}) + f(p_{\lambda+1})]$. This proves (1) by induction since (1) is trivial in cases $\lambda = 0$ and $\lambda = 1$.

c) Let λ, μ be positive integers and suppose that $\|x - y\| = \frac{\lambda\varrho}{\mu}$ for $x, y \in \mathbb{R}^n$. Then $\|f(x) - f(y)\| = \frac{\lambda\varrho}{\mu}$ holds true: Because of $n > 1$ and $2\lambda\varrho > \|x - y\|$ there exists a point $z \in \mathbb{R}^n$ with $\|z - x\| = \lambda\varrho = \|z - y\|$. With such a fixed z define a, b by

$$x = z + \lambda(a - z), \quad y = z + \lambda(b - z) \quad (2)$$

and put

$$x' := z + \mu(a - z), \quad y' := z + \mu(b - z). \quad (3)$$

Since $\|a - z\| = \varrho = \|b - z\|$ we hence have the corresponding formulas to (2), (3) for the images because of b). Now

$$\|x' - y'\| = \varrho = \|f(x') - f(y')\| = \mu \|f(a) - f(b)\|$$

and

$$\|f(x) - f(y)\| = \lambda \|f(a) - f(b)\| \quad \text{imply} \quad \|f(x) - f(y)\| = \frac{\lambda\varrho}{\mu}.$$

d) Let r, s be positive rational numbers and let x, y be points such that $r\varrho < \|x - y\| < s\varrho$. Then $r\varrho \leq \|f(x) - f(y)\| \leq s\varrho$: Since $n > 1$ and $s\varrho > \|x - y\|$ there exists a point z with $\|z - x\| = \frac{s\varrho}{2} = \|z - y\|$. Now c) implies $\|f(z) - f(x)\| = \frac{s\varrho}{2} = \|f(z) - f(y)\|$ and hence $\|f(x) - f(y)\| \leq \|f(x) - f(z)\| + \|f(z) - f(y)\| = s\varrho$.

Put $w := x + \frac{s\varrho}{\|x - y\|}(y - x)$ and observe $\|w - x\| = s\varrho$ and

$$\|w - y\| = \left(\frac{s\varrho}{\|x - y\|} - 1 \right) \|y - x\| = s\varrho - \|y - x\| < (s - r)\varrho.$$

Hence $\|f(w) - f(x)\| = s\varrho$ by c) and $\|f(w) - f(y)\| \leq (s - r)\varrho$ by the already proved part of d). This implies

$$\|f(x) - f(y)\| \geq \|f(x) - f(w)\| - \|f(y) - f(w)\| \geq s\varrho - (s - r)\varrho = r\varrho.$$

4. Throughout this section let $k > 0$ be a fixed real number and f be a mapping of \mathbb{R}^n ($1 < n < \infty$) into itself such that distance k is preserved under f , i.e. $\|x - y\| = k$ implies $\|f(x) - f(y)\| = k$ for all $x, y \in \mathbb{R}^n$.

Lemma: Suppose that α, β are positive real numbers such that $\gamma(\alpha, \beta) > 0$ (compare section 2). Suppose moreover that f preserves distances α and β and that x, y are points with $\|x - y\| = \varepsilon := \sqrt{\gamma(\alpha, \beta)}$. Then $\|f(x) - f(y)\| \in \{0, \varepsilon\}$ and in case $2\varepsilon > \alpha$ we even have $\|f(x) - f(y)\| = \varepsilon$.

Proof: This is trivial for $\varepsilon = \alpha$ since distance α is preserved. So assume $\varepsilon \neq \alpha$. Let P be a β -set such that x, y are the α -associated points of P (compare 3) of section 2). It is $P' := f(P)$ also a β -set since distance β is preserved. If we denote the α -associated points of P' by x', y' we get $f(x), f(y) \in \{x', y'\}$ since distance α is also preserved under f and since the α -associated points of P' are uniquely determined. This implies $\|f(x) - f(y)\| \in \{0, \|x' - y'\|\} = \{0, \varepsilon\}$ according to 2) in section 2. Assume now $2\varepsilon > \alpha$. We have to show that $f(x) \neq f(y)$. Assume $f(x) = f(y)$ and take a $z \in \mathbb{R}^n$ with $\|z - x\| = \varepsilon$ and $\|y - z\| = \alpha$ which exists since $n > 1$ and $2\varepsilon > \alpha$. The already proved part of the lemma yields $\|f(x) - f(z)\| \in \{0, \varepsilon\}$, i.e. $\|f(y) - f(z)\| \in \{0, \varepsilon\}$ because of $f(x) = f(y)$. Hence $\alpha = \|y - z\| = \|f(y) - f(z)\| \in \{0, \varepsilon\}$. This contradicts $\varepsilon \neq \alpha > 0$.

We note the following three consequences of our Lemma:

- a) Putting $\alpha = k = \beta$ we realize that distance $\sqrt{\gamma(\alpha, \beta)} = k \sqrt{2 \left(1 + \frac{1}{n}\right)}$ is preserved.
- b) Putting $\alpha = \beta = k \sqrt{2 \left(1 + \frac{1}{n}\right)}$ we realize that distance $\sqrt{\gamma(\alpha, \beta)} = (n+1) \cdot \frac{2k}{n}$ is preserved.
- c) Put $\alpha = k$ and $\beta = k \sqrt{2 \left(1 + \frac{1}{n}\right)}$. Then $\|x - y\| = \sqrt{\gamma(\alpha, \beta)} = \frac{2k}{n}$ implies $\|f(x) - f(y)\| \in \left\{0, \frac{2k}{n}\right\}$, i.e. $\|f(x) - f(y)\| \leq \frac{2k}{n}$ for all $x, y \in \mathbb{R}^n$.

If we now take $\varrho := \frac{2k}{n}$ in the Proposition of section 3 and $N := n+1$ we realize that f is an isometry according to c), b) and $n > 1$.

Walter Benz, Math. Seminar, Universität Hamburg

REFERENCES

- 1 F. S. Beckman and D. A. Quarles, Jr.: On isometries of Euclidean spaces. Proc. Amer. Math. Soc. **4**, 810–815 (1953).
- 2 W. Benz: Isometrien in normierten Räumen. Aeq. Math. **29**, 204–209 (1985)
- 3 W. Benz: Eine Beckman-Quarles-Charakterisierung der Lorentztransformationen des \mathbb{R}^n . Archiv Math. **34**, 550–559 (1980).
- 4 A. S. Parhomenko and P. S. Modenov: Geometric Transformations I, II. Academic Press 1965.
- 5 E. M. Schröder: Eine Ergänzung zum Satz von Beckman und Quarles. Aeq. Math. **19**, 89–92 (1979).