

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 42 (1987)
Heft: 3: Archimedes was right. Part one

Rubrik: Kleine Mitteilung

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Kleine Mitteilung

Note on the diophantine equation $1 + x + x^2 + \dots + x^n = y^m$

This equation has occurred from time to time in the literature. In this note, we shall treat some aspects of the lowest cases $m=2$ and $m=3$. Throughout we shall assume $|x| > 1$ and $n > 1$.

Theorem. *The Diophantine equation $1 + x + \dots + x^n = y^2$ has solutions only for $n=3$ and $n=4$, the solutions being $(7, \pm 20)$ and $(3, \pm 11)$, respectively. The Diophantine equation $1 + x + \dots + x^n = y^3$ has solutions only for $n=2$ unless n is of the form $6k+4$. The respective solutions are $(18, 7)$ and $(-19, 7)$.*

This was shown by W. Ljunggren [4] in 1943*, using results by D. Schepel [7] and T. Nagell [6]. Actually, the case $n=3$ for the first equation follows from a result already known to Fermat and subsequently proved by E. Lucas and A. Genocchi [2]. (See also [1], vol. II, p. 487.)

Considering only primes for x , the sum $1 + x + \dots + x^n$ can be interpreted as $\sigma(p^n)$, the sum of the divisors of p^n . The question for $m=2$ now reads: can $\sigma(p^n)$ ever be a square? It is easily seen that for $n=1$, $1 + p = y^2$ if and only if $p=3$, $y=2$. Hence the answer to our question is simply, that $\sigma(p^n)$ is a square only in the cases $p=3$, $n=1$ or 4, and $p=7$, $n=3$.

Recently, Takaku [8] has shown that $\sigma(p^n)$ a square requires p to be less than $2^{2^{n+1}}$. In view of Ljunggren's theorem this is quite staggering.

Returning to the second part of the theorem, let us remark that R. Guy ([3], p. 7) poses the following question on so-called repunits, i.e. numbers $1111\dots 11 = 1 + 10 + \dots + 10^n$: Can such a number ever be a cube? (It is known that, except 1, they can never be squares.) As $1 + 10$ is not a cube, a positive answer to Guy's question can only stem from $n = 6k + 4$. Using the elementary geometric summation formula, one obtains $10^{6k+6} - 10 = 90y^3$, which gives $y^3 \equiv 2 \pmod{7}$ and hence is impossible. Therefore, the answer to Guy's question is «no».

A. Rotkiewicz, Mathematics Institute PAN, Warszawa

REFERENCES

- 1 L. E. Dickson: History of the theory of numbers, New York 1952.
- 2 A. Genocchi: Nouvelles Annales de Math. (2), t. 2 (1885), p. 306.
- 3 Richard K. Guy: Unsolved problems in number theory. New York: Springer-Verlag 1981.
- 4 W. Ljunggren: Noen setninger om ubestemte likninger av formen $\frac{x^n-1}{x-1} = y^q$, Norsk Mat. Tidsskr. 1 Hefte, 25 (1943), pp. 17–20.
- 5 K. Mahler: Über den grössten Primteiler spezieller Polynome zweiten Grades. Archiv for Math. og Naturv. B. XLI. Nr. 6 (1935).
- 6 T. Nagell: Sur l'équation indéterminée $\frac{x^n-1}{x-1} = y^2$, Norsk. Mat. Forenings Skrifter I, No. 3 (1921), 17 pp.
- 7 D. Schepel: On the Pell equation (Dutch), Nieuw Arch. Wiskunde 18 (1935), p. 1–30.
- 8 A. Takaku: Prime numbers such that the sums of the divisors of their powers are perfect squares. Colloq. Math. 49, 117–121 (1984).

1* The author wishes to thank Prof. J. Brzezinski for a copy of Ljunggren's paper which was not available in Poland.