

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 42 (1987)  
**Heft:** 5

**Artikel:** Some inequalities for the triangle  
**Autor:** Miloševi, M.  
**DOI:** <https://doi.org/10.5169/seals-40040>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

- 4 D. H. Lehmer: Coloring the platonic solids. Amer. Math. Monthly 93, 288–292 (1986).  
 5 K. Wagner: Graphentheorie. BI Mannheim, 1970.  
 6 J. M. Wills: Mathematik und Kunst: Die platonischen Körper. Siegener Studien (Liter. und Naturwiss.) 35, 17–27 (1983/84).

## Some inequalities for the triangle

**Notation.**  $a, b, c$  – sides BC, CA, AB of a triangle ABC;  $\alpha, \beta, \gamma$  – its angles;  $s$  – semi-perimeter;  $F$  – area;  $R$  – radius of circumcircle;  $r$  – radius of incircle;  $h_a, h_b, h_c$  – altitudes;  $m_a, m_b, m_c$  – medians and  $r_a, r_b, r_c$  – radii of excircles.

**Theorem 1.** *In every triangle the following equalities are valid:*

$$\frac{h_b + h_c}{r_a} + \frac{h_c + h_a}{r_b} + \frac{h_a + h_b}{r_c} = 6 \quad (1)$$

$$\cos^3 \frac{\alpha}{2} \cos \frac{\beta - \gamma}{2} + \cos^3 \frac{\beta}{2} \cos \frac{\gamma - \alpha}{2} + \cos^3 \frac{\gamma}{2} \cos \frac{\alpha - \beta}{2} = \frac{3s}{4R}. \quad (2)$$

**Proof.** Since

$$h_a = \frac{2F}{a}, \quad h_b = \frac{2F}{b}, \quad h_c = \frac{2F}{c} \quad \text{and}$$

$$r_a = \frac{F}{s-a}, \quad r_b = \frac{F}{s-b}, \quad r_c = \frac{F}{s-c},$$

we have

$$\begin{aligned} & \frac{h_b + h_c}{r_a} + \frac{h_c + h_a}{r_b} + \frac{h_a + h_b}{r_c} \\ &= \frac{2}{abc} (a(s-a)(b+c) + b(s-b)(c+a) + c(s-c)(a+b)) \\ &= \frac{2}{abc} (a^3 + b^3 + c^3 - 2s(a^2 + b^2 + c^2) + 2s(ab + bc + ca)). \end{aligned}$$

From  $abc = 4Rrs$ ,  $ab + bc + ca = r^2 + s^2 + 4Rr$ ,  $a^2 + b^2 + c^2 = 2(s^2 - 4Rr - r^2)$  and  $a^3 + b^3 + c^3 = 2s(s^2 - 6Rr - 3r^2)$ , we get

$$\begin{aligned} & \frac{h_b + h_c}{r_a} + \frac{h_c + h_a}{r_b} + \frac{h_a + h_b}{r_c} \\ &= \frac{1}{2Rrs} (2s(r^2 + s^2 + 4Rr) - 2s \cdot 2(s^2 - 4Rr - r^2) + 2s(s^2 - 6Rr - 3r^2)) = 6, \end{aligned}$$

i.e. (1) holds.

Moreover, since

$$\begin{aligned} h_b + h_c &= 8R \sin \frac{\alpha}{2} \cos^2 \frac{\alpha}{2} \cos \frac{\beta-\gamma}{2}, \quad h_c + h_a = 8R \sin \frac{\beta}{2} \cos^2 \frac{\beta}{2} \cos \frac{\gamma-\alpha}{2}, \\ h_a + h_b &= 8R \sin \frac{\gamma}{2} \cos^2 \frac{\gamma}{2} \cos \frac{\alpha-\beta}{2} \quad \text{and} \\ r_a &= s \cdot \tan \frac{\alpha}{2}, \quad r_b = s \cdot \tan \frac{\beta}{2}, \quad r_c = s \cdot \tan \frac{\gamma}{2}, \end{aligned}$$

we obtain

$$\begin{aligned} & \frac{h_b + h_c}{r_a} + \frac{h_c + h_a}{r_b} + \frac{h_a + h_b}{r_c} \\ &= \frac{8R}{s} \left( \cos^3 \frac{\alpha}{2} \cos \frac{\beta-\gamma}{2} + \cos^3 \frac{\beta}{2} \cos \frac{\gamma-\alpha}{2} + \cos^3 \frac{\gamma}{2} \cos \frac{\alpha-\beta}{2} \right), \end{aligned}$$

which is equivalent to

$$\cos^3 \frac{\alpha}{2} \cos \frac{\beta-\gamma}{2} + \cos^3 \frac{\beta}{2} \cos \frac{\gamma-\alpha}{2} + \cos^3 \frac{\gamma}{2} \cos \frac{\alpha-\beta}{2} = \frac{3}{4} \cdot \frac{s}{R},$$

i.e., (2).

**Corollary 1.** *In every triangle*

$$3 \cdot 2^n \leq \left( \frac{h_b + h_c}{r_a} \right)^n + \left( \frac{h_c + h_a}{r_b} \right)^n + \left( \frac{h_a + h_b}{r_c} \right)^n \leq 6^n, \quad n \geq 1. \quad (\text{A})$$

*Equality in (A) holds if and only if the triangle is equilateral and  $n = 1$ .*

**Proof.** The inequality (A) follows from (1) putting  $x_1 = \frac{h_b + h_c}{r_a}$ ,  $x_2 = \frac{h_c + h_a}{r_b}$  and  $x_3 = \frac{h_a + h_b}{r_c}$  in ([7], 47)

$$3 \left( \sum_{i=1}^3 \frac{x_i}{3} \right)^n \leq \sum_{i=1}^3 x_i^n \leq \left( \sum_{i=1}^3 x_i \right)^n, \quad (n \geq 1, x_i > 0).$$

**Remark 1.** a) The left part of (A) holds for  $n \leq 0$ . b) If  $0 < n < 1$ , then the inequality (A) is reversed.

**Remark 2.** For  $n = -1$ , we have the inequality 6.26 in [3]:

$$\frac{r_a}{h_b + h_c} + \frac{r_b}{h_c + h_a} + \frac{r_c}{h_a + h_b} \geq \frac{3}{2}.$$

**Corollary 2.** *The following inequality*

$$\cos^3 \frac{\alpha}{2} \cos \frac{\beta - \gamma}{2} + \cos^3 \frac{\beta}{2} \cos \frac{\gamma - \alpha}{2} + \cos^3 \frac{\gamma}{2} \cos \frac{\alpha - \beta}{2} \leq \frac{9\sqrt[3]{3}}{8}$$

*holds, with inequality only for  $\alpha = \beta = \gamma$ .*

**Proof.** This follows from (2) and  $2s = a + b + c \leq 3R\sqrt{3}$  ([3], 5.3).

**Theorem 2.** *In every triangle*

$$\left( \frac{m_a}{r_a} \right)^n + \left( \frac{m_b}{r_b} \right)^n + \left( \frac{m_c}{r_c} \right)^n \geq 3^{1-3n/2} \cdot \left( \frac{s}{r} \right)^n \geq 3, \quad n \geq 2/3, \quad (\text{B})$$

*and the equality is true if and only if the triangle is equilateral.*

**Proof.** Since ([5], 156)

$$m_a \geq \sqrt{s(s-a)}, \quad m_b \geq \sqrt{s(s-b)}, \quad m_c \geq \sqrt{s(s-c)},$$

we have

$$\left( \frac{m_a}{r_a} \right)^n + \left( \frac{m_b}{r_b} \right)^n + \left( \frac{m_c}{r_c} \right)^n \geq \frac{1}{r^n s^{n/2}} ((s-a)^{3n/2} + (s-b)^{3n/2} + (s-c)^{3n/2}).$$

Let us consider the function

$$f(x) = (s-x)^{3n/2}, \quad (0 < x < s, n \geq 2/3)$$

and its second derivative

$$f''(x) = \frac{3n}{4} (3n-2) (s-x)^{\frac{3n}{2}-2} \geq 0.$$

Hence, the function  $f$  is convex, so that

$$\sum_{i=1}^3 (s-x_i)^{3n/2} \geq 3^{1-3n/2} \left( 3s - \sum_{i=1}^3 x_i \right)^{3n/2}, \quad n \geq \frac{2}{3}. \quad (4)$$

Putting  $x_1 = a$ ,  $x_2 = b$  and  $x_3 = c$  in (4), we get

$$(s-a)^{3n/2} + (s-b)^{3n/2} + (s-c)^{3n/2} \geq 3^{1-3n/2} \cdot s^{3n/2}. \quad (5)$$

Now (5) and (3) imply that

$$\left( \frac{m_a}{r_a} \right)^n + \left( \frac{m_b}{r_b} \right)^n + \left( \frac{m_c}{r_c} \right)^n \geq 3^{1-3n/2} \left( \frac{s}{r} \right)^n, \quad n \geq \frac{2}{3}. \quad (6)$$

From (6) and  $s^2 \geq 27r^2$  ([3], 5.11), we obtain:

$$\left( \frac{m_a}{r_a} \right)^n + \left( \frac{m_b}{r_b} \right)^n + \left( \frac{m_c}{r_c} \right)^n \geq 3.$$

**Theorem 3.** *In every triangle*

$$4\sqrt[3]{3}(3R-4r) \leq \frac{a^3}{r_b r_c} + \frac{b^3}{r_c r_a} + \frac{c^3}{r_a r_b} \leq \frac{2R\sqrt[3]{3}}{r} (3R-4r). \quad (\text{C})$$

*Equality occurs if and only if the triangle is equilateral.*

**Proof.** We have

$$\begin{aligned} & \frac{a^3}{r_b r_c} + \frac{b^3}{r_c r_a} + \frac{c^3}{r_a r_b} \\ &= \frac{1}{F^2} (a^3(s-b)(s-c) + b^3(s-c)(s-a) + c^3(s-a)(s-b)) \\ &= \frac{1}{F^2} (s(a^4+b^4+c^4) - s^2(a^3+b^3+c^3) + abc(a^2+b^2+c^2)). \end{aligned}$$

From  $F = rs$ ,  $abc = 4Rrs$ ,  $a^2+b^2+c^2 = 2 \cdot (s^2 - 4Rr - r^2)$ ,  $a^3+b^3+c^3 = 2s(s^2 - 6Rr - 3r^2)$  and  $a^4+b^4+c^4 = 2(s^2 - 4Rr - r^2)^2 - 2(2rs)^2$ , we obtain

$$\begin{aligned} & \frac{a^3}{r_b r_c} + \frac{b^3}{r_c r_a} + \frac{c^3}{r_a r_b} = \frac{2}{rs} (4Rr^2 + 2Rs^2 - 3rs^2 + r^3) \\ &= \frac{2}{F} (r \cdot r(4R+r) + s \cdot s(2R-3r)). \end{aligned} \quad (7)$$

Then, since ([3], 7.2, 5.11)

$$r(4R+r) \geq F\sqrt{3}$$

and

$$s^2 \geq 27r^2,$$

(7) implies

$$\frac{a^3}{r_b r_c} + \frac{b^3}{r_c r_a} + \frac{c^3}{r_a r_b} \geq \frac{2}{F} (r \cdot F\sqrt{3} + s \cdot 3r\sqrt{3}(2R - 3r)) = 4\sqrt{3}(3R - 4r),$$

i.e. the first part of the inequality (C).

By 5.3 and 5.6 in [3] and (7) we get the second part of (C).

**Theorem 4.** *In every triangle*

$$\frac{4\sqrt{3}}{3} \cdot \frac{7R - 2r}{9R - 2r} \leq \frac{a}{h_b + h_c} + \frac{b}{h_c + h_a} + \frac{c}{h_a + h_b} \leq \sqrt{\frac{3R}{2r}}. \quad (\text{D})$$

*Equality holds if and only if the triangle is equilateral.*

**Proof.** Since

$$h_a = \frac{2F}{a}, \quad h_b = \frac{2F}{b}, \quad h_c = \frac{2F}{c},$$

we have

$$\begin{aligned} \frac{a}{h_b + h_c} + \frac{b}{h_c + h_a} + \frac{c}{h_a + h_b} &= \frac{abc}{2F} \left( \frac{1}{b+c} + \frac{1}{c+a} + \frac{1}{a+b} \right) \\ &= \frac{abc}{2F} \cdot \frac{(a^2 + b^2 + c^2 + 3(ab + bc + ca))}{((a+b+c)(ab + bc + ca) - abc)}. \end{aligned}$$

From  $abc = 4Rrs$ ,  $F = rs$ ,  $a+b+c = 2s$ ,  $ab+bc+ca = r^2 + s^2 + 4Rr$  and  $a^2 + b^2 + c^2 = 2(s^2 - 4Rr - r^2)$ , we get

$$\begin{aligned} \frac{a}{h_b + h_c} + \frac{b}{h_c + h_a} + \frac{c}{h_a + h_b} &= \frac{R}{s} \cdot \frac{5s^2 + 4Rr + r^2}{s^2 + 2Rr + r^2} \\ &= \frac{R}{s} \left( 5 - \frac{2r(3R + 2r)}{s^2 + 2Rr + r^2} \right). \end{aligned}$$

Using ([3], 5.8, 5.3)

$$s^2 \geq r(16R - 5r)$$

and

$$2s = a + b + c \leq 3R\sqrt{3},$$

we obtain

$$\begin{aligned} \frac{a}{h_b + h_c} + \frac{b}{h_c + h_a} + \frac{c}{h_a + h_b} &\geq \frac{R}{s} \left( 5 - \frac{3R + 2r}{9R - 2r} \right) \\ &= \frac{R}{s} \cdot \frac{6(7R - 2r)}{9R - 2r} \\ &\geq \frac{4\sqrt{3}}{3} \cdot \frac{7R - 2r}{9R - 2r}, \end{aligned}$$

i.e. the left part of the inequality (D).

From the inequalities

$$h_b + h_c \geq 2\sqrt{h_b h_c}, \quad h_c + h_a \geq 2\sqrt{h_c h_a}, \quad h_a + h_b \geq 2\sqrt{h_a h_b},$$

it follows that

$$\begin{aligned} \frac{a}{h_b + h_c} + \frac{b}{h_c + h_a} + \frac{c}{h_a + h_b} &= 2F \left( \frac{1}{h_a(h_b + h_c)} + \frac{1}{h_b(h_c + h_a)} + \frac{1}{h_c(h_a + h_b)} \right) \\ &\leq \frac{F}{\sqrt{h_a h_b h_c}} \left( \frac{1}{\sqrt{h_a}} + \frac{1}{\sqrt{h_b}} + \frac{1}{\sqrt{h_c}} \right). \end{aligned} \quad (8)$$

If in the well-known inequality

$$(x + y + z)^2 \leq 3(x^2 + y^2 + z^2), \quad (x, y, z > 0), \quad (9)$$

we put  $x = \frac{1}{\sqrt{h_a}}$ ,  $y = \frac{1}{\sqrt{h_b}}$  and  $z = \frac{1}{\sqrt{h_c}}$ , we obtain

$$\left( \frac{1}{\sqrt{h_a}} + \frac{1}{\sqrt{h_b}} + \frac{1}{\sqrt{h_c}} \right)^2 \leq 3 \left( \frac{1}{h_a} + \frac{1}{h_b} + \frac{1}{h_c} \right). \quad (10)$$

From  $\frac{1}{h_a} + \frac{1}{h_b} + \frac{1}{h_c} = \frac{1}{r}$ ,  $h_a h_b h_c = \frac{2F^2}{R}$ , (8) and (10), we get

$$\frac{a}{h_b + h_c} + \frac{b}{h_c + h_a} + \frac{c}{h_a + h_b} \leq \frac{F}{\sqrt{h_a h_b h_c}} \sqrt{\frac{3}{r}} = \sqrt{\frac{3R}{2r}}.$$

This proves the right part of (D).

**Remark 4.** From (D) using the inequality 5.1 in [3] we have the inequality

$$\sqrt{3} \leq \frac{a}{h_b + h_c} + \frac{b}{h_c + h_a} + \frac{c}{h_a + h_b} \leq \frac{R}{2r} \sqrt{3},$$

shown in [4].

**Theorem 5.** *In every triangle*

$$\frac{m_a^2}{bc} + \frac{m_b^2}{ca} + \frac{m_c^2}{ab} \leq \frac{5}{4} + \frac{R}{2r}, \quad (\text{E})$$

*with equality holding if and only if the triangle is equilateral.*

**Proof.** Using Bager's identity ([2], 39)

$$am_a^2 + bm_b^2 + cm_c^2 = \frac{s}{2} (s^2 + 2Rr + 5r^2),$$

we get

$$\begin{aligned} \frac{m_a^2}{bc} + \frac{m_b^2}{ca} + \frac{m_c^2}{ab} &= \frac{1}{abc} (am_a^2 + bm_b^2 + cm_c^2) \\ &= \frac{s}{2abc} (s^2 + 2Rr + 5r^2). \end{aligned}$$

Then, since ([3], 5.8, 5.1)

$$s^2 \leq 4R^2 + 4Rr + 3r^2$$

and

$$2r \leq R,$$

we have

$$\begin{aligned} \frac{m_a^2}{bc} + \frac{m_b^2}{ca} + \frac{m_c^2}{ab} &\leq \frac{1}{8Rr} (4R^2 + 6Rr + 8r^2) \\ &\leq \frac{1}{8Rr} \left( 4R^2 + 6Rr + 8r \cdot \frac{R}{2} \right) \\ &= \frac{5}{4} + \frac{R}{2r}, \end{aligned}$$

i.e. (E) holds.

**Theorem 6.** *For acute triangle the following inequality holds*

$$a^2 \tan \alpha + b^2 \tan \beta + c^2 \tan \gamma \geq \frac{4rs^3}{2R^2 + r^2}. \quad (\text{F})$$

**Proof.** If in Jensen's inequality for a convex function

$$f\left(\frac{\sum_{i=1}^3 p_i}{\sum_{i=1}^3 x_i}\right) \leq \frac{\sum_{i=1}^3 x_i f\left(\frac{p_i}{x_i}\right)}{\sum_{i=1}^3 x_i}, \quad p_i \text{ and } \frac{p_i}{x_i} \in \bar{I},$$

we put

$$x_1 = a, \quad x_2 = b, \quad x_3 = c, \quad p_1 = a^2, \quad p_2 = b^2 \quad \text{and} \quad p_3 = c^2,$$

then, for  $f(x) = 1/x$ , from ([1], 43; [3] 5.14)

$$a^2 + b^2 + c^2 = 4F(\cot \alpha + \cot \beta + \cot \gamma)$$

and

$$a^2 + b^2 + c^2 \leq 8R^2 + 4r^2,$$

we obtain (F).

**Remark 5.** This method shown in [6].

**Theorem 7.** *In every triangle*

$$3\sqrt[3]{3r} \leq \sqrt{h_a} + \sqrt{h_b} + \sqrt{h_c} \leq \left(1 + \frac{r}{R}\right)\sqrt{6R}. \quad (\text{G})$$

*Equality holds if and only if the triangle is equilateral.*

**Proof.** By means of the arithmetic-geometric inequality, we get

$$\begin{aligned} \sqrt{h_a} + \sqrt{h_b} + \sqrt{h_c} &\geq 3\sqrt[6]{\frac{2F}{a} \cdot \frac{2F}{b} \cdot \frac{2F}{c}} \\ &= 3\sqrt[6]{\frac{2r^2s^2}{R}}. \end{aligned} \quad (11)$$

Then, since ([3], 5.12)

$$2s^2 \geq 27Rr,$$

(11) implies

$$\sqrt{h_a} + \sqrt{h_b} + \sqrt{h_c} \geq 3\sqrt{3r},$$

i.e. the left part of the inequality (G).

If in the inequality (9) we put  $x = \frac{1}{\sqrt{a}}$ ,  $y = \frac{1}{\sqrt{b}}$  and  $z = \frac{1}{\sqrt{c}}$ , then, from ([3], 5.17)

$$ab + bc + ca \leq 4(R+r)^2,$$

we obtain

$$\begin{aligned} \sqrt{h_a} + \sqrt{h_b} + \sqrt{h_c} &\leq \sqrt{2F} \cdot \sqrt{3\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)} \\ &= \sqrt{\frac{3}{2R}(ab + bc + ca)} \\ &\leq \left(1 + \frac{r}{R}\right) \sqrt{6R}, \end{aligned}$$

i.e. the right part of (G).

**Remark 6.** The second part of the inequality (G) is more precise than

$$\sqrt{h_a} + \sqrt{h_b} + \sqrt{h_c} \leq \frac{3}{2} \sqrt{6R},$$

shown in [3].

**Theorem 8.** *In every triangle the following inequality is valid:*

$$\frac{r_b r_c}{a^2} + \frac{r_c r_a}{b^2} + \frac{r_a r_b}{c^2} \geq \frac{9}{4}. \quad (\text{H})$$

*Equality holds if and only if the triangle is equilateral.*

**Proof 1.** Inequality (H) is equivalent to

$$\frac{s-a}{a^2} + \frac{s-b}{b^2} + \frac{s-c}{c^2} \geq \frac{9}{4s}. \quad (12)$$

The function

$$f(x) = \frac{s}{x^2} - \frac{1}{x} \quad (0 < x < s)$$

is convex, so that

$$\sum_{i=1}^3 f(x_i) \geq 3 \cdot f\left(\frac{\sum_{i=1}^3 x_i}{3}\right), \quad x_i \in \bar{I}.$$

If we put  $x_1 = a$ ,  $x_2 = b$  and  $x_3 = c$ , we get the inequality (H).

**Proof 2.** As

$$\begin{aligned} & \frac{r_b r_c}{a^2} + \frac{r_c r_a}{b^2} + \frac{r_a r_b}{c^2} \\ &= \frac{s(s-a)}{a^2} + \frac{s(s-b)}{b^2} + \frac{s(s-c)}{c^2} \\ &= \frac{s^2}{a^2 b^2 c^2} (b^2 c^2 + c^2 a^2 + a^2 b^2) - \frac{s}{abc} (ab + bc + ca) \\ &= \frac{1}{16 R^2 r^2} (b^2 c^2 + c^2 a^2 + a^2 b^2 - 4Rr(ab + bc + ca)). \end{aligned} \quad (13)$$

From ([3], 5.17)

$$\begin{aligned} ab + bc + ca &\geq 4r(5R - r), \\ (bc + ab + ca)^2 &\leq 3(b^2 c^2 + c^2 a^2 + a^2 b^2) \end{aligned}$$

and (13), we have

$$\begin{aligned} \frac{r_b r_c}{a^2} + \frac{r_c r_a}{b^2} + \frac{r_a r_b}{c^2} &\geq \frac{ab + bc + ca}{48 R^2 r^2} (ab + bc + ca - 12Rr) \\ &\geq \frac{(5R - r)(2R - r)}{3R^2}. \end{aligned} \quad (14)$$

By (14) and the inequality 5.1 in [3], we obtain (H).

D. M. Milošević, Pranjani, YU

#### REFERENCES

- 1 Š. Z. Arslanagić, D. M. Milošević: Some inequalities for a triangle. Radovi matematički 2, 35–44 (1986).
- 2 A. Bager: Some inequalities for the medians of a triangle. Publication de la Faculté d'Electrotechnique de l'Université à Belgrade, série: Mathématiques et physique N° 338–352, 37–40 (1971).
- 3 O. Bottema, R. Ž. Djordjević, R. R. Janić, D. S. Mitrinović, P. M. Vasić: Geometric inequalities. Wolters-Noordhoff, Groningen 1969.
- 4 R. R. Janić, M. R. Tasković: Quelques inégalités relatives à un triangle. These Publications, N° 247–273, 131–134 (1969).

- 5 M. Jovanović: Inequalities for the triangle. These Publications № 412–460, 155–157 (1973).  
 6 D. S. Mitrinović, J. E. Pečarić, V. Volenec: Recent advances in geometric inequalities (the book) (to appear).  
 7 M. Petrović: Računanje sa brojnim razmacima. Gradjevinska knjiga, Beograd 1969.


## Ceva-Dreiecke

Ein einem Dreieck  $ABC$  einbeschriebenes Dreieck  $A_1B_1C_1$  (Fig. 1) möge *Ceva-Dreieck* heißen, wenn sich die drei Transversalen  $AA_1$ ,  $BB_1$ ,  $CC_1$  in einem Punkt  $S$  schneiden. Nach dem Satz von *Ceva* tritt dies genau dann ein, wenn die drei Teilverhältnisse

$$[ABC_1] = -\tau < 0; \quad [BCA_1] = -\varrho < 0; \quad [CAB_1] = -\sigma < 0 \quad (1)$$

der Gleichung genügen

$$\varrho \cdot \sigma \cdot \tau = 1. \quad (2)$$



Im folgenden soll gezeigt werden:

**Satz 1:** Gegeben sei ein beliebiges Dreieck  $ABC$  und ein ebenfalls beliebiges Dreieck  $A_0B_0C_0$  (Fig. 1). Dann gibt es genau ein dem Dreieck  $ABC$  einbeschriebenes Ceva-Dreieck  $A_1B_1C_1$ , das dem Dreieck  $A_0B_0C_0$  ähnlich ist.

**Beweis:** Zunächst weisen wir die Existenz eines solchen Dreiecks  $A_1B_1C_1$  nach. Dazu führen wir ein Koordinatensystem ein, das dem vorgegebenen Dreieck  $ABC$  nach Figur 2 angepasst ist.

Aus der Figur können die Bezeichnungen für die Masszahlen der Winkel (im Bogenmass) sowie die Koordinaten der Punkte (in Abhängigkeit von den Teilverhältnissen  $\varrho$ ,  $\sigma$ ,  $\tau$ ) abgelesen werden. Dabei dürfen wir ohne Einschränkung der Allgemeinheit voraussetzen, dass

$$b, c, b-a \in \mathbb{R}^+; \quad a \in \mathbb{R}; \quad \alpha_1, \beta_1, \gamma_1 \in \mathbb{R}^+; \quad \alpha_1 + \beta_1 + \gamma_1 = \pi. \quad (3)$$