Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 42 (1987)
Heft: 6

Artikel: A generalization of a two triangle inequality
Autor: Tsintsifas, G.
DOI: https://doi.org/10.5169/seals-40043

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

LITERATURVERZEICHNIS

1 L. Bieberbach: Theorie der geometrischen Konstruktionen. Birkhäuser, Basel 1952.
2 G. Fischer, R. Sacher: Einführung in die Algebra. Teubner, Stuttgart 1978.
3 S. Lang: Algebra. Addison-Wesley, 1967.
4 H. Lebesgue: Leçons sur les constructions géométriques. Gauthier-Villars, Paris 1950.
5 O. Perron: Algebra II. Göschen, Berlin 1951.
6 A. Schinzel: On linear dependence of roots. Acta Arithmetica (1975) *28*, 161–175.

© 1987 Birkhäuser Verlag, Basel 0013-6018/87/060000-00$1.50+0.20/0

A generalization of a two triangle inequality

1. Introduction

By coupling an inequality of Bottema [1] together with one of Pedoe [1], O. Bottema and M. S. Klamkin obtained the two chain inequality

$$a'x + b'y + c'z \geqq \left[\frac{P}{2} + 8FF'\right]^{1/2} \geqq 4\sqrt{FF'}\,,$$

see [2], where $P = \sum a'^2(b^2 + c^2 - a^2)$, x, y, z the distances of an interior point M of the triangle ABC from the vertices A, B, C, a, b, c and a', b', c' the sides of the triangles ABC and $A'B'C'$ and F, F' their area respectively.
In this note the author will generalize the part

$$a'x + b'y + c'z \geqq 4\sqrt{FF'}$$

of the above inequality for two simplices $s^{(n)} = (A_1A_2 \ldots A_{n+1})$ and $s'^{(n)} = (A'_1A'_2 \ldots A'_{n+1})$.

2. Notations

We denote by $V(W)$ the volume of the simplex W, $s^{(n)} = (A_1A_2 \ldots A_{n+1})$, $s'^{(n)} = (A'_1A'_2 \ldots A'_{n+1})$ two simplices of E^n. The facets

$$(A_1A_2 \ldots A_{i-1}A_{i+1} \ldots A_{n+1}),\ (A'_1A'_2 \ldots A'_{i-1}A'_{i+1} \ldots A'_{n+1})$$

will be denoted by $s_i^{(n-1)}$, $s_i'^{(n-1)}$ respectively.
Suppose that M is an interior point of the simplex $s^{(n)}$ with distances $A_iM = x_i$. We put:

$$D = \sum_{i=1}^{n+1} x_i V(s_i'^{(n-1)})\,.$$



Figure 1

The quantity D depends on the labeling of the vertex set but for every case we can prove:

3. Theorem

For two simplices $s^{(n)}$ and $s'^{(n)}$ holds:

$$D^2 \geqq n^4 \cdot V^{\frac{n}{2}}(s^{(n)}) \cdot V^{\frac{2n-2}{2}}(s'^{(n)}) .$$

Proof: We assume that the simplex $p^{(n)} = (B_1 B_2 \ldots B_n)$ is similar to $s'^{(n)}$ and inscribed in $s^{(n)}$, so that $B_i \in s_i^{(n-1)}$. We also consider the simplex $q^{(n)} = (C_1 C_2 \ldots C_{n+1})$ similar to $s'^{(n)}$ and circumscribed to $s^{(n)}$ so that $A_i \in q_i^{(n-1)}$, where $q_i^{(n-1)} = (C_1 C_2 \ldots C_{i-1} C_{i+1} \ldots C_{n+1})$. It is known that:

$$V^n(s^{(n)}) \geqq V^{n-1}(p^{(n)}) \cdot V(q^{(n)}) , \tag{1}$$

see [3].
Suppose $W' = kW$ is the homothetical image of W with ratio k. We put:

$$p^{(n)} = \lambda s'^{(n)} \quad \text{and} \quad q^{(n)} = \mu s'^{(n)} . \tag{2}$$

We can easily see that: $x_i \cdot V(p_i^{(n-1)}) \geqq n V(A_i B_1 B_2 \dots B_{i-1} B_{i+1} \dots B_{n+1} M)$ and $x_i \cdot V(q_i^{(n-1)}) \geqq n V(C_1 C_2 \dots C_{i-1} C_{i+1} \dots C_{n+1} M)$. From the above follows

$$\sum_{i=1}^{n+1} x_i V(p_i^{(n-1)}) \geqq n V(s^{(n)}), \tag{3}$$

$$\sum_{i=1}^{n+1} x_i V(q_i^{(n-1)}) \geqq n V(q^n), \tag{4}$$

where $p_i^{(n-1)} = (B_1 B_2 \dots B_{i-1} B_{i+1} \dots B_{n+1})$.
See the figure, above, for the elementary case $n = 2$. From (2) follows,

$$V(p_i^{(n-1)}) = \lambda^{n-1} V(s'^{(n-1)}), \tag{5a}$$

$$V(q_i^{(n-1)}) = \mu^{n-1} V(s'^{(n-1)}) \quad \text{and} \tag{5b}$$

$$V(q^{(n)}) = \mu^n V(s'^{(n)}), \tag{5c}$$

see (2).
Therefore from (3), (4), (5) we have:

$$\lambda^{n-1} D \geqq n V(s^{(n)}), \quad \mu^{n-1} D \geqq n \mu^n V(s'^{(n)}) \tag{6}$$

or,

$$\frac{\lambda^{n-1}}{\mu} D^2 \geqq n^2 V(s^{(n)}) V(s'^{(n)}). \tag{7}$$

Using (1), (5c) we take:

$$\frac{V^n(s^{(n)})}{V^2(q^{(n)})} \geqq \frac{\lambda^{n(n-1)} V^{n-1}(s'^{(n)})}{\mu^n V(s'^{(n)})} = \left(\frac{\lambda^{n-1}}{\mu}\right)^n V^{n-2}(s'^{(n)})$$

or,

$$V^{n-2}(s^{(n)}) \left(\frac{V(s^{(n)})}{V(q^{(n)})}\right)^2 \geqq \left(\frac{\lambda^{n-1}}{\mu}\right)^n V^{n-2}(s'^{(n)}). \tag{8}$$

But it is known that if $V(q^{(n)})$ is minimum then A_i is the centroid of $q_i^{(n-1)}$, see [4] or [5], therefore

$$\frac{1}{n^n} \geqq \frac{V(s^{(n)})}{V(q^{(n)})}. \tag{9}$$

From (8), (9) follows:

$$\frac{V^{n-2}(s^{(n)})}{n^{2n}} \geqq \left(\frac{\lambda^{n-1}}{\mu}\right)^n V^{n-2}(s'^{(n)})$$

or,

$$\frac{1}{n^2}\frac{V^{\frac{n-2}{n}}(s^{(n)})}{V^{\frac{n-2}{n}}(s'^{(n)})} \geqq \frac{\lambda^{n-1}}{\mu}. \tag{10}$$

Therefore from (7) and (10) we obtain

$$D^2 \geqq n^4 V^{\frac{2}{n}}(s^{(n)})\, V^{\frac{2n-2}{n}}(s'^{(n)}).$$

The author is grateful to the referee for his helpful suggestions.

G. Tsintsifas, Thessaloniki, Greece

REFERENCES

1 O. Bottema, R. Z. Djordjević, R. R. Janić, D. S. Mitrinović, and P. M. Vasić: Geometric inequalities. Wolters-Noordhoff, Groningen 1969.
2 O. Bottema and M. S. Klamkin: Joint triangle inequalities. Simon Stevin, wis-en Natuur'Kundig Tijdschrift 48e Jeargang (1974–1975), Afleviving I, II (Juli–October 1974).
3 G. D. Chakerian: Minimum area of circumscribed polygons. Elemente der Mathematik, Vol. 28, Heft 5, 1973.
4 M. M. Day: Polygons circumscribed about closed convex sets. Trans. Amer. Math. Soc. *62*, pp. 315–319, 1957.
5 G. D. Chakerian and L. H. Lange: Geometric extremum Problems. Math. Magazine *44*, N° 2, 1971.

© 1987 Birkhäuser Verlag, Basel 0013-6018/87/060000-00$1.50+0.20/0

A tournament result deduced from harems

There is a large class of difficult problems of the type: "does there exist a graph with n vertices having prescribed degrees $d_1, \ldots, d_n$?" Restricting the problem to particular types of graphs can lead to some very neat characterisations. For example, it is a straightforward exercise to show that a tree exists on $n\,(\geqq 2)$ vertices with degrees $d_1, \ldots, d_n$ if and only if the $d_1, \ldots, d_n$ are positive integers with

$$d_1 + \ldots + d_n = 2(n-1).$$

We shall now restrict attention to 'tournaments'. A *tournament* is a directed graph in which each pair of distinct vertices is joined precisely once (in one direction or the other). Alternatively it can be thought of as a competition of a set of players in which each pair plays once resulting in a win for one of the players. Before proceeding, note that, for example, there exists a tournament of 4 players in which their numbers of wins are 1, 1, 2 and 2 (e.g. A beats B, B beats D, C beats A, C beats B, D beats A and