

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 43 (1988)
Heft: 2

Rubrik: Kleine Mitteilung

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Zahlreiche Eigenschaften von E findet man in [5–7], [10] und in der dort zitierten Literatur.

Gould und Mays [4] haben gezeigt, dass die einzigen Mittelwerte, die sowohl $E(r, s; x, y)$ als auch $L_r(x, y)$ angehören, das arithmetische, das geometrische und das harmonische Mittel von x und y sind.

Der Redaktion möchte ich für Verbesserungsvorschläge herzlich danken.

Horst Alzer, Waldbröl

LITERATUR

- 1 E. F. Beckenbach: A class of mean value functions. Amer. Math. Monthly 57, 1–6 (1950).
- 2 D. Farnsworth and R. Orr: Gini means. Amer. Math. Monthly 93, 603–607 (1986).
- 3 C. Gini: Di una formula comprensiva delle medie. Metron 13, 3–22 (1938).
- 4 H. W. Gould and M. E. Mays: Series expansions of means. J. Math. Anal. Appl. 101, 611–621 (1984).
- 5 E. B. Leach and M. C. Sholander: Extended mean values. Amer. Math. Monthly 85, 84–90 (1978).
- 6 E. B. Leach and M. C. Sholander: Extended mean values II. J. Math. Anal. Appl. 92, 207–223 (1983).
- 7 E. B. Leach and M. C. Sholander: Multi-variable extended mean values. J. Math. Anal. Appl. 104, 390–407 (1984).
- 8 D. H. Lehmer: On the compounding of certain means. J. Math. Anal. Appl. 36, 183–200 (1971).
- 9 D. S. Mitrinović: Analytic Inequalities. Springer Verlag, Berlin 1970.
- 10 K. B. Stolarsky: Generalizations of the logarithmic mean. Math. Mag. 48, 87–92 (1975).
- 11 E. M. Wright: A generalization of Schur's inequality. Math. Gaz. 40, 217 (1956).

Kleine Mitteilung

Sums of a certain family of series

By identifying the sum

$$S_n = \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \frac{1}{2^k (n+k+1)} \quad (n \in \mathcal{N} = \{1, 2, 3, \dots\}) \quad (1)$$

with the integral

$$S_n = \int_0^1 \left(1 - \frac{t}{2}\right)^{n-1} t^n dt, \quad (2)$$

and evaluating this Eulerian integral, M. Vowe and H.-J. Seiffert [3] have recently shown that

$$S_n = \frac{2^n (n-1)! n!}{(2n)!} - \frac{2^{-n}}{n} \quad (n \in \mathcal{N}). \quad (3)$$

In our attempt to find the sum in (1), *without* evaluating the integral in (2), we are led naturally to the fact that the formula (3) is just one of numerous interesting (and useful) consequences of a known result in the theory of the Gaussian hypergeometric series

$$F(a, b; c; z) = 1 + \frac{a b}{c} \frac{z}{1!} + \frac{a(a+1) b(b+1)}{c(c+1)} \frac{z^2}{2!} + \dots, \quad (4)$$

which, for $a = 1$ and $b = c$ (or, alternatively, for $a = c$ and $b = 1$), reduces immediately to the familiar geometric series. In one of his 1836 memoirs [1], Ernst Eduard Kummer (1810–1893) proved the summation theorem [1, p. 134, Theorem 3]:

$$F(a, 1-a; c; \frac{1}{2}) = \frac{\Gamma\left(\frac{c}{2}\right) \Gamma\left(\frac{c+1}{2}\right)}{\Gamma\left(\frac{c+a}{2}\right) \Gamma\left(\frac{c-a+1}{2}\right)} \quad (c \neq 0, -1, -2, \dots), \quad (5)$$

where, as usual, $\Gamma(z)$ denotes the familiar Gamma function satisfying the relationships:

$$\begin{cases} \Gamma(z+1) = z \Gamma(z), & \sqrt{\pi} \Gamma(2z) = 2^{2z-1} \Gamma(z) \Gamma(z+1/2), \\ \Gamma(n+1) = n! & (n \in \mathbb{N} \cup \{0\}), \Gamma(1/2) = \sqrt{\pi} \end{cases} \quad (6)$$

(see also Srivastava and Karlsson [2, pp. 18–19]).

From the definition

$$\binom{\lambda}{0} = 1; \quad \binom{\lambda}{k} = \frac{\lambda(\lambda-1)\dots(\lambda-k+1)}{k!} \quad (k \in \mathbb{N}), \quad (7)$$

for an arbitrary (real or complex) λ , it follows readily that

$$\binom{\lambda+k-1}{k} = \frac{\lambda(\lambda+1)\dots(\lambda+k-1)}{k!} \quad (k \in \mathbb{N} \cup \{0\}). \quad (8)$$

Making use of (8), and the second relationship in (6), it is fairly easy to state Kummer's summation theorem (5) in the (more relevant) form:

$$S_{\lambda, \mu} \equiv \sum_{k=0}^{\infty} (-1)^k \binom{\lambda-1}{k} \frac{\binom{\lambda+k-1}{k}}{2^k \binom{\mu+k-1}{k}} = \frac{2^{1-\mu} \sqrt{\pi} \Gamma(\mu)}{\Gamma\left(\frac{\mu+\lambda}{2}\right) \Gamma\left(\frac{\mu-\lambda+1}{2}\right)} \quad (9)$$

Since $(\mu \neq 0, -1, -2, \dots)$.

$$\binom{n-1}{k} = 0, \quad k = n, n+1, n+2, \dots, \quad (10)$$

the sum in (9) would terminate at $k = n - 1$ in the special case when $\lambda = n \in \mathcal{N}$. In particular, we have

$$S_{n,n} \equiv \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} 2^{-k} = 2^{1-n} \quad (n \in \mathcal{N}). \quad (11)$$

$$S_{n,n+1} \equiv \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \frac{1}{2^k (n+k)} = \frac{2^n (n-1)! n!}{(2n)!} \quad (n \in \mathcal{N}), \quad (12)$$

and

$$S_{n,n+2} \equiv \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \frac{1}{2^k (n+k)(n+k+1)} = \frac{2^{-n}}{n} \quad (n \in \mathcal{N}). \quad (13)$$

Formula (11) is an obvious consequence of the familiar binomial theorem:

$$\sum_{k=0}^n \binom{n}{k} a^k b^{n-k} = (a+b)^n \quad (n \in \mathcal{N} \cup \{0\}), \quad (14)$$

or, more generally,

$$\sum_{k=0}^{\infty} \binom{\lambda}{k} z^k = (1+z)^\lambda \quad (|z| < 1; \lambda \text{ arbitrary}), \quad (15)$$

which incidentally is related to (4) with $a = -\lambda$, $b = c$, and z replaced by $-z$. Formulas (12) and (13), together, yield

$$\begin{aligned} S_n &\equiv S_{n,n+1} - S_{n,n+2} = \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \frac{1}{2^k (n+k+1)} \\ &= \frac{2^n (n-1)! n!}{(2n)!} - \frac{2^{-n}}{n} \quad (n \in \mathcal{N}), \end{aligned} \quad (16)$$

which is precisely the summation formula (3) given by Vowe and Seiffert [3]. It is not difficult to deduce from (9) the following generalization of (3):

$$\sum_{k=0}^{\infty} (-1)^k \binom{\lambda-1}{k} \frac{1}{2^k (\lambda+k+1)} = \frac{2^\lambda \Gamma(\lambda) \Gamma(\lambda+1)}{\Gamma(2\lambda+1)} - \frac{2^{-\lambda}}{\lambda} \quad (\lambda \neq 0, -1, -2, \dots), \quad (17)$$

which holds true for an essentially arbitrary (real or complex) λ .

Some further consequences of the general result (9) are worthy of note. Indeed, for every non-negative integer l , we obtain

$$\begin{aligned} S_{\lambda, \lambda+2l} &\equiv \sum_{k=0}^{\infty} (-1)^k \binom{\lambda-1}{k} \frac{1}{2^k \prod_{j=1}^{2l} (\lambda+k+j-1)} \\ &= \frac{2^{1-\lambda} l!}{(2l)! \prod_{j=1}^l (\lambda+j-1)} \quad (\lambda \neq 0, -1, -2, \dots) \end{aligned} \quad (18)$$

and

$$\begin{aligned} S_{\lambda, \lambda+2l+1} &\equiv \sum_{k=0}^{\infty} (-1)^k \binom{\lambda-1}{k} \frac{1}{2^k \prod_{j=0}^{2l} (\lambda+k+j)} \\ &= \frac{2^\lambda \Gamma(\lambda) \Gamma(\lambda+l+1)}{l! \Gamma(2\lambda+2l+1)} \quad (\lambda \neq 0, -1, -2, \dots), \end{aligned} \quad (19)$$

where, as usual, an empty product is to be interpreted as 1.

Upon subtracting (18) from (19) with l replaced by $l-1$, we find that

$$\begin{aligned} &\sum_{k=0}^{\infty} (-1)^k \binom{\lambda-1}{k} \frac{\lambda+k+2l-2}{2^k \prod_{j=1}^{2l} (\lambda+k+j-1)} \\ &= \frac{2^\lambda \Gamma(\lambda) \Gamma(\lambda+l)}{(l-1)! \Gamma(2\lambda+2l-1)} - \frac{2^{1-\lambda} l!}{(2l)! \prod_{j=1}^l (\lambda+j-1)} \quad (l \in \mathcal{N}), \end{aligned} \quad (20)$$

which evidently yields (17) when $l = 1$.

Each of the summation formulas (18), (19), and (20) would terminate, by virtue of (10), in its special case when $\lambda = n \in \mathcal{N}$. Formula (20) thus yields

$$\begin{aligned} &\sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \frac{n+k+2l-2}{2^k \prod_{j=1}^{2l} (n+k+j-1)} \\ &= \frac{2^n (n-1)! (n+l-1)!}{(l-1)! (2n+2l-2)!} - \frac{2^{1-n} l!}{(2l)! \prod_{j=1}^l (n+j-1)} \quad (n, l \in \mathcal{N}), \end{aligned} \quad (21)$$

which provides us with yet another generalization of the summation formula (3).

H. M. Srivastava, Department of Mathematics, University of Victoria, Canada

REFERENCES

- 1 E. E. Kummer: Über die hypergeometrische Reihe J. Reine Angew. Math. 15, 39–83 and 127–172 (1836).
- 2 H. M. Srivastava, P. W. Karlsson: Multiple Gaussian Hypergeometric Series. Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto 1985.
- 3 M. Vowe, H.-J. Seiffert: Aufgabe 946. Elem. Math. 42, 111–112 (1987).

© 1988 Birkhäuser Verlag, Basel

0013-6018/88/020054-05 \$ 1.50 + 0.20/0

Aufgaben

Aufgabe 957. Man beweise die Ungleichung

$$\sin(x/2) + \cos x < (\pi - x)/2; \quad 0 < x < \pi.$$

P. Ivady, Budapest, Ungarn

Lösung. Für $0 < x < \pi$ gilt nach einer bekannten Identität

$$\begin{aligned} \sin(x/2) + \cos x &= 2 \sin((\pi - x)/4) \cos((3x - \pi)/4) \\ &< 2 \sin((\pi - x)/4) \\ &< (\pi - x)/2. \end{aligned}$$

A. A. Jagers, Enschede, NL

Weitere Lösungen sandten S. Arslanagic (Trebinje, YU), A. Bender (Zürich), H. Bopp (Illingen), E. Braune (Linz, A), P. Bracken (Toronto, CD), P. Bundschuh (Köln, BRD), F. Götze (Jena, DDR), M. Hübner (Leipzig, DDR), W. Janous (Innsbruck, A), L. Kuipers (Sierre), Kee-wai Lau (Hongkong), I. Merenyi (Berveni, RU), A. Müller (Zürich), P. Müller (Nürnberg, BRD), H.-J. Seiffert (Berlin), Tsen-Pao Shen (München, BRD), H. M. Smid (Amsterdam, NL), M. Vowe (Therwil), R. Wyss (Flumenthal).

Aufgabe 958. Es seien

$$x_n := \sum_{k=1}^n (-1)^{k+1} \frac{1}{k} \binom{n}{k} \quad \text{und} \quad y_n := \sum_{k=1}^n (-1)^{k+1} \frac{1}{k} \binom{n-1}{k-1}.$$

Man berechne: $\lim_{n \rightarrow \infty} (x_n + \log y_n)$.

H. Alzer, Waldbröl, BRD