

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 43 (1988)
Heft: 4

Artikel: Periodicity of p-adic continued fractions
Autor: Weger, B.M.M. de
DOI: <https://doi.org/10.5169/seals-40807>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Wir verzichten notgedrungen auf weitere Darstellungen und auch auf spezielle Literaturangaben. Statt dessen nennen wir lediglich einige Übersichtswerke. Sollte ein Leser noch nicht völlig verschreckt sein (hoffentlich gibt es solche!) und gar den Wunsch haben, noch mehr zu erfahren, so sei er auf die ausführliche Bibliographie zur Designtheorie in [3] verwiesen (sie wurde inzwischen auf über 1000 Titel erweitert!). Dort kann er unter den im vorliegenden Text angegebenen Autorennamen genaue Literaturhinweise finden und dann mit dem Studium dieses hochaktuellen, aber auch äusserst interessanten Gebietes beginnen. Viel Vergnügen und viel Erfolg!

H. Zeitler, Math. Institut, Universität Bayreuth

LITERATUR

- 1 Beth T., Jungnickel J., Lenz H.: *Design Theory*. Mannheim-Wien-Zürich (1985).
- 2 Hughes D. R., Piper F. C.: *Design Theory*. Cambridge (1985).
- 3 Lindner C. C., Rosa A.: *Topics on Steiner Systems*. Amsterdam-New York-Oxford (1980).

© 1988 Birkhäuser Verlag, Basel

0013-6018/88/040097-16\$1.50 + 0.20/0

Periodicity of p -adic continued fractions

A well known theorem of Lagrange states that the simple continued fraction expansion of a real number is periodic if and only if that real number is quadratic irrational. Several authors have tried to establish analogous results for continued fractions of p -adic numbers. The present author [3] showed that such a result is possible if one starts with sequences of approximation lattices of p -adic numbers, instead of continued fractions. We note that from a periodic sequence of approximation lattices of a p -adic number ξ it is easy to construct a periodic continued fraction expansion of ξ .

This process of constructing the continued fraction such that it is periodic, is the reverse of the process in the real case, where one starts by defining the continued fraction, and then tries to prove its periodicity. It would be interesting to obtain periodicity results for a given p -adic continued fraction expansion method, e.g. that introduced by Schneider [2]. Bundschuh [1] remarks that for this type of p -adic continued fractions a periodic continued fraction represents either a rational p -adic number of special type, or a quadratic irrational p -adic number (analogous to Euler's theorem). Further, he gives some numerical evidence indicating that the converse (analogous to Lagrange's theorem) may not be true.

It is the purpose of this note to show that for Schneider's continued fraction algorithm for p -adic numbers, it may indeed happen that quadratic irrational numbers in \mathbb{Q}_p have non-periodic continued fraction expansions. Thus for this type of continued fractions an

analogue of Lagrange's theorem is not true. Some criteria (but not exhaustive) will be given for (non)-periodicity. The examples that Bundschuh treated numerically are proved to be non-periodic.

Let p be prime. Let $\xi \in \mathbb{Q}_p$ be a nonzero p -adic integer. Schneider [2] defines the continued fraction expansion of ξ as follows. Put $\xi_0 = \xi$. Given a p -adic integer ξ_n for some $n \in \mathbb{N}_0$, let $a_n \in \{0, 1, \dots, p-1\}$ be such that $b_n = \text{ord}_p(\xi_n - a_n)$ is positive. We continue only if $\xi_n \neq a_n$. Then define ξ_{n+1} by

$$\xi_n = a_n + \frac{p^{b_n}}{\xi_{n+1}}.$$

Then $|\xi_n|_p = 1$ and $a_n \neq 0$ for all $n > 0$. Now the continued fraction expansion of ξ is

$$\xi = a_0 + \frac{p^{b_0}}{\lceil a_1 \rceil} + \frac{p^{b_1}}{\lceil a_2 \rceil} + \dots$$

Note that Bundschuh [1] has a slightly, but not essentially, different definition and notation.

We say that the continued fraction expansion of ξ is periodic if there exist $m_0 \in \mathbb{N}_0$, $k \in \mathbb{N}$ such that $a_{m+k} = a_m$, $b_{m+k} = b_m$ for all $m \geq m_0$. Bundschuh [1] asks to prove or disprove periodicity for the continued fractions of $\xi = \sqrt{c}$, for $c \in \mathbb{Z}$ not a square, but a quadratic residue $(\bmod p)$, so that $\xi \in \mathbb{Q}_p$, a.o. for the four examples $(c, p) = (-1, 5), (2, 7), (5, 11), (3, 13)$.

There exist unique rational numbers P_n, Q_n such that

$$\xi_n = \frac{P_n + \sqrt{c}}{Q_n}$$

for $n \in \mathbb{N}_0$. Then $P_0 = 0$, $Q_0 = 1$, and we have the recursion formulas

$$\begin{aligned} P_{n+1} &= -(P_n - a_n \cdot Q_n), \\ Q_{n+1} &= (c - P_{n+1}^2)/p^{b_n} \cdot Q_n. \end{aligned}$$

We show that P_n and Q_n are integers, and that $Q_n | c - P_n^2$, for all $n \in \mathbb{N}_0$. It is obvious for $n = 0$. Suppose it is true for some $n \geq 0$. Then P_{n+1} is obviously an integer. Further,

$$c - P_{n+1}^2 = c - P_n^2 + 2 \cdot a_n \cdot P_n \cdot Q_n - a_n^2 \cdot Q_n^2 \equiv 0 \pmod{Q_n},$$

and

$$\begin{aligned} c - P_{n+1}^2 &= (P_{n+1} + \sqrt{c}) \cdot (-P_{n+1} + \sqrt{c}) \\ &= (P_{n+1} + \sqrt{c}) \cdot (P_n - a_n Q_n + \sqrt{c}) \\ &= (P_{n+1} + \sqrt{c}) \cdot (\xi_n - a_n) \cdot Q_n \equiv 0 \pmod{p^{b_n}}, \end{aligned}$$

hence Q_{n+1} is integral. It follows at once that $Q_{n+1} | c - P_{n+1}^2$. We now prove the following lemma.

Lemma. If for some n the signs of P_n and Q_n are different, and $P_{n+1}^2 > c$, then the continued fraction of $\sqrt{c} \in \mathbb{Q}_p$ is non-periodic.

Proof. Since P_n and Q_n have different signs, it follows by the recursion formula for P_{n+1} that P_n and P_{n+1} have different signs. By the recursion formula for Q_{n+1} and by $P_{n+1}^2 > c$ it follows that $Q_{n+1} \neq 0$, and Q_n and Q_{n+1} have different signs. Hence the signs of P_{n+1} and Q_{n+1} are different. Further, by $a_{n+1} \neq 0$,

$$|P_{n+2}| = |P_{n+1}| + a_{n+1} \cdot |Q_{n+1}| > |P_{n+1}|,$$

so that $P_{n+2}^2 > c$. Consequently, the conditions for the lemma hold for $n+1$ as well. Hence, by induction, $|P_n| \rightarrow \infty$ as $n \rightarrow \infty$. But periodicity of the continued fraction implies periodicity of P_n and Q_n , which contradicts that $|P_n|$ is unbounded. \square

Corollary. If $c < 0$ then the continued fraction of $\sqrt{c} \in \mathbb{Q}_p$ is non-periodic.

Proof. If $c < 0$ then always $P_n^2 > c$. Further, $P_1 = a_0 > 0$, and $Q_1 = (c - P_1^2)/p^{b_0} < 0$. Apply the lemma for $n = 1$. \square

Examples. The four examples of Bundschuh all satisfy the conditions for the above lemma with $n = 1$. We give a few details below.

1. $\sqrt{-1} \in \mathbb{Q}_5$. The non-periodicity follows from the corollary. We have in fact, if we take $\sqrt{-1} \equiv 2 \pmod{5}$,

n	P_n	Q_n	a_n	b_n
0	0	1	2	1
1	2	-1	1	1
2	-3	2	2	2

and the continued fraction starts with

$$2 + \frac{5}{1} + \frac{5}{2} + \frac{25}{1} + \frac{5}{3} + \frac{5}{4} + \frac{5}{2} + \frac{5}{1} + \frac{125}{1} + \frac{5}{4} + \dots$$

2. $\sqrt{2} \in \mathbb{Q}_7$. We take $\sqrt{2} \equiv 3 \pmod{7}$. Then we have

n	P_n	Q_n	a_n	b_n
0	0	1	3	1
1	3	-1	1	1
2	-4	2	3	2

3. $\sqrt{5} \in \mathbb{Q}_{11}$. We take $\sqrt{5} \equiv 4 \pmod{11}$. Then we have

n	P_n	Q_n	a_n	b_n
0	0	1	4	1
1	4	-1	3	1
2	-7	4	2	1

4. $\sqrt{3} \in \mathbb{Q}_{13}$. We take $\sqrt{3} \equiv 4 \pmod{13}$. Then we have

n	P_n	Q_n	a_n	b_n
0	0	1	4	1
1	4	-1	5	1
2	-9	6	10	1

Next we show that there are also many p -adic quadratic irrationals that do have periodic continued fraction expansions, as defined above. Let $c \in \mathbb{N}$ be a non-square, that can be written as $c = e^2 + d \cdot p^k$ for $d, e, k \in \mathbb{N}$ with $1 \leq e \leq \frac{1}{2}(p - 1)$ and $d \nmid 2e, p \nmid d$. Then we find

n	P_n	Q_n	a_n	b_n
0	0	1	e	k
1	e	d	$2e/d$	k
2	e	1	$2e$	k
3	e	d	$2e/d$	k

so that the continued fraction is periodic with period length 2.

We conclude by giving some «exceptional» pairs (c, p) which do have periodic continued fractions, but seem not to fit in an infinite sequence, such as given above. They are: $(c, p) = (136, 3), (376, 5), (148, 7), (388, 11)$. We have (with the bar denoting the repeating part):

$$\sqrt{136} = 1 + \overline{[27; 1, 3, 1, 27]},$$

$$\sqrt{376} = 1 + \overline{[125; 4, 5, 3, 1, 5, 4, 2]},$$

$$\sqrt{148} = 1 + \overline{[49; 3, 4, 3, 2]},$$

$$\sqrt{388} = 5 + \overline{[121; 7, 8, 7, 10]}.$$

Suggestions for further research. We observe that in all cases presented above, as in the real case, the periodic continued fractions are symmetric. Namely, for period length k we have $a_j = a_{k-j}$ and $b_j = b_{k+1-j}$ for $j = 1, 2, \dots, k-1$. Further, $a_k = 2 \cdot a_0$ appears to hold, as in the real case. We found no examples of odd period length.

There is also a connection to Pell-like equations. We illustrate this with the example $\sqrt{376} \in \mathbb{Q}_5$. Let p_n/q_n be the n th convergent of the continued fraction expansion, defined by

$$\begin{aligned} p_{-1} &= 1, \quad p_0 = a_0, \quad p_n = a_n \cdot p_{n-1} + 5^{b_n} \cdot p_{n-2} \quad \text{for } n \geq 1, \\ q_{-1} &= 0, \quad q_0 = 1, \quad q_n = a_n \cdot q_{n-1} + 5^{b_n} \cdot q_{n-2} \quad \text{for } n \geq 1. \end{aligned}$$

Put

$$p_n^2 - 376 \cdot q_n^2 = d_n \cdot 5^{c_n}, \quad c_n \in \mathbb{N}, \quad d_n \in \mathbb{Z}, \quad 5 \nmid d_n.$$

Then we find that $c_n = \sum_{j=0}^n b_j$, and the sequence $\{d_n\}_{n=-1}^{\infty}$ is given by 1, -3, 17, -4, 17, -3, 1, -3, 17, ..., which is symmetric. The fifth convergent $p_5/q_5 = 12103/603$, for which $d_5 = 1$, gives rise to a sort of 5-adic fundamental unit $12103 + 603 \cdot \sqrt{376}$, in the sense that $p_{i+6j} + q_{i+6j} \cdot \sqrt{376} = (p_i + q_i \cdot \sqrt{376}) \cdot (p_5 + q_5 \cdot \sqrt{376})^j$ for $i = -1, 0, \dots, 4$, and $j = 0, 1, 2, \dots$. It would be interesting to have a more general theory of these matters.

B. M. M. de Weger, Faculty of Applied Mathematics,
University of Twente, Enschede

REFERENCES

- 1 Bundschuh P.: p -adische Kettenbrüche und Irrationalität p -adischer Zahlen. *Elem. Math.* 32, 36–40 (1977).
- 2 Schneider Th.: Über p -adische Kettenbrüche. *Symposia Math.* Vol. IV, 181–189 (1970).
- 3 de Weger B. M. M.: Approximation Lattices of p -adic Numbers. *J. Number Th.* 24, 70–88 (1986).

A further remark concerning Nagel cevians

Dedicated to Professor D. S. Mitrinović on his 80th birthday

In El. Math., vol. 41/5 and 42/4, R. H. Eddy and D. S. Milošević considered the cases $L = 14$ and (the improvement) $L = 10$, resp., of

$$\sum n_a \leq 9r + L(R - 2r). \tag{1}$$

Here n_a, n_b, n_c are the Nagel cevians and R, r the circumradius and inradius, resp., of the given triangle.