Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 43 (1988)
Heft: 6

Artikel: Some integral inequalities
Autor: Sándor, Joseph
DOI: https://doi.org/10.5169/seals-40815

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 08.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

REFERENCES

1 Berger J. O.: Statistical decision theory and Bayesian analysis, 2nd ed., Springer, Berlin – Heidelberg – New York 1985.
2 Hodges J. L. and Lehmann E. L.: Some problems in minimax point estimation. Ann. Math. Statist. *21*, 182–197 (1950).
3 Lehmann E. L.: Theory of point estimation. Wiley, New York 1983.
4 Loeffel H.: Statistische Inferenz und strategisches Spiel. El. Math. *40*, 109–120 (1985).
5 Neumann J. v.: Zur Theorie der Gesellschaftsspiele. Math. Annalen *100*, 295–320 (1928).
6 Rauhut B., Schmitz N. und Zachow E.-W.: Spieltheorie. Teubner, Stuttgart 1979.

© 1988 Birkhäuser Verlag, Basel 0013-6018/88/060170-08 $1.50 + 0.20/0

Some integral inequalities

The aim of this note is to prove some integral inequalities and to find interesting applications for the logarithmic and exponential functions. These relations have some known corollaries ([3], [4], [5], [8]).

Theorem 1. *Let $f:[a,b]\to\mathbb{R}$ $(a<b)$ be a differentiable function with increasing (strictly increasing) derivative on $[a,b]$. Then one has the following inequalities:*

$$\int_a^b f(t)\,dt \underset{(>)}{\geq} (b-a)\,f\left(\frac{a+b}{2}\right) \tag{1}$$

$$2\cdot\int_a^b f(t)\,dt \underset{(<)}{\leq} (b-a)\,f(\sqrt{ab}) + (\sqrt{b}-\sqrt{a})(\sqrt{b}\,f(b)+\sqrt{a}\,f(a))$$

$$(\textit{Here } 0\leq a<b). \tag{2}$$

Proof. The Lagrange mean-value theorem implies: $f(y)-f(x)\underset{(>)}{\geq}(y-x)\,f'(x)$ for all $x,y\in[a,b]$. Take $x=(a+b)/2$ and integrate the obtained inequality:

$$\int_a^b f(y)\,dy-(b-a)\,f\left(\frac{a+b}{2}\right)\underset{(>)}{\geq} f'\left(\frac{a+b}{2}\right)\cdot\int_a^b\left(y-\frac{a+b}{2}\right)dy=0,$$

i.e. relation (1).

In order to prove (2) consider as above the inequality $f(y)-f(x)\underset{(<)}{\leq}(y-x)\,f'(y)$ with $x=\sqrt{ab}$. Integrating by parts on $[a,b]$ we get

$$\int_a^b f(y)\,dy-(b-a)\,f(\sqrt{ab})\underset{(<)}{\leq}(y-\sqrt{ab})\,f(y)\Big|_a^b-\int_a^b f(y)\,dy$$

which easily implies (2).

Remark. Inequality (1) is called sometimes "Hadamard's inequality" and it is valid for convex functions f as well with the same proof, but using $f'_+\left(\frac{a+b}{2}\right)$ instead of $f'\left(\frac{a+b}{2}\right)$ (see also [1]).

In applications is useful the following generalization of (1) (see [9])

Theorem 2. *Let $f:[a,b]\to\mathbb{R}$ be a $2k$-times differentiable function, having continuous $2k$-th derivative on $[a,b]$ and satisfying $f^{(2k)}(t)\underset{(>)}{\geq}0$ for $t\in(a,b)$. Then one has the inequality:*

$$\int_a^b f(t)\,dt \underset{(>)}{\geq} \sum_{p=1}^{k} \frac{(b-a)^{2p-1}}{2^{2p-2}(2p-1)!} f^{(2p-2)}\left(\frac{a+b}{2}\right). \tag{3}$$

Proof. Apply Taylor's formula (with Lagrange remainder term) for f around $\left(\frac{a+b}{2}\right)$ and integrate term by term this relation. Remarking that $\int_a^b \left(x-\frac{a+b}{2}\right)^{2m-1} dx = 0$ for $m = 1, 2, 3, \ldots$, we obtain

$$\int_a^b f(x)\,dx = (b-a)\,f\left(\frac{a+b}{2}\right) + \frac{(b-a)}{2^2\,3!} f''\left(\frac{a+b}{2}\right)$$
$$+ \ldots + \frac{(b-a)^{2k-1}}{2^{2k-2}(2k-1)!} f^{(2k-2)}\left(\frac{a+b}{2}\right) + \int_a^b \frac{(x-(a+b)/2)^{2k}}{(2k)!} f^{(2k)}(\xi)\,dx.$$

Taking into account $f^{(2k)}(\xi)\underset{(>)}{\geq}0$, we get the desired inequality (3).

Applications. 1) Let $a>0$, $b=a+1$, $f_1(t)=\frac{1}{t}$ and $f_2(t)=-\ln t$ in (1). We can easily deduce the following double inequality:

$$\frac{2a+2}{2a+1} < \frac{e}{\left(1+\frac{1}{a}\right)^a} < \sqrt{1+\frac{1}{a}} \tag{4}$$

containing inequalities studied by E. R. Love [4] and G. Pólya – G. Szegö [7]. Using Bernoulli's inequality we have $(1+1/(2a+1))^{5/2} > 1+5/(4a+2) \geq 1+1/a$, for $a\geq 2$. Hence we have:

$$\left(1+\frac{1}{a}\right)^{a+2/5} < e < \left(1+\frac{1}{a}\right)^{a+1/2} \quad (a\geq 2) \tag{5}$$

2) By repeating the same argument in (3) for $k = 2$, $b = a + 1\,(a > 0)$, $f_1(t) = \frac{1}{t}$, $f_2(t) = -\ln t$, we obtain:

$$\frac{2a+2}{2a+1}\, e^{\frac{1}{6(2a+1)^2}} < \frac{e}{\left(1+\frac{1}{a}\right)^a} < \sqrt{1+\frac{1}{a}} \cdot e^{\frac{1}{3(2a+1)^2}}. \tag{6}$$

This inequality implies for $a > 0$ e.g. that

$$e^{\frac{1}{2a}\left(1-\frac{1}{a}\right)} < \frac{e}{\left(1+\frac{1}{a}\right)^a} < e^{\frac{1}{2a}\left(1-\frac{1}{2a}\right)} \tag{7}$$

and so

$$A_n = \left(\frac{1}{2} - n \ln e \Big/ \left(1+\frac{1}{n}\right)^n\right) = 0(1/n)$$

which can be compared with the more familiar $\lim\limits_{n\to\infty} A_n = 0$.

3) Apply (1), (2) for $f(t) = \frac{1}{t}$ to deduce

$$\sqrt{ab} < L(a,b) < \frac{a+b}{2}, \tag{8}$$

Where $L(a,b) = \dfrac{b-a}{\ln b - \ln a}$ denotes the logarithmic means (see [2], [3]). The right-hand side of this inequality is due to B. Ostle and H. L. Terwilliger [6]. The left-hand inequality was stated by B. C. Carlson [2]. (8) was rediscovered also by A. Lupaş [5].

4) Select $f(t) = -\ln t$ in (2). This application yields the following improvement of the right-hand side of (8):

$$L(a,b) < \left(\frac{a+b}{2} + \sqrt{ab}\right)/2. \tag{9}$$

5) An interesting remark is that one can use (8) (and also (9)) to obtain refinements of this inequality. Indeed, let us consider $a = \sqrt{x}$, $b = \sqrt{y}$ in (8). It follows that

$$\sqrt{xy} < \sqrt[4]{xy}\left(\frac{\sqrt{x}+\sqrt{y}}{2}\right) < L(x,y) < \left(\frac{\sqrt{x}+\sqrt{y}}{2}\right)^2 < \frac{x+y}{2}. \tag{10}$$

With the same argument we can derive (on base of (9)):

$$L(x,y) < \frac{1}{2}\left(\frac{\sqrt{x}+\sqrt{y}}{2}\right)^2 + \frac{1}{4}(\sqrt{x}+\sqrt{y})\sqrt[4]{xy}. \tag{11}$$

6) In order to arrive to a better refinement, we can consider the relation (3) for $f(t) = 1/t$, $k = 2 (0 < a < b)$. It results $L(a, b) < \frac{3}{8}(a + b)^3/(a^2 + ab + b^2)$. Letting $a = \sqrt[3]{x}$, $b = \sqrt[3]{y}$, this is just one of the Lin [3] and Rüthing [8] inequalities:

$$L(x, y) < \left(\frac{\sqrt[3]{x} + \sqrt[3]{y}}{2}\right)^3 \tag{12}$$

The author wishes to thank the referee for his valuable suggestions.

Joseph Sándor,
Jud. Harghita, Romania

REFERENCES

1 Beckenbach E. F.: Convex functions. Bull. Amer. Math. Soc. *54*, 439–460 (1948).
2 Carlson B. C.: Some inequalities for hypergeometric functions. Proc. Amer. Math. Soc. *17*, 32–39 (1966).
3 Lin T. P.: The power mean and the logarithmic mean. Amer. Math. Monthly *81*, 879–883 (1974).
4 Love E. R.: Some logarithm inequalities. Math. Gaz. *64*, 55–57 (1980).
5 Lupaş A.: Problem 12739. Gaz. Mat. (Bucureşti), Number 2, 1973.
6 Ostle B. and Terwilliger H. L.: A comparison of two means. Proc. Montana Acad. Sci. *17*, 69–70 (1957).
7 Pólya G. and Szegö G.: Aufgaben und Lehrsätze aus der Analysis, vol. I, Ch. 4, Springer 1924.
8 Rüthing D.: Eine allgemeine logarithmische Ungleichung. El. Math. *41*, 14–16 (1986).
9 Sándor J.: On Hadamard's inequality (Hungarian). Mat. Lapok (Cluj) *87*, 427–430 (1982).

© 1988 Birkhäuser Verlag, Basel 0013-6018/88/060177-04$1.50+0.20/0

Kleine Mitteilungen

Eine komplexe Ungleichung aus elementarer Sicht

Aus der Operatorentheorie ist folgende Ungleichung bekannt:

Für $f(z_1, z_2) = 1 + 2(z_1 + z_2) + (z_1 - z_2)^2$ mit $z_1, z_2 \in C$ und $|z_1| = |z_2| = 1$ gilt

$$|f(z_1, z_2)| \leq 5. \tag{1}$$

Man kann (1) mit Hilfe der Ableitungen beweisen, indem man $z_1 = \exp(i\varphi_1)$, $z_2 = \exp(i\varphi_2)$ setzt und dann die Paare φ_1, φ_2 sucht, für die

$$\frac{\partial}{\partial\varphi_1}|f(z_1, z_2)|^2 = \frac{\partial}{\partial\varphi_2}|f(z_1, z_2)|^2 = 0$$