

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 44 (1989)
Heft: 1

Artikel: Some inequalities for the triangle
Autor: Corach, G. / McGowan, J. / Porta, H.
DOI: <https://doi.org/10.5169/seals-41603>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ELEMENTE DER MATHEMATIK

Revue de mathématiques élémentaires – Rivista di matematica elementare

*Zeitschrift zur Pflege der Mathematik
und zur Förderung des mathematisch-physikalischen Unterrichts*

El. Math.

Vol. 44

Nr. 1

Seiten 1–24

Basel, Januar 1989

Some inequalities for the triangle

Inequalities involving the average side, the average altitude, and the diameter of the circumscribed circle of a triangle

I. Introduction

Let a, b, c be the sides of a triangle inscribed in a circle of diameter D . Denote by l and h the averages of the sides $l = (a + b + c)/3$ and the altitudes $h = (h_a + h_b + h_c)/3$, respectively. The quotients h/l and D/l satisfy the following sharp inequalities (with equality for the equilateral triangle):

$$0 < h/l \leq \sqrt{3}/2 \quad (1)$$

$$2/\sqrt{3} \leq D/l < +\infty. \quad (2)$$

The first inequality is due to Santaló [S] and the second to Nakajima [N] and Padoa [P] (see also [B], 6.1 and 5.3). Inequality (1) was also proposed as a Monthly problem (E 1427, page 692, vol 67, 1960; solution on page 296, vol 68, 1961). Apparently none of the 19 solutions received mentioned [S], which had appeared 17 years earlier.

Averaging the inequalities in (1) and (2) gives

$$1/\sqrt{3} \leq \frac{1}{2}h/l + \frac{1}{2}D/l < +\infty, \quad (3)$$

but the sharp lower bound 1 is not hard to obtain (see [C]). The objective of this paper is to study such inequalities.

Two questions arise naturally: (a) what happens when other convex combinations of h/l and D/l are considered; and (b) what happens when only acute triangles are considered? We obtain sharp inequalities for all combined cases of (a) and (b) (in particular we prove the upper bound for (3) in the case of acute triangles conjectured in [C]).

More precisely, for $0 < \theta < 1$ denote by g_θ and G_θ the infimum and supremum of $E_\theta = \theta h/l + (1 - \theta)D/l$ taken over all triangles and by g'_θ and G'_θ the infimum and supremum taken over all acute triangles. Our main result is the following theorem whose proof is given below.

4. Theorem. *With θ_4 and θ_{10} given below, we have:*

$$\begin{aligned} \text{If } \frac{3}{4} \leq \theta < 1 \text{ then } g_\theta = \sqrt{3} \sqrt{\theta(1-\theta)}; \\ \text{if } \frac{1}{2} \leq \theta \leq \frac{3}{4} \text{ then } g_\theta = \frac{3}{2} - \theta; \\ \text{if } \theta_{10} \leq \theta \leq \frac{1}{2} \text{ then } g_\theta = \theta(3u+2)\sqrt{(1-u)/(1+u)} \end{aligned} \quad (4.1)$$

where u is the smaller root of

$$\begin{aligned} (u^2 - 1)(2u + 3) + 3(1 - \theta)/\theta = 0 \quad \text{in } 0 < u < 1; \\ \text{if } 0 < \theta \leq \theta_{10} \text{ then } g_\theta = (\sqrt{3}/6)(4 - \theta). \end{aligned}$$

Only when $0 < \theta < \frac{1}{2}$ is g_θ attained (at the equilateral triangle for $0 < \theta < \theta_{10}$ and at the isosceles triangle with angle opposite the base equal to $\pi - 2 \arccos(u)$ for $\theta_{10} \leq \theta < \frac{1}{2}$).

$$\begin{aligned} \text{If } \frac{3}{4} \leq \theta < 1 \text{ then } g'_\theta = \frac{3}{2} - \theta; \\ \text{if } 0 < \theta \leq \frac{3}{4} \text{ then } g'_\theta = g_\theta. \end{aligned} \quad (4.2)$$

It is attained only for $0 < \theta < \frac{1}{2}$.

$$\begin{aligned} \text{If } \theta_4 \leq \theta < 1 \text{ then } G'_\theta = \frac{\sqrt{3}}{6}(4 - \theta); \\ \text{if } \frac{3}{23}(5 - \sqrt{2}) \leq \theta \leq \theta_4 \text{ then } G'_\theta = 3(-1 + \sqrt{2}) + \frac{1}{2}(9 - 7\sqrt{2})\theta; \\ \text{if } 0 < \theta \leq \frac{3}{23}(5 - \sqrt{2}) \text{ then } G'_\theta = \frac{3}{2} - \theta. \end{aligned} \quad (4.3)$$

Only when $\frac{3}{23}(5 - \sqrt{2}) \leq \theta \leq 1$ is the supremum attained (at the equilateral triangle for $\theta_4 \leq \theta < 1$ and at the right isosceles triangle for $\frac{3}{23}(5 - \sqrt{2}) \leq \theta \leq \theta_4$).

$$G_\theta = +\infty \quad \text{for } 0 < \theta < 1. \quad (4.4)$$

In the above

$$\begin{aligned} \theta_4 &= (1/382)(508 + 153\sqrt{2} - 138\sqrt{3} - 113\sqrt{6}) = 0.5460, \\ \theta_{10} &= 3/(3 + K_{10}) = 0.4799, \end{aligned}$$

where $K_{10} = 3 + 2u_0 - 3u_0^2 - 2u_0^3 = 3.2512$, with

$$A = \frac{1}{12}(22,429 + 243\sqrt{5793})^{1/3} + \frac{1}{12}(22,429 - 243\sqrt{5793})^{1/3} + \frac{19}{48}$$

and

$$u_0 = \frac{1}{2} \left(\sqrt{A} + \sqrt{\frac{297}{32\sqrt{A}} + \frac{19}{16} - A} \right) - \frac{11}{8} = 0.1801.$$

We close this section with some particularly attractive inequalities obtained from Theorem 4 by assigning special values to θ . In all cases the upper bounds hold for acute triangles, and they are sharp. The lower bounds hold for all triangles, and they are sharp for both acute or arbitrary triangles.

$$2l \leq h + D \leq (1/2)(-3 + 5\sqrt{2}) \cdot l$$

$$(5/2)\sqrt{3} \cdot l \leq h + 3D \leq 5 \cdot l$$

$$3l \leq 3h + D \leq (13/6)\sqrt{3} \cdot l$$

$$7l \leq 5h + 3D \leq (9/2)\sqrt{3} \cdot l$$

$$(\sqrt{3}/6)(7 + 4\sqrt{2}) \cdot l \leq h + (1 + \sqrt{2})D \leq (1/2)(4 + 3\sqrt{2}) \cdot l$$

$$(1/6)\sqrt{3}(-3 + 7\sqrt{2}) \cdot l \leq (\sqrt{2} - 1)h + \sqrt{2}D \leq (1/2)(-1 + 4\sqrt{2}) \cdot l$$

$$405\sqrt{5/7} \cdot l \leq 162h + 175D \leq (-282 + 444\sqrt{2}) \cdot l$$

$$608\sqrt{7} \cdot l \leq 768h + 819D \leq (-1305 + 2073\sqrt{2}) \cdot l$$

$$(1725/\sqrt{11}) \cdot l \leq 250h + 264D \leq (-417 + 667\sqrt{2}) \cdot l.$$

The corresponding values of θ are: $1/2, 1/4, 3/4, 5/8, 1 - (\sqrt{2}/2), (1/7)(3 - \sqrt{2}), 162/337, (16/23)^2$, and $125/257$, respectively. Such nice expressions can not be expected for all values of θ , of course. In particular if $\theta = 12/25$ for example (so $K = 13/4$), given any triangle T one can construct with straightedge and compass the number $E_\theta(T)$ but it is impossible to construct the lower bound $g_{12/25}$ or the triangle where it is attained.

II. Reduction to special cases

Since h/l and D/l are invariant under dilations we will assume in the rest of this paper that $D = 1$. Then if α, β, γ are the angles opposite sides a, b, c we have $a = \sin \alpha, b = \sin \beta, c = \sin \gamma, h_a = b c, h_b = c a, h_c = a b$. Abbreviate

$$\begin{aligned} P &= a + b + c = \sin \alpha + \sin \beta + \sin \gamma \\ Q &= a b + b c + c a = \sin \alpha \sin \beta + \sin \beta \sin \gamma + \sin \gamma \sin \alpha \end{aligned} \tag{5}$$

and introduce the convenient parameter $K = 3(1 - \theta)/\theta$ (so that $\theta = 3/(3 + K)$); we will frequently use K or a combination of both K and θ in lieu of θ . Then $E_\theta = \theta(Q + K)/P$ and our problem is to find the extrema of E_θ as a function of α, β, γ subject to

$$\alpha, \beta, \gamma \geq 0, \quad \alpha + \beta + \gamma = \pi \tag{6.a}$$

for the case of all triangles or

$$\frac{\pi}{2} \geq \alpha, \beta, \gamma \geq 0, \quad \alpha + \beta + \gamma = \pi \quad (6.b)$$

for acute triangles.

7. Proposition. *The interior critical points of E_θ subject to (6.a) (or (6.b)) satisfy $\alpha = \beta$, $\beta = \gamma$, or $\gamma = \alpha$, i.e., they correspond to isosceles triangles.*

Proof: Apply Lagrange multipliers as follows. Denote $f(\alpha, \beta, \gamma) = \alpha + \beta + \gamma - \pi$. Using (5) we get

$$\frac{\partial}{\partial \alpha} (E_\theta + \lambda f) = \frac{\theta}{P^2} \{ (P^2 - Q - K) \cos \alpha - P \sin \alpha \cos \alpha + \lambda P^2 \}$$

and similar expressions for the other partial derivatives. Thus the simultaneous vanishing of $\partial(E_\theta + \lambda f)/\partial \alpha$, $\partial(E_\theta + \lambda f)/\partial \beta$, and $\partial(E_\theta + \lambda f)/\partial \gamma$, is equivalent to a system of the form

$$L \cos \alpha + M \sin \alpha \cos \alpha + N = 0$$

$$L \cos \beta + M \sin \beta \cos \beta + N = 0$$

$$L \cos \gamma + M \sin \gamma \cos \gamma + N = 0$$

(with $M = P \neq 0$ for interior points, etc.). Therefore the critical points are zeros of

$$\delta(\alpha, \beta, \gamma) = \det \begin{pmatrix} \cos \alpha & \sin \alpha \cos \alpha & 1 \\ \cos \beta & \sin \beta \cos \beta & 1 \\ \cos \gamma & \sin \gamma \cos \gamma & 1 \end{pmatrix}.$$

From the assumptions $0 < \alpha, \beta, \gamma, \alpha + \beta + \gamma = \pi$ one can evaluate $\delta(\alpha, \beta, \gamma)$ (see VI):

$$\delta(\alpha, \beta, \gamma) = 4 \sin \frac{\alpha - \beta}{2} \sin \frac{\beta - \gamma}{2} \sin \frac{\gamma - \alpha}{2} \left(1 + 2 \sin \frac{\alpha}{2} \cdot \sin \frac{\beta}{2} \cdot \sin \frac{\gamma}{2} \right)$$

and since the last factor cannot vanish (in fact, ≥ 2 by 2.16 in [B]), two of α, β, γ must agree, and the proof of Proposition 7 is complete.

This reduces the calculations to isosceles and degenerate triangles (or isosceles, degenerate and right triangles if we restrict ourselves to acute triangles). In fact interpreting the simplex $\alpha + \beta + \gamma = \pi$, $0 \leq \alpha, \beta, \gamma$ as an equilateral triangle (see Figure 1) we conclude from the proposition above that in order to minimize or maximize E_θ over all triangles it suffices to do so over the intervals $V_1'' V_1'$ and $V_1'' V_2'$ (the remainder of the boundary simply repeats congruent copies of degenerate triangles already contained in $V_1'' V_2'$). For acute triangles it suffices to consider RV_1' and RV_2' , for analogous reasons. Thus from now on only critical points of functions of one variable are considered.




Figure 1.

To avoid repetition we use V' to denote the degenerate triangle with two right angles, and E and R to denote the equilateral and the right isosceles triangles, respectively. In Figure 1 congruent copies of V' appear three times as V'_1 , V'_2 and V'_3 . We have

$$E_\theta(V') = \frac{3}{2} - \theta$$

$$E_\theta(E) = \frac{2\sqrt{3}}{3} - \frac{\sqrt{3}}{6}\theta$$

$$E_\theta(R) = 3(-1 + \sqrt{2}) + \frac{1}{2}(9 - 7\sqrt{2})\theta.$$

We will also introduce thirteen constants K_1, \dots, K_{13} whose values are given in Table 1.

III. Critical points

In this section we determine the extrema of E_θ on each of the relevant segments of Figure 1.

III.1. Case of $V''_1 V''_2$

These degenerate triangles have $\beta = 0$, $0 \leq \gamma \leq \pi/2$, $\alpha = \pi - \gamma$ so that

$$E_\theta = \theta(\sin^2 \gamma + K)/2 \sin \gamma.$$

The critical point W' on $V''_1 V''_2$ is characterized by that value of γ at which

$$\frac{dE_\theta}{d\gamma} = \frac{\theta \cos \gamma}{2 \sin^2 \gamma} (\sin^2 \gamma - K) = 0$$

so that $K = \sin^2 \gamma$ is between 0 and 1. Also $E_\theta(W') = \theta \sqrt{K}$, so it is easy to compare the values of E_θ at the endpoints $V_1''(E_\theta(V_1'') = +\infty)$, $V_2'(E_\theta(V_2') = E_\theta(V'))$ and $E_\theta(W')$.

III.2. Case of RV_2'

These are right triangles with $\alpha = \pi/2$, $0 \leq \beta \leq \pi/4$, $\gamma = \pi/2 - \beta$. Setting $y = 1 + \sin \beta + \cos \beta$ we have $E_\theta = y \theta/2 + \theta(K - 1)/y$ and therefore the only critical points $dE_\theta/d\beta = 0$ satisfy $y^2 = 2(K - 1)$ or $dy/d\beta = 0$. Now $dy/d\beta = 0$ corresponds to R and for the triangle R_1 corresponding to the other solution we have $E_\theta(R_1) = \theta y$. Since $2 \leq y \leq 1 + \sqrt{2}$ there is an internal ($0 < \beta < \pi/2$) critical point only for $K_5 \leq K \leq K_{13}$. Calculating $dE_\theta/d\beta$ between $\beta = 0(V_2')$ and $\beta = \pi/4(R)$ we see easily that for $K_5 < K < K_{13}$ we have $dE_\theta/d\beta < 0$ at $\beta = 0$ and $dE_\theta/d\beta > 0$ at $\beta = \pi/4 - \varepsilon$ for $\varepsilon > 0$ sufficiently small. This shows that $E(R_1) \leq E(V_2')$ and $E(R_1) \leq E(R)$.

III.3. Case of $V_1'' V_1'$

These are isosceles triangles with $\beta = \gamma$. Introduce the parameter $u = \sqrt{1 - b^2} = \sqrt{1 - c^2} = \cos \beta = \cos \gamma$ so that $\alpha = 2u\sqrt{1 - u^2}$, and

$$E_\theta(a, b, c) = E_\theta(u) = \theta \frac{(1 + 4u)(1 - u^2) + K}{2(1 + u)\sqrt{1 - u^2}}.$$

The triangles V_1' , E , R and V_1'' correspond to $u = 0, 1/2, \sqrt{2}/2$ and 1, respectively. Also abbreviate

$$\eta(x) = (1 - x^2)(1 + 4x),$$

$$\zeta(x) = (1 + x)\sqrt{1 - x^2},$$

$$\psi(x) = (1 - x^2)(3 + 2x),$$

$$H(x, y) = (\eta(x) + y)/\zeta(x),$$

$$\phi(x) = 2(2 + 3x)\sqrt{\frac{1 - x}{1 + x}} = H(x, \psi(x)).$$

Then $E_\theta(u) = (\theta/2)H(u, K)$ and

$$\frac{dE_\theta}{du} = \frac{\theta}{2} \frac{(1 - 2u)(\psi(u) - K)}{\zeta(u)(1 - u^2)}.$$

Thus the critical points of $E_\theta(u)$ are given by $u = 1/2$ and any roots of

$$\psi(u) = K \tag{8}$$




Figure 2.

in the interval $0 \leq u < 1$ for general triangles, $0 \leq u < \sqrt{2}/2$ for acute triangles. (Note that by symmetry the directional derivative of E_θ perpendicular to $V_1'' V_1'$ vanishes so that every root of (8) is indeed an interior critical point.)

Observing Figure 2 we obtain: For $K > K_{11}$ there are no roots of (8) in $(0, 1)$. For $3 = K_5 < K < K_{11}$ there are two roots

$$0 < u_2 = u_2(K) < \frac{1}{6}(-3 + \sqrt{21}) < u_1 = u_1(K) < 1/2$$

and for $0 < K < K_5$ there is only one root $1/2 < u_1 < 1$. Furthermore, in the last case, we have $\sqrt{2}/2 > u_1$ (i.e., u_1 represents an acute triangle) exactly when $K_3 < K$. We will denote by I_1 and I_2 the triangles corresponding to u_1 and u_2 , respectively.

From the calculation

$$\psi'(x) = 2(2 + 3x)(1 - x^2)\phi'(x)/\phi(x)$$

and the relation $K = \psi(u_i(K))$ for $i = 1, 2$ we get

$$\begin{aligned} \frac{d}{dK} \log \phi(u_i) &= \frac{\phi'(u_i)}{\phi(u_i)} \frac{du_i}{dK} \\ &= \frac{\psi'(u_i)}{2(2 + 3u_i)(1 - u_i^2)} \frac{du_i}{dK} \\ &= \frac{1}{2(2 + 3u_i)(1 - u_i^2)} \\ &= \frac{1}{2K} \frac{3 + 2u_i}{2 + 3u_i}. \end{aligned}$$

We draw two conclusions from this formula. First,

$$\frac{d}{dK} \log \frac{\phi(u_2)}{\phi(u_1)} = \frac{5}{2K} \frac{u_1 - u_2}{(2 + 3u_1)(2 + 3u_2)} > 0.$$

Second,

$$\begin{aligned} \frac{d}{dK} \log E_\theta(I_i) &= \frac{d}{dK} \log (\theta \phi(u_i)) \\ &= \frac{\theta'}{\theta} + \frac{1}{2K} \frac{3 + 2u_i}{2 + 3u_i} \\ &= -\frac{1}{K + 3} + \frac{1}{2K} \frac{3 + 2u_i}{2 + 3u_i}, \end{aligned}$$

which is positive when $K = 3$ for both $u_1 = 1/2$ and $u_2 = 0$. Using $\psi(u_i) = K$, this derivative could only vanish if u_i were a root of $8x^4 + 14x^3 - 5x^2 - 8x + 6 = (2x + 3)(4x^3 + x^2 - 4x + 2) = 0$. But this equation has no positive root. Hence from $E_\theta(u_2)/E_\theta(u_1) = \phi(u_2)/\phi(u_1)$ and the foregoing we get:

III.4. $E_\theta(I_1)$, $E_\theta(I_2)$, and the ratio $E_\theta(I_2)/E_\theta(I_1)$, are strictly increasing functions of K . Since $E_\theta(I_2) = E_\theta(I_1)$ for $K = K_{11}$ it follows that $E_\theta(I_2) < E_\theta(I_1)$ for $3 < K < K_{11}$.

IV. Intersections

According to sections II and III the extrema of E_θ are achieved for general triangles at one of E, V', V'', W', I_1, I_2 and for acute triangles at one of R, E, V', R_1, I_2 and I_1 (if I_1 corresponds to an acute triangle). To determine which points correspond to extrema we consider the seven functions $f^T(K) = E_\theta(T)$ of K where, T is one of $V', W', R, R_1, E, I_1, I_2$ and $\theta = 3/(3 + K)$. Our strategy is a brute force approach: First we determine the values of K at which each pair of functions f^T and f^S intersect by solving

$$f^T(K) = f^S(K) \tag{9}$$

and then we rank them in the resulting contiguous intervals. All the information is summarized in Figure 3 at the end of this section and Tables 1 and 2.

IV.1. $T, S \in \{R, E, V'\}$. These are isosceles triangles with u_T and u_S among $u_R = \sqrt{2}/2$, $u_E = 1/2$, $u_{V'} = 0$ so the intersections occur at the roots of

$$\theta \frac{(1 + 4u_T)(1 - u_T^2) + K}{2(1 + u_T)(1 - u_T^2)^{1/2}} = \theta \frac{(1 + 4u_S)(1 - u_S^2) + K}{2(1 + u_S)(1 - u_S^2)^{1/2}}$$

which is linear in K . The solutions K_4, K_6, K_{12} are listed in Table 1.

IV.2. $T \in \{I_1, I_2\}$, $S \in \{R, E, V'\}$. Equation (9) reads

$$\frac{\theta \eta(u) + \psi(u)}{2 \zeta(u)} = \theta \frac{(1 + 4u_S)(1 - u_S^2) + K}{2(1 + u_S)(1 - u_S^2)^{1/2}}$$

where $K = \psi(u)$. Upon squaring this yields a seventh order polynomial in u . The roots r in $((-3 + \sqrt{21})/6, 1)$ correspond to $T = I_1$ and the roots s in $(0, (-3 + \sqrt{21})/6)$ correspond to $T = I_2$.

IV.2a. When $S = R$ the equation obtained is

$$(u - 1/\sqrt{2})^2 (8u^5 + 8(4 + \sqrt{2})u^4 + 2(19 + 16\sqrt{2})u^3 + 2(-1 + 13\sqrt{2})u^2 + (1 - 2\sqrt{2})u + 9 - 4\sqrt{2}) = 0$$

where the fifth order factor has no positive roots and the root $r = 1/\sqrt{2}$ corresponds to the intersection of f^{I_1} and f^R at K_3 .

IV.2b. When $S = E$ the equation is $(u - 1/2)^3 (8u^4 + 44u^3 + 86u^2 + 33u - 9) = 0$ and $r = 1/2 > (-3 + \sqrt{21})/6$ gives the intersection of f^{I_1} and f^E at $K_5 = \psi(1/2) = 3$. The quartic factor has one positive root $u_0 = 0.180125573$ whose closed form expression is given in the statement of Theorem 4. It corresponds to the intersection of f^{I_2} and f^E at K_{10} .

IV.2c. When $S = V'$ the equation obtained is $u^2(4u^5 + 16u^4 + 13u^3 - 27u^2 - 12u + 8) = 0$ and the second factor has two positive roots: $r_1 = 0.961103259$ and $r_2 = 0.407160288$ (both are $> (-3 + \sqrt{21})/6$). They correspond to the intersections of f^{I_1} and $f^{V'}$ at $K_1 = \psi(r_1)$ and $K_7 = \psi(r_2)$. The root $u=0$ corresponds to the intersection of $f^{I_2}(K)$ and $f^{V'}(K)$ at $K_5 = \psi(0) = 3$.

IV.3. $T = I_1$, $S = I_2$. As stated in III.4 these intersect only once, at K_{11} .

IV.4. $T = V'$, $S = R_1$. Equation (9) is $3/2 - 3/(3 + K) = (3/(3 + K))\sqrt{2K - 2}$ so $K = K_5 = 3$.

IV.5. $T = E$, $S = R_1$. Recall that f^{R_1} is only defined for $3 \leq K \leq 5/2 + \sqrt{2}$. The equation (9) is $2\sqrt{3}/3 - (\sqrt{3}/6)(3/(3 + K)) = (3/(3 + K))\sqrt{2K - 2}$, or $16K^2 - 144K + 297 = 0$, and $K_8 = (18 - 3\sqrt{3})/4$ is the only root in the range.

IV.6. $T = R$, $S = R_1$. Equation (9) is $3(-1 + \sqrt{2}) + \frac{1}{2}(9 - 7\sqrt{2})(3/(3 + K)) = (3/(3 + K))\sqrt{2K - 2}$ or $K = K_{13} = 5/2 + \sqrt{2}$.

IV.7. $T = R_1$, $S \in \{I_1, I_2\}$. The equation is

$$\frac{\theta \eta(u) + \psi(u)}{2 \zeta(u)} = \theta \sqrt{2\psi(u) - 2}$$

where $K = \psi(u)$. Thus $u^2(4u^2 + u - 1) = 0$, and the positive root $r = (-1 + \sqrt{17})/8$ corresponds to the intersection of f^{I_1} and f^{R_1} at $K = K_9 = \psi(r) = (135 + 17\sqrt{17})/64$. The root $u = 0$ corresponds to the intersection of f^{I_2} and f^{R_1} at $K_5 = 3 = \psi(0)$.

IV.8. $T = V'$, $S = W'$. The equation is $3/2 - 3/(3+K) = (3/(3+K))\sqrt{K}$ so $K = K_2 = 1$.

IV.9. $T \in \{E, R\}$, $S = W'$. Equation (9) leads to quadratic equations $(16K^2 - 36K + 81 = 0$ for $T = E$, $4K^2 - 8K + 9 + 4\sqrt{2} = 0$ for $T = R$) with no real roots.

IV.10. $T \in \{I_1, I_2\}$, $S = W'$. Equation (9) reads

$$\theta(\eta(u) + \psi(u))/2\zeta(u) = \theta\sqrt{\psi(u)}$$

or $(u - 1)(2u^3 - 2u^2 - 4u - 1) = 0$, with no root in $(0, 1)$.

IV.11. $T = R_1$, $S = W'$. The corresponding functions have disjoint domains so there is no intersection.

The proof of Theorem 4 results now from inspections of Tables 1, 2, 3, and 4.




Figure 3. Abstract sketch of the ranks and intersections of $f^T(K)$.

V. Miscellaneous remarks

V.1. In Theorem 4 we gave a “closed form” radical expression for K_{10} . This is possible for all K_i except K_1 and K_7 . In fact the polynomial $\tau(u) = 4u^5 + 16u^4 + 13u^3 - 27u^2 - 12u + 8$ is irreducible over the rationals and has precisely two non-real roots. Thus its Galois group is S_5 (the symmetric group on five elements) (see [K], Theorem 29). If $K = K_1$ or K_7 , then for some root u of τ , $K = \psi(u) \in \mathbb{Q}(u) \setminus \mathbb{Q}$. As $[\mathbb{Q}(u):\mathbb{Q}] = 5$, we have $\mathbb{Q}(K) = \mathbb{Q}(u)$. Thus the Galois group of the minimal polynomial for K is S_5 as well ($= \text{Aut}_{\mathbb{Q}} N$, where N is the normal closure of $\mathbb{Q}(K)$ over \mathbb{Q}). Hence K can not lie in a radical extension of \mathbb{Q} (by Theorem 28 in [K]).

Table 1. Values of K_i and θ_i .

$K_0 = 0.0000$	$= 0$	$\theta_0 = 1.0000$
$K_1 = 0.3755$	$= \psi(u)[1^*]$ where u is the <i>larger</i> root of $r(u)[2^*]$ in $(0, 1)$ ($u = 0.9611$)	$\theta_1 = 0.8888$
$K_2 = 1.0000$	$= 1$	$\theta_2 = 0.7500$
$K_3 = 2.2071$	$= (3 + \sqrt{2})/2$	$\theta_3 = 0.5761$
$K_4 = 2.4948$	$= (3/194)(39 - 18\sqrt{2} + 33\sqrt{3} + 37\sqrt{6})$	$\theta_4 = 0.5460$
$K_5 = 3.0000$	$= 3$	$\theta_5 = 0.5000$
$K_6 = 3.1801$	$= (3/11)(3 + 5\sqrt{3})$	$\theta_6 = 0.4854$
$K_7 = 3.1820$	$= \psi(u)$ where u is the <i>smaller</i> root of $r(u)$ in $(0, 1)$ ($u = 0.4072$)	$\theta_7 = 0.4853$
$K_8 = 3.2010$	$= (1/4)(18 - 3\sqrt{3})$	$\theta_8 = 0.4838$
$K_9 = 3.2046$	$= (1/64)(135 + 17\sqrt{17})$	$\theta_9 = 0.4835$
$K_{10} = 3.2512$	$= \psi(u_0)$ where u_0 is the root of $\sigma(u)[3^*]$ in $(0, 1)$ ($u_0 = 0.1801[4^*]$)	$\theta_{10} = 0.4799$
$K_{11} = 3.2821$	$= (1/18)(27 + 7\sqrt{21})$	$\theta_{11} = 0.4775$
$K_{12} = 3.4142$	$= 2 + \sqrt{2}$	$\theta_{12} = 0.4677$
$K_{13} = 3.9142$	$= (1/2)(5 + 2\sqrt{2})$	$\theta_{13} = 0.4339$
$K_\infty = \infty$		$\theta_\infty = 0$

[1*] $\psi(u) = -2u^3 - 3u^2 + 2u + 3$,

[2*] $r(u) = 4u^5 + 16u^4 + 13u^3 - 27u^2 - 12u + 8$,

[3*] $\sigma(u) = 8u^4 + 44u^3 + 86u^2 + 33u - 9$,

[4*] the exact value of u_0 is given in Theorem 4.

Table 2. Values of K at which $f^T(K)$ and $f^S(K)$ intersect. Table entries are the subscripts from Table 1 (e.g., f^{I_1} and $f^{V'}$ intersect at $K = K_1$ and $K = K_7$).

	V'	I_2	I_1	E	R	R_1	W'
V'	—	5	1.7	6	12	5	2
I_2	—	—	11	10	[1*]	5	[1*]
I_1	—	—	—	5	3[2*]	9	[1*]
E	—	—	—	—	4	8	[1*]
R	—	—	—	—	—	13	[1*]
R_1	—	—	—	—	—	—	[1*]
W'	—	—	—	—	—	—	—

[1*] no intersection,

[2*] intersect, do not cross.

Table 3. (All triangles.) Orderings of $E_\theta(T)$, for $T \in \{W', V', I_1, E_2, E\}$ and $K_i < K < K_j$. The critical point of lowest E_θ appears on the left; E_θ increases to the right.

i, j				
0, 1	W'	I_1	V'	E
1, 2	W'	V'	I_1	E
2, 5		V'	I_1	E
5, 6	I_2	V'	E	I_1
6, 7	I_2	E	V'	I_1
7, 10	I_2	E	I_1	V'
10, 11	E	I_2	I_1	V'
11, ∞	E			V'

Table 4. (Acute triangles.) Orderings of $E_\theta(T)$ for $T \in \{I_1, I_2, R, R_1, E, V'\}$ for $K_i < K < K_j$. The critical point with smallest value of E_θ appears on the left; E_θ increases to the right. (Remark: I_1 is obtuse for $K < K_3$).

i, j						
0, 3			V'		R	E
3, 4			V'	I_1	R	E
4, 5			V'	I_1	E	R
5, 6	I_2	R_1	V'	E	I_1	R
6, 7	I_2	R_1	E	V'	I_1	R
7, 8	I_2	R_1	E	I_1	V'	R
8, 9	I_2	E	R_1	I_1	V'	R
9, 10		I_2	E	I_1	R_1	V'
10, 11		E	I_2	I_1	R_1	V'
11, 12		E			R_1	V'
12, 13		E			R_1	R
13, ∞		E			R	V'

Setting $\sigma(u) = 8u^4 + 44u^3 + 86u^2 + 33u - 9$, note that the cubic resolvent of σ is the irreducible cubic with two non-real roots

$$(4y)^3 - 43(4y)^2 + 435(4y) - 2007$$

with Galois group S_3 (the symmetric group on three elements) (see [K], Theorem 29). The Galois group of $\sigma (= \text{Aut}_Q N$ where N is the normal closure of $Q(u_0)$ where $\sigma(u_0) = 0$) is S_4 (by Theorem 43 in [K]). Since $K_{10} = \psi(u_0) \in Q(u_0) \setminus Q$, $Q(\theta_{10}) = Q(K_{10}) = Q(u_0)$ ([K], Ex. 5, p. 53).

Therefore

- i) θ_{10} , K_{10} and u_0 are of degree 4 over Q (σ is irreducible!).
- ii) The minimal polynomials of θ_{10} , K_{10} and u_0 all have Galois group S_4 .
- iii) None of θ_{10} , K_{10} , u_0 are constructible with straightedge and compass ([K], p. 195, remark at page bottom).

Thus none of the following are constructible with straightedge and compass (over Q):

- i) The minimum relative variation of $E_\theta(T)$ over acute triangles (see V.3):

$$G'_{\theta_{10}}/g'_{\theta_{10}}$$

- ii) The upper (lower) bound of $E_\theta(T)$ at the point of minimum relative variation:

$$G'_{\theta_{10}}, g'_{\theta_{10}}.$$

- iii) The non-equilateral, isosceles triangle with minimum value of E_θ at the point of minimum relative variation (angle opposite the base equal to $\pi - 2 \arccos u_0$).
- iv) The relative weights for minimum relative variation in E_θ : θ_{10} and $1 - \theta_{10}$.

All of the above are of course constructible given θ_{10} (or K_{10}). For completeness we give the minimal polynomial of θ_{10} over \mathbb{Q} :

$$4871\theta^4 - 5939\theta^3 + 1356\theta^2 + 112\theta + 32.$$

V.2. The conjecture in [C] that for any acute triangle T we have $E_{1/2}(V) \leq E_{1/2}(T) \leq E_{1/2}(R)$ follows from Theorem 4. Observe that it can be written as

$$l \leq \frac{1}{2}(h + D) \leq \frac{-3 + 5\sqrt{2}}{4} \cdot l$$

where $E_{1/2}(R) = (-3 + 5\sqrt{2})/4 = 1.017766953$, i.e.: *in any acute triangle the average of the mean altitude and the circumdiameter is within 2% of the mean side.*

V.3. From Theorem 4 and III.4 we see that G'_θ , g'_θ and g_θ are decreasing functions of θ , so that $\sqrt{3}/2 \leq G'_\theta \leq 3/2$, $1/2 \leq g'_\theta \leq 2/\sqrt{3}$ and $0 < g_\theta \leq 2/\sqrt{3}$ where the lower limits correspond to $\theta = 1$ and the upper limits to $\theta = 0$. It also follows from Theorem 4 that G'_θ/g'_θ is a decreasing function of K for $0 < K \leq K_5$ and increasing for $K \geq K_{10}$. Over the interval $K_5 \leq K \leq K_{10}$, we have $G'_\theta/g'_\theta = E_\theta(R)/E_\theta(I_2) = 2E_\theta(R)/\theta\phi(u_2(K))$ where ϕ and u_2 are as in III.3. The logarithmic derivative

$$\frac{d}{dK} \log \frac{G'_\theta}{g'_\theta} = \frac{2(-1 + \sqrt{2})}{3 - \sqrt{2} + 2(-1 + \sqrt{2})K} - \frac{3 + 2u_2(K)}{2K(3u_2(K) + 2)}$$

can only vanish when (using $K = \psi(u_2)$):

$$(u_2 + 3/2)(u_2 - 1/\sqrt{2})(8u_2^2 + 2(1 + 2\sqrt{2})u_2 - 4 + \sqrt{2}) = 0.$$

However the smallest positive root of this equation is $(\sqrt{41 - 4\sqrt{2}} - (2\sqrt{2} + 1))/8$ which is greater than $(-3 + \sqrt{21})/6$, and therefore it can not be u_2 . Hence the quotient G'_θ/g'_θ is monotone over $K_5 \leq K \leq K_{10}$ and an evaluation of the logarithmic derivative at $K = K_5$ shows it to be a decreasing function of K there. Thus

$$\begin{aligned} \text{for } K \leq K_{10}: G'_{\theta_{10}}/g'_{\theta_{10}} &\leq G'_\theta/g'_\theta \leq \sqrt{3}, \\ \text{for } K \geq K_{10}: G'_{\theta_{10}}/g'_{\theta_{10}} &\leq G'_\theta/g'_\theta \leq 3\sqrt{3}/4, \end{aligned}$$

where the upper limits correspond to $\theta = 1$ and to $\theta = 0$, respectively. The minimum value of G'_θ/g'_θ is then achieved at $K = K_{10}$ with value

$$\frac{18(-1 + \sqrt{2}) + 3(9 - 7\sqrt{2})\theta_{10}}{\sqrt{3}(4 - \theta_{10})} = 1.010471349$$

(where $\theta_{10} = 3/(3 + K_{10})$), or using the values of $g'_{\theta_{10}}$ and $G'_{\theta_{10}}$:

$$1.01616367 \cdot l \leq \theta_{10}h + (1 - \theta_{10})D \leq 1.026804277 \cdot l.$$

VI. Calculation of Determinant

Here is the calculation used in II. of the determinant

$$\delta = \det \begin{pmatrix} 1 & \cos \alpha & \sin \alpha \cos \alpha \\ 1 & \cos \beta & \sin \beta \cos \beta \\ 1 & \cos \gamma & \sin \gamma \cos \gamma \end{pmatrix}.$$

Set $x = e^{i\alpha}$, $y = e^{i\beta}$, $z = e^{i\gamma}$. Then $x\bar{x} = y\bar{y} = z\bar{z} = 1$ and $xyz = -1$. Also substituting in the determinant and using linearity in each column we get $8i\delta = \delta_1 - \delta_2 + \delta_3 - \delta_4$, where

$$\delta_1 = \det \begin{pmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{pmatrix}$$

$$\delta_2 = \det \begin{pmatrix} 1 & x & \bar{x}^2 \\ 1 & y & \bar{y}^2 \\ 1 & z & \bar{z}^2 \end{pmatrix}$$

$$\delta_3 = \det \begin{pmatrix} 1 & \bar{x} & x^2 \\ 1 & \bar{y} & y^2 \\ 1 & \bar{z} & z^2 \end{pmatrix}$$

$$\delta_4 = \det \begin{pmatrix} 1 & \bar{x} & \bar{x}^2 \\ 1 & \bar{y} & \bar{y}^2 \\ 1 & \bar{z} & \bar{z}^2 \end{pmatrix}.$$

Now δ_1 is a Vandermonde determinant so $\delta_1 = (y-x)(z-y)(z-x)$ and using $\bar{x} = 1/x$, etc., and $xyz = -1$ we get

$$\bar{\delta}_1 = (\bar{y}-\bar{x})(\bar{z}-\bar{y})(\bar{z}-\bar{x}) = \frac{x-y}{xy} \cdot \frac{y-z}{yz} \cdot \frac{x-z}{xz} = -\delta_1$$

so δ_1 is purely imaginary. Since $\delta_4 = \bar{\delta}_1$ we have $\delta_1 - \delta_4 = 2\delta_1$.

On the other hand

$$\delta_2 = (xyz)^2 \delta_2 = \det \begin{pmatrix} 1 & x^2 & x^3 \\ 1 & y^2 & y^3 \\ 1 & z^2 & z^3 \end{pmatrix}$$

and by successive subtraction of columns we find $\delta_2 = (xy + yz + zx)(y-x)(z-x)(z-y) = -(\bar{z} + \bar{x} + \bar{y})\delta_1$. Also $\delta_3 = \bar{\delta}_2$ so $-\delta_2 + \delta_3 = 2i\mathfrak{T}((\bar{z} + \bar{x} + \bar{y})\delta_1) = 2\delta_1\Re(\bar{z} + \bar{x} + \bar{y})$. Thus

$4i\delta = (1 + \Re(\bar{z} + \bar{x} + \bar{y}))\delta_1$. Finally observe that

$$y - x = e^{i\beta} - e^{i\alpha} = e^{i(\alpha+\beta)/2} 2i \sin \frac{\beta - \alpha}{2}$$

and similar formulas hold for $z - x$ and $z - y$. Hence

$$\begin{aligned} \delta_1 &= e^{i(\alpha+\beta)/2} e^{i(\alpha+\gamma)/2} e^{i(\gamma+\beta)/2} (2i)^3 \sin \frac{\beta - \alpha}{2} \sin \frac{\gamma - \alpha}{2} \sin \frac{\gamma - \beta}{2} \\ &= 8i \sin \frac{\beta - \alpha}{2} \sin \frac{\gamma - \alpha}{2} \sin \frac{\gamma - \beta}{2}, \end{aligned}$$

and since $\Re(\bar{z} + \bar{x} + \bar{y}) = \cos \alpha + \cos \beta + \cos \gamma$ we get for δ the value

$$\delta = 2 \sin \frac{\beta - \alpha}{2} \sin \frac{\gamma - \alpha}{2} \sin \frac{\gamma - \beta}{2} (1 + \cos \alpha + \cos \beta + \cos \gamma).$$

According to [B, 2.16 and 2.12] this can also be written as

$$\begin{aligned} \delta &= 4 \sin \frac{\beta - \alpha}{2} \sin \frac{\gamma - \alpha}{2} \sin \frac{\gamma - \beta}{2} \left(1 + 2 \sin \frac{\alpha}{2} \cdot \sin \frac{\beta}{2} \cdot \sin \frac{\gamma}{2} \right) \\ &= 4 \sin \frac{\beta - \alpha}{2} \sin \frac{\gamma - \alpha}{2} \sin \frac{\gamma - \beta}{2} (1 + r) \end{aligned}$$

where r is the inradius of the triangle.

G. Corach, I.A.M., Buenos Aires
 J. McGowan, University of Missouri, Columbia, Missouri, USA
 H. Porta, University of Illinois, Urbana, Illinois, USA

REFERENCES

- [B] Bottema O. et al.: Geometric Inequalities. Wolters-Noordhoff, Groningen, 1969.
- [C] Corach G. and Porta H.: Algunas desigualdades geométricas. Revista de Educación 1, 3–12 (1983).
- [K] Kaplansky I.: Fields and Rings (Second Edition). University of Chicago Press, 1972.
- [N] Nakajima S.: Some inequalities between the fundamental elements of the triangle. Tôhoku Math. J. 25, 115–121 (1925).
- [P] Padoa A.: Una questione di minimo. Periodico di Matematiche 5, 80–85 (1925).
- [S] Santaló L.: Algunas desigualdades entre los elementos de un triángulo. Mathematicae Notae 3, 65–73 (1943).