Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 44 (1989)
Heft: 3

Artikel: On cubic polynomials giving many primes
Autor: Goetgheluck, P.
DOI: https://doi.org/10.5169/seals-41611

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 03.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

also modulo $\frac{1}{2}\pi$ reduziert $= 0$, beträgt. Es bleiben die Terme längs $A_0 A_3$ (mit dem Diederwinkel $\frac{1}{2}\pi - \alpha$ und der Kantenlänge $l = \int_{2\sin\alpha}^{u} 1/z\,dz$ zwischen den gelegten Horosphären), und längs $A_0 A_1$ (mit dem Diederwinkel α und der Länge $\int_1^u 1/z\,dz$ ausserhalb der Horosphären). Die elementare Ermittlung der Integrale führt auf

$$\Psi(L(\alpha)) = (\log u - \log|2\sin\alpha|) \otimes (\tfrac{\pi}{2} - \alpha) + (\log u) \otimes \alpha,$$

also nach Reduzierung mod $\frac{1}{2}\pi$ und Addition auf

$$\Psi(L(\alpha)) = \log|2\sin\alpha| \otimes \alpha, \quad \text{w.z.b.w.}$$

Hans E. Debrunner, Math. Institut, Universität Bern

LITERATURVERZEICHNIS

1 Böhm J. und Schwulow H.: Eine Zerlegung von vierdimensionalen euklidischen und nichteuklidischen Simplexen in Orthoscheme. Wiss. Z. Friedrich-Schiller-Univ. Jena Math. Natur. Reihe *31*, 545–555 (1982); MR *84d*: 52008.
2 Böhm J. und Hertel E.: Polyedergeometrie in *n*-dimensionalen Räumen konstanter Krümmung. Birkhäuser, Basel-Boston-Stuttgart 1981; MR *82k*: 52001a.
3 Debrunner H. E.: Dissecting orthoschemes into orthoschemes, im Druck.
4 Dupont J. L. and Sah C.-H.: Scissors congruences, II. J. Pure Appl. Algebra *25*, 159–195 (1982); *30* 217 (1983); MR *84b*: 53062b.
5 Hilbert D. und Cohn-Vossen S.: Anschauliche Geometrie. Springer, Berlin 1932.
6 Lobatschefskij N. I.: Über die Anfangsgründe der Geometrie, Kasaner Bote 1830. Übersetzt und kommentiert von F. Engel, Teubner, Leipzig 1898.
7 Milnor J. W.: Hyperbolic Geometry: the first 150 years. Bull. Amer. Math. Soc. *6*, 9–24 (1982); MR *82m*: 57005.
8 Sah C.-H.: Scissors congruences, I. The Gauss-Bonnet map, Math. Scand *49*, 181–210 (1982); *53*, 62 (1983); MR *84b*: 53062a.
9 Schläfli L.: Theorie der vielfachen Kontinuität, 1852. In: L. Schläfli, Gesammelte Mathematische Abhandlungen, Band I, 177–392, Birkhäuser, Basel 1950.
10 Wythoff W. A.: The rule of Neper in the four dimensional space. K. Akad. Wet. Amsterdam, Proc. Sect. of Sci. *9*, 529–534 (1907).

© 1989 Birkhäuser Verlag, Basel 0013-6018/89/030066-05$1.50 + 0.20/0

On cubic polynomials giving many primes

1. Introduction

In the following «prime» means positive or negative prime, that is an integer of the sequence $\ldots -7, -5, -3, -2, 2, 3, 5, 7, 11 \ldots$

The question of finding quadratic polynomials giving many primes has been widely investigated (see [2, p. 115–117] and [4, p. 141–143]) and we can cite for example $2n^2 - 199$ (Karst, 1973, see [3]) giving 87 primes for $n = 0, 1, \ldots, 99$ and $n^2 + n + 41$ (Euler, 1772) giving 86 primes in the same range.
The purpose of this paper is to exhibit some polynomials $a n^3 + b n^2 + c n + d$ giving many primes and to explain the way they were discovered. In the mathematical literature we found only one result about this subject in [1, p. 420]: «Escott found that $x^3 + x^2 + 17$ is a prime for $-14, -13, \ldots, +10$» (In fact this is not true since making $x = -3$ yields the number -1).
Arbitrarily we shall say that

«$p(n) = a n^3 + b n^2 + c n + d$ gives many primes»

if and only if:

Card $\{n \in [0, 1, 2, \ldots, 99]$ such that $p(n)$ is prime$\} \geq 75$.

2. Method of computation

All calculations were performed with a micro-computer, using PASCAL and assembly language.

First step

We determine all polynomials $p(n) = a n^3 + b n^2 + c n + d$ satisfying:

(i) $a = 1$ or $a = 2$.
(ii) $-1 \leq b \leq 1$ if $a = 1$;
$-2 \leq b \leq 3$ if $a = 2$.
(iii) $|c| \leq 15$.
(iv) $p(n) \not\equiv 0 \pmod{m}$ $(m = 2, 3, 5, 7;\ n \in \mathbb{N})$.
(v) $|d| \leq 10\,000$.
(vi) Card $\{n \in [0, 1, \ldots, 99]$ such that $p(n)$ is prime$\} \geq 60$.

Comments

1) Condition (ii) is not a real restriction since replacing n by $n + 1$ transforms $p(n)$ into $a n^3 + (b + 3a) n^2 + (c + 2b + 3a) n + (d + a + b + c)$ showing that b can be reduced mod $3a$.
2) Without condition (iv) we should have 5,219,739 polynomials to examine, that is 521,973,900 primality tests to perform. With this condition the amount of computation is reduced by a factor 40 and the percentage of interesting polynomials lost is very low.

Practically, for given a, b, and c, to get $d \in [0, \ldots, 209]$ $(210 = 2 \cdot 3 \cdot 5 \cdot 7)$ such that $a n^3 + b n^2 + c n + d$ is never divisible by 2, 3, 5 or 7 we run the following algorithm:

```
For  p = 2, 3, 5 and 7  do
     for  n = 0 to p − 1  do
          for  d = 0 to p − 1  do
               if  (a n^3 + b n^2 + c n + d) mod p = 0
               then for j = d to 209 step p do suppress j.
```

The convenient values of $d \pmod{210}$ are the remaining $j \in [0, \ldots, 209]$.

Example: For $a = 2$, $b = 1$ and $c = -7$, we have only to consider $d = 41$, 107, 167, 191 (mod 210).

Second step

For every polynomial provided by step 1:

- We search in the interval $[-100, \ldots, 100]$ a sequence of 100 consecutive integers j, $j+1, \ldots, j+99$ satisfying.
  Card $\{n \in [j, j+1, \ldots, j+99]$ such that $p(n)$ is prime$\}$ is maximum.
- If the maximum is greater than or equal to 75, changing n into $n+j$ in $p(n)$ yields a new polynomial giving at least 75 primes values for $n = 0, 1, \ldots, 99$.
- If the maximum is less than 75, the polynomial is rejected.

Third step

In the list of polynomials we get from step 2, there are pairs of polynomials $(p(n), q(n))$ such that $p(n) \equiv -q(-n-k)$ for some integer k. Geometrically this means that graphs of p and q are symmetric with respect to a vertical axis. For every pair of such polynomials we cancel one of them.

3. Results

From step 1 we get 215 polynomials. After step 2 it remains 34 polynomials and after step 3, 21 polynomials giving at least 75 prime values for $n = 0, 1, 99$. They are listed below with the number of prime values they have for $0 \leq n < 100$, 200, ..., 500.
Polynomials from step 1 have also been exploited to search the longest sequences of consecutive integers n such that all corresponding values of $p(n)$ are primes:

- Formulas (14) and (15) give 26 primes respectively for $n = 16, \ldots, 41$ and $n = 74, \ldots, 99$;
- Formula $|2n^3 - 83n^2 + 1157n - 4999|$ gives 34 distinct primes for $n = 0, \ldots, 33$.

					Number of primes given by an^3+bn^2+cn+d for				
No	a	b	c	d	$n<100$	$n<200$	$n<300$	$n<400$	$n<500$
(1)	1	−220	16119	−392723	75	134	179	219	261
(2)	1	−199	13190	−290869	75	124	163	206	235
(3)	1	−160	8547	−142811	75	130	179	221	264
(4)	1	−159	8420	−148153	76	124	164	203	238
(5)	1	−151	7610	−129097	76	125	168	204	245
(6)	1	−150	7493	−124277	76	128	170	217	250
(7)	1	−137	6270	−95203	75	121	168	197	233
(8)	1	−125	5196	−73291	79	130	174	212	254
(9)	1	−119	4718	−71741	75	118	158	190	224
(10)	1	−114	4343	−54829	76	125	171	200	232
(11)	1	−111	4100	−49367	76	119	150	199	246
(12)	1	−97	3126	−32603	75	115	161	195	235
(13)	1	−96	3059	−32563	75	127	162	192	225
(14)	1	−82	2237	−20407	75	112	149	183	218
(15)	2	−489	39847	−1084553	75	134	176	222	267
(16)	2	−372	23050	−476027	75	128	174	211	239
(17)	2	−292	14202	−231551	76	124	160	206	252
(18)	2	−289	13917	−221891	75	124	172	210	247
(19)	2	−281	13157	−204487	78	129	169	214	250
(20)	2	−280	13072	−203857	76	123	168	204	240
(21)	2	−199	6595	−79657	79	125	174	217	257

P. Goetgheluck, Université de Paris-sud

REFERENCES

1 Dickson L. E.: History of the theory of numbers, vol. I. Carnegie Inst. Washington, Chelsea, New York 1952.
2 Guy R. K.: Reviews in number theory 1973–1983, vol. I. AMS, Providence RI, 1984.
3 Karst E.: New quadratic forms with high density of primes. Elem. Math. *28*, 116–118 (1973).
4 Leveque W. J.: Reviews in number theory 1940–1972, vol. I. AMS, Providence RI, 1974.

© 1989 Birkhäuser Verlag, Basel 0013-6018/89/030070-04$1.50+0.20/0

A remark on the gamma function

According to Problem 188, Part II of G. Pólya and G. Szegő [2] for each integrable function $f(x)$ on $0 \leqq x \leqq 1$ we have:

$$\lim_{n\to\infty} \frac{1}{\varphi(n)} \sum_{\substack{k=1 \\ (k,n)=1}}^{n} f\left(\frac{k}{n}\right) = \int_0^1 f(x)\,dx,$$