

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 45 (1990)  
**Heft:** 3

**Rubrik:** Kleine Mitteilung

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## LITERATURVERZEICHNIS

- 1 Kamke E.: Differentialgleichungen, Bd. 1, Teubner, Stuttgart 1977, p. 560.  
 2 Wong J. S. W.: On the generalized Emden-Fowler Equation, SIAM Review 17 (2), 339–360 (1975).

© 1990 Birkhäuser Verlag, Basel

0013-6018/90/030075-06\$1.50 + 0.20/0

## Kleine Mitteilung

### Zeros of characters and the Frattini subgroup

Let  $G$  be a finite group and let  $\text{Irr}(G)$  be the set of its (complex) irreducible characters. Of course, the Frattini subgroup  $\Phi(G)$  being normal in  $G$ , it must be an intersection of certain kernels of elements of  $\text{Irr}(G)$ . However, it seems that the problem of describing  $\Phi(G)$  in terms of characters is still open. The aim of this short Note ist to give a sufficient condition for the nontriviality of  $\Phi(G)$  in terms of vanishing sets of nonlinear irreducible characters of  $G$ .

Our notation is standard and follows that of [2]. Throughout,  $G$  will be a finite group and  $Z(G)$ ,  $G'$  will denote its centre and its derived subgroup, respectively. If  $\chi \in \text{Irr}(G)$  is nonlinear, the vanishing set  $A(\chi)$  of  $\chi$  is defined by  $A(\chi) := \{g \in G : \chi(g) = 0\}$ . A well-known result of Burnside asserts that  $A(\chi) \neq \Phi$ ; moreover, it's clear that  $A(\chi)$  is a union of conjugate classes of elements of  $G$ .

We prove the following

**Theorem.** *Let  $G$  be a finite group with  $1 < Z(G) < G$ . Suppose that there exists a nonlinear  $\chi \in \text{Irr}(G)$  such that  $A(\chi)$  contains fewer than  $|Z(G)|$  conjugate classes of elements. Then  $\Phi(G) \neq 1$ .*

**Proof:** The key observation is that actually  $A(\chi)$  is a union of cosets modulo  $Z(G)$ . To prove this, note that by Problem 3.12 of [2] it follows that for every  $g \in G$ ,

$$|\chi(g)|^2 = \frac{\chi(1)}{|G|} \sum_{n \in G} \chi([g, h]). \quad (*)$$

Let now  $g, h \in G$  and  $z \in Z(G)$ ; since  $[g, h] = [gz, h]$ , it follows from  $(*)$  that  $g \in A(\chi)$  iff  $gz \in A(\chi)$  for every  $z \in Z(G)$ . This means that  $A(\chi)$  is a union of cosets modulo  $Z(G)$ . Suppose, by way of contradiction, that  $\Phi(G) = 1$ . By a well-known result of [1],  $G' \cap Z(G) \leq \Phi(G)$ , so  $G' \cap Z(G) = 1$ .

Denote by  $s(\chi)$  and  $t(\chi)$  the number of conjugate classes of  $G$  contained in  $A(\chi)$  and the number of cosets modulo  $Z(G)$  lying in  $A(\chi)$ , respectively. We shall reach a contradiction by applying the pigeonhole principle. Suppose that  $g, h \in A(\chi)$ ,  $g \neq h$  and there exist  $u \in G$  and  $z \in Z(G)$  such that  $g = h^u = hz$ . Then  $z = h^{-1}g = h^{-1}h^u = [h, u] \in G' \cap Z(G)$ , whence  $g = h$ . This contradicts the choice of  $g$  and  $h$  and shows that  $A(\chi)$  contains at most  $s(\chi)t(\chi)$

elements. Taking now into account that  $t(\chi) := |A(\chi)|/|Z(G)|$ , it results that  $|Z(G)| \geq s(\chi)$ . But this contradicts the hypothesis and we are done.

**Corollary.** Let  $G$  be a finite group with  $1 < Z(G) < G$  and  $\Phi(G) = 1$ . Let

$$c := \max \{|C_G(x)| / x \in G \setminus Z(G)\} \quad \text{and} \quad b(G) := \max \{\chi(1) / \chi \in \text{Irr}(G)\}.$$

$$\text{Then } b(G) \geq \left( \frac{c}{c - |Z(G)|} \right)^{\frac{1}{2}}.$$

**Proof:** As a direct consequence of the Theorem, we obtain that  $c|A(\chi)| \geq |Z(G)||G|$  for every nonlinear  $\chi \in \text{Irr}(G)$ . On the other hand, it's a simple exercise to prove that  $|A(\chi)| < |G| \frac{\chi(1)^2 - 1}{\chi(1)^2}$ . Since  $\frac{(b(G))^2 - 1}{(b(G))^2} \geq \frac{\chi(1)^2 - 1}{\chi(1)^2}$  for every  $\chi \in \text{Irr}(G)$ , the result follows by combining these inequalities.

We have already seen that if  $\chi \in \text{Irr}(G)$  is nonlinear, then  $|Z(G)| \mid |A(\chi)|$ . This result may be refined in certain very special cases. For example, suppose that  $G$  is a finite group and  $\chi \in \text{Irr}(G)$  is faithful such that  $\chi(1) = 2$  and  $\chi(g) \in \mathbb{Q}$  for every  $g \in G$ . It is a matter of simple calculations to show that  $|A(\chi)| = 3|Z(G)|$ . If, moreover,  $Z(G) = 1$ , then  $|A(\chi)| = 3$  and  $G$  has a maximal subgroup of index 3 (the centralizer of an element of  $A(\chi)$ ).

Marian Deaconescu, Department of Mathematics,  
University of Timisoara, Romania

## REFERENCES

- 1 Gaschütz W.: Über die  $\Phi$ -Untergruppe endlicher Gruppen. Math. Z. 58, 160–170 (1953).
- 2 Isaacs I. M.: Character theory of finite groups. Academic Press, New York (1976).

## Aufgaben

**Aufgabe 1009.**  $n$  Zahlen  $x_1, \dots, x_n$  mit  $x_i \in \{0, 1, \dots, k\}$  ( $k \geq 2$ ) werden einmal linear, ein andermal kreisförmig so angeordnet, dass die Summe zweier Nachbarglieder stets von  $k + 1$  verschieden ist. Für beide Fälle bestimme man die Anzahl der zulässigen Anordnungen.

J. Binz, Bolligen

**Lösung des Aufgabenstellers (Bearbeitung der Redaktion).**

$s_n$  bzw.  $t_n$  bezeichnen die Anzahlen der zulässigen linearen bzw. kreisförmigen Anordnungen.

a) Es gilt  $s_n = a_n + b_n$ , wobei  $a_n$  die Anzahl der auf 0 endenden,  $b_n$  diejenige der übrigen zulässigen linearen Anordnungen bedeuten. Dann ist

$$a_{n+1} = a_n + b_n, \quad b_{n+1} = k a_n + (k - 1) b_n, \quad a_2 = k + 1, \quad b_2 = k^2,$$