Elemente der Mathematik		
Schweizerische Mathematische Gesellschaft		
46 (1991)		
6		
Dissections into equilateral triangles		
Tuza, Zsolt		
https://doi.org/10.5169/seals-43279		

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 09.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

ELEMENTE DER MATHEMATIK

Revue de mathématiques élémentaires - Rivista di matematica elementare

Zeitschrift zur Pflege der Mathematik und zur Förderung des mathematisch-physikalischen Unterrichts

El. Math.	Vol. 46	Nr. 6	Seiten 153-180	Basel, November 1991
-----------	---------	-------	----------------	----------------------

Dissections into equilateral triangles

Abstract. In a recent paper [El. Math., Vol. 46/4] H. Kaiser proved that every non-equilateral triangle has a dissection into finitely many (at most eight) similar but pairwise incongruent triangles. In this note we show that an equilateral triangle – and, more generally, any convex polygon – has no finite dissection into incongruent equilateral triangles. We also prove that the maximum number f(n) of distinct sizes of equilateral triangles in a dissection into n parts is equal to cn - o(n) for some constant $c \ge 5/7$.

1. Introduction

Let Π and Π' be polygons in the Euclidean plane. A dissection of Π into Π' is a decomposition of Π into finitely many, internally disjoint polygons Π'_1, \ldots, Π'_n $(n \ge 2)$ such that all of the Π'_i are similar to Π' . A dissection is perfect if the Π'_i are pairwise incongruent. Those Π'_i will be called the tiles of the dissection. The symbol Δ will be a shorthand for «equilateral triangle» (possibly with distinct side lengths at different places).

In the past fifty years it was extensively studied how squares and rectangles can be dissected into smaller squares. A detailed account on the history of this problem, with numerous references, can be found in the survey [2].

Relatively little is known, however, about dissections of polygons other than the square. Tutte [4, § 2] proved that a Δ has no perfect dissection into smaller Δ 's (this result was stated without proof in the classic paper [1]), and quite recently Kaiser [3] observed that in fact Δ is the only «exceptional case», i.e. every non-equilateral triangle has a perfect dissection (into at most eight tiles).

In this note, applying an argument much shorter than the original one in [4], we prove the following extension of Tutte's theorem.

Theorem 1. Every dissection of a convex polygon into equilateral triangles contains two triangles of the same size.

Knowing that every finite dissection of Δ contains at least two congruent tiles, it is natural to raise the following problem.

Problem 1. Given a positive integer $n \ge 6$, dermine the largest number f(n) of distinct side lengths in a dissection $\Delta_1 \cup \ldots \cup \Delta_n = \Delta$ of an equilateral triangle Δ .

Figure 1 shows that f(n) is well-defined for every $n \ge 6$, and also that $f(n) \ge 2$. (The dissection into 4 tiles is unique, with f(4) = 1, and no dissection exists with 5 tiles.) Here we prove

Theorem 2. There is a positive constant c, $5/7 \le c \le 1$, such that f(n) = cn - o(n) as $n \to \infty$.

It would be interesting to determine the exact value of $\lim_{n \to \infty} f(n)/n$. In particular, decide whether or not the limit is less than 1.



Figure 1. Dissection into an odd or even number of triangles.

We note that Δ has a «perfect dissection» into countably many tiles whose total area is equal to the area of Δ . The idea to show this fact is that a large part of a «long» trapezoid can be filled with finitely many Δ 's of nearly equal (but distinct) side lengths. It remains an open problem, however, to prove or disprove that no dissection into pairwise incongruent Δ 's is locally finite apart from the neighborhoods of a finite number of points.

2. Proofs

Proof of Theorem 1.

Consider a dissection $\Delta_1 \cup \ldots \cup \Delta_n = \Pi$ of a convex polygon Π into $n \ge 2$ internally disjoint Δ 's. We say that a Δ_i is split if a vertex of a Δ_j is an internal point of some side of Δ_i . If none of the Δ_i is split, then the (one, two, or three) neighbors of each Δ_i have the same size as Δ_i itself, and the connectedness of the dissection implies that all Δ_i are congruent. Hence, from now on we assume that there is at least one split triangle. Note that, by the assumption on convexity, each internal point splitting a side of some Δ_i is the vertex of precisely three tiles.

Suppose that $\Delta_1 = ABC$ is the smallest split triangle. We prove that an internal point of some side of Δ_1 is the vertex of two triangles Δ_i , $\Delta_{i'}$ of the same size.

A triangle Δ_j is said to overhang Δ_1 at A if the boundary of Δ_j entirely contains the side AB or AC of Δ_1 , and A is not a vertex of Δ_j . Certainly, at each vertex of a triangle at most one neighboring triangle can overhang, and if Δ_i overhangs Δ_j then $|\Delta_i| > |\Delta_j|$ (where $|\Delta_i|$ denotes any measure – side length, area, etc.).

We find two Δ_i of the same size, depending on the distribution of vertices on the periphery of Δ_1 . In the first three of the four possible cases we assume that each side of Δ_1 contains at most one internal vertex. In order to simplify some technical details of the argument, and to insure that each tile be surrounded completely by its neighbors, we artificially place an «external» Δ on each side of Π . Note that placing or removing external Δ 's does not change the status of split triangles in the dissection.

Case 1: The periphery of Δ_1 contains just one internal vertex.

Say, D is an internal vertex on AB. The two neighbors of Δ_1 containing C cannot be smaller than Δ_1 . Suppose that both of them are larger than Δ_1 . Since at most one of them can overhang at C, the other must overhang at, say, A (see Fig. 2(a)). Then AD is the side of some triangle $\Delta_j = ADE$. Since $|\Delta_j| < |\Delta_1|$, Δ_j is not split by the choice of Δ_1 . Moreover, at most one neighbor of Δ_j can overhang at E. Thus, AE or DE is the side of two triangles of the same size.



Case 2: The periphery of Δ_1 contains precisely two internal vertices, on distinct sides.

Let D and E be internal vertices in AC and BC, respectively. Since the edge AB is not split, some neighbor Δ_i of Δ_1 entirely contains AB, and if $|\Delta_i| \neq |\Delta_1|$ then Δ_i overhangs Δ_1 at, say, B (see Fig. 2(b)). Then the triangle BEF is in a similar situation as ADE in Case 1, so that it has a neighbor of the same size.

Case 3: The periphery of Δ_1 contains precisely three internal vertices, on pairwise distinct sides.

Let D_i (i = 1, 2, 3) be the internal vertices on the periphery of Δ_1 . Consider the triangles whose boundaries contain the segments CD_1 and CD_2 . Some of them does not contain C as an internal point of its side; say, $\Delta_j = CD_1E_1$ is a triangle of the dissection (see Fig. 3). By our assumptions, Δ_j is not split. Thus, if CD_1E_1 has no neighbor of the same size, then some triangle overhangs at each of its vertices. We denote by Δ_1 , the triangle that overhangs at C. The presence of Δ_1 implies that CD_2 is the side of some triangle CD_2F_1 , i.e. E_1F_1 is a side of Δ_1 . Repeating this argument for the triangles incident to Aand B, we obtain that the three vertices of Δ_1 are internal points on the peripheries of three triangles $\Delta_{1'}$, $\Delta_{2'}$, $\Delta_{3'}$ as exhibited in Fig. 3.



Observing the six neighbors of Δ_1 , we obtain $|CD_1| = |CE_1|$, $|CD_2| = |CF_1|$, etc., implying $|E_1F_1| + |E_2F_2| + |E_3F_3| = |AB| + |BC| + |CA|$ and $|\Delta_{1'}| + |\Delta_{2'}| + |\Delta_{3'}| = 3 |\Delta_1|$. On the other hand, since Δ_1 is a smallest split triangle, $|\Delta_{i'}| \ge |\Delta_1|$ holds for $1 \le i \le 3$. Consequently, $|\Delta_{i'}| = |\Delta_1|$ for $1 \le i \le 3$, i.e. the dissection contains four mutually congruent triangles.

Case 4: There are at least two internal vertices on the same side of Δ_1 .

Let D, E be two consecutive internal vertices in AB (see Fig. 2 (c)). Since the triangle DEF is not split, and at most one of its neighbors can overhang at F, its other neighbor has the same size. This fact completes the proof of Theorem 1. \Box

Proof of Theorem 2.

We prove the theorem in the slightly different – but equivalent – form that the limit

 $\lim_{n\to\infty} (f(n)-1)/(n-1)$

exists. Put g(n) = (f(n) - 1)/(n - 1). We are going to show that

 $c:=\liminf g\left(n\right)\geq g\left(k\right)$

for all $k \ge 6$. This inequality will imply the theorem.

Suppose that two dissections D_k and D_n are available, where D_i (i = k, n) has *i* tiles with f(i) distinct sizes. Substituting D_k into the smallest triangle of D_n , we obtain a dissection D_{n+k-1} with n + k - 1 tiles and f(n) + f(k) - 1 or f(n) + f(k) distinct sizes; the number of sizes after substitution is f(n) + f(k) if and only if the smallest size in D_n occurs at least twice. Applying this substitution t times, we obtain:

- (a) For every k, t, and n, $f(n + kt t) \ge f(n) + tf(k) t$, and
- (b) if there is a dissection of Δ into k tiles of f(k) distinct sizes such that the smallest size occurs at least twice, then $f(n + kt t) \ge f(n) + tf(k) 1$.

The reason is that the size of every triangle in the image of D_k is strictly smaller than the sizes in D_n .

Fixing the value of k $(k \ge 6)$, let n = (t + 1)(k - 1) + i, $0 \le i \le k - 2$, with t tending to infinity. Then (a) implies

$$g(n) = (f(n) - 1)/(n - 1)$$

$$\geq (f(i + k - 1) - 1 + t(f(k) - 1))/(i + k - 1 + t(k - 1))$$

$$= (f(k) - 1)/(k - 1) - o(1)$$

$$= g(k) - o(1).$$

This inequality holds independently of the value of *i*, so that $\liminf g(n) \ge g(k)$ follows, implying f(n) = c n + o(n) for some constant *c*.

The lower bound of $c \ge 5/7$ is obtained by repeatedly substituting the dissection D_{15} – taken from [4] – with 10 distinct sizes (the smallest one occurring twice), shown in Fig. 4. \Box



Figure 4. The dissection D_{15} .

As we have seen, property (a) is strong enough to prove the asymptotic result of Theorem 2. The advantage of part (b) is that it can be applied to obtain sharper lower bounds when a fairly good initial construction – like D_{15} – is available.

Acknowledgements. I am grateful to H. Kaiser for stimulating discussions related to Problem 1, and to the organizers of the conference «Geometry» for local hospitality at Rauischholzhausen (Germany), October 1990, where the bulk of this research was done.

Zsolt Tuza, Computer and Automation Institute, Budapest

REFERENCES

- 1 Brooks R. L., Smith C. A. B., Stone A. H., Tutte W. T.: The dissection of rectangles into squares. Duke Math. J. 7, 1940, 312-340.
- 2 Federico P. J.: Squaring rectangles and squares A historical review with annotated bibliography. In: Graph Theory and Related Topics (A. J. Bondy and U. S. R. Murty, Eds.), Academic Press, 1979, 173–196.
- 3 Kaiser H.: Perfekte Dreieckszerlegungen. El. Math. 46, No. 4, 106-111 (1991).
- 4 Tutte W. T.: The dissection of equilateral triangles into equilateral triangles. Proc. Cambridge Philos. Soc. 44 (4), 1948, 463-482.

© 1991 Birkhäuser Verlag, Basel

0013-6018/91/060153-06\$1.50 + 0.20/0

Klassische Beleuchtungsgeometrie im E^d $(d \ge 2)$ II. Kinematik in der Beleuchtungsgeometrie des E^d $(d \ge 2)$

In Teil I (vgl. [6]) wurden im \mathbb{R}^d ($d \ge 2$) Scharen aus isophotischen Flächenelementen bezüglich einer klassischen Zentralbeleuchtung (q, 1) – d.i. Lichtstärke 1 in jeder von der in q placierten Lichtquelle ausgehenden Richtung – konstruiert. Gemäss einer in [5] «kinematisch» genannten Vorgehensweise lassen sich nun aus der Zentralbeleuchtung (q, 1) und solch einer Schar isophotischer Elemente neue isophotische Scharen erzeugen. Diese Methode wird hier, nach kurzer Darlegung, verwendet, um Zusammenhänge zwischen Kurvenklassen aufzudecken, die die klassische Beleuchtungsgeometrie im E^d ($d \ge 2$) aufgrund ihres dimensionsabhängigen Beleuchtungsstärkegesetzes liefert.

1. Ein kinematisches Erzeugungsprinzip

Hat das orientierte Flächenelement (x, n) mit Trägerpunkt x und Einheitsnormalenvektor n die Beleuchtungsstärke E > 0 bezüglich der Zentralbeleuchtung (q, 1), so auch das Element (q, -n) bezüglich der Beleuchtung (x, 1). Die Ersetzung

 $((q, 1), (x, n)) \rightarrow ((x, 1), (q, -n))$