

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 47 (1992)

**Artikel:** A class of singular continuous functions  
**Autor:** Pei Hsu, Elton  
**DOI:** <https://doi.org/10.5169/seals-43922>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

---

# A Class of Singular Continuous Functions

---

Elton Pei Hsu

Northwestern University, Evanston, Illinois, USA

Pei Hsu obtained his Ph.D. in mathematics from Stanford University. Before joining the Department of Mathematics of Northwestern University, he has done research and taught at New York University and University of Illinois at Chicago. His main mathematical interests are probability theory and its applications to analysis.

A nondecreasing continuous function  $F(x)$  on the real line is *singular* if its derivative  $F'(x)$  is zero almost everywhere (a.e.). The most famous singular continuous function is Cantor's function discussed in every textbook on real analysis. In this short note we construct a class of singular continuous functions which includes Cantor's function as a

Auch wenn es dem Mathematik-Studenten manchmal so vorkommen mag, die grundlegenden mathematischen Begriffe, wie "Funktion", "Konvergenz", "Stetigkeit", etc. sind nicht plötzlich in ihrer heutigen Eindeutigkeit und Schärfe vom Himmel gefallen. Vielmehr haben sie sich aus intuitiven Anfängen im Laufe langer Jahre langsam entwickelt: nur schrittweise haben sie an Schärfe gewonnen, erst nach und nach wurde ihr Verhältnis zur bestehenden Begriffswelt geklärt. Intensive Bemühungen führten schliesslich zu den heute gängigen ausgefeilten Definitionen dieser Konzepte. Ein Beispiel, an dem sich diese Entwicklung gut verfolgen lässt, ist der Begriff der Stetigkeit einer Funktion. Die Beziehungen der Eigenschaft der Stetigkeit zu anderen, wie etwa zur Differenzierbarkeit, und zur Geometrie waren während langer Zeit unklar. Immer wieder wurden Beispiele entdeckt, deren Eigenschaften die Fachwelt überraschten und die zur Korrektur von damals üblichen, aber eben irrgänigen Annahmen zwangen. — Pei Hsu geht in seinen Beitrag auf eine spezielle derartige Frage ein. Ausgehend von der bekannten Cantor-Funktion konstruiert er eine Klasse von monoton zunehmenden stetigen Funktionen, welche fast überall eine triviale Ableitung besitzen. Solche Funktionen, man nennt sie *singuläre stetige Funktionen*, widersprechen natürlich einer naiven Anschauung der Stetigkeit. — Eine sehr gut zugängliche Darstellung des ganzen in diesem Beitrag behandelten Themenkreises ist im Buch von Ralph P. Boas, jr. zu finden *A primer of real functions*, Carus Mathematical Monographs 13, Mathematical Association of America (1972). Für die Cantor-Funktion vergleiche man S. 135. *ust*

special case. Unlike Cantor's function, most functions in this class are strictly increasing on the interval  $[0, 1]$ .

We construct these singular functions as the distribution functions of certain random variables. We also give an explicit, but less instructive, formula for these functions. Readers not familiar with probability theory may be more comfortable with the explicit formula.

Let  $N \geq 2$  be a positive integer and  $\mathbf{p} = [p_0, \dots, p_{N-1}]$  be an  $N$ -dimensional probability vector, namely,

$$p_i \geq 0, \quad i = 0, \dots, N-1; \quad \sum_{i=0}^{N-1} p_i = 1.$$

We exclude the trivial case where one of the  $p_i$ 's is equal to 1. Next let  $X_n, n = 1, 2, \dots$  be a sequence of independent, identically distributed random variables with the common distribution

$$P[X_1 = i] = p_i, \quad i = 0, \dots, N-1.$$

Define a new random variable  $X$  with values in the interval  $[0, 1]$ :

$$X = \sum_{n=1}^{\infty} \frac{X_n}{N^n}.$$

Thus  $X$  is a number in  $[0, 1]$  whose digits in base  $N$  expansion  $X = 0.X_1X_2\dots$  are chosen randomly and independently according to the probability vector  $\mathbf{p}$ . Let  $F(x; \mathbf{p})$  be the distribution function of  $X$ ,

$$F(x; \mathbf{p}) = P[X \leq x].$$

We now derive an explicit formula for  $F(x; \mathbf{p})$ . In the following we simply write  $F(x)$  instead of  $F(x; \mathbf{p})$  if no confusion is possible.

It is clear that  $F(x) = 0$  for  $x \leq 0$  and  $F(x) = 1$  for  $x \geq 1$ . For  $0 < x < 1$ , suppose that  $x = 0.x_1x_2\dots$  is the expansion of  $x$  in base  $N$ . For definitiveness, when two expansions are possible, for example  $0.13\bar{0} = 0.12\bar{N-1}$  ( $\bar{k}$  means repeating the digit  $k$  indefinitely), unless indicated otherwise we always take the nonterminating one, namely, the one with only the digit  $N-1$  from some digit on. Let  $x^n = 0.x_1x_2\dots x_n$  be the truncation of  $x$  at the  $n$ th place. Then since  $P[X = x] = P[X \leq 0] = 0$ , we have

$$F(x) = \sum_{n=0}^{\infty} P[x^n < X \leq x^{n+1}].$$

Now  $x^n < X \leq x^{n+1}$  means that  $X$  must have the form  $X = 0.x_1x_2\dots x_nX_{n+1}X_{n+2}\dots$ , where  $X_{n+1}$  is equal to one of the digits  $0, 1, \dots, x_{n+1}-1$ . Note that we may assume that the expansion  $X = 0.X_1X_2\dots$  is always nonterminating, for the probability that  $X$  has a terminating expansion is zero. Thus

$$P[x^n < X \leq x^{n+1}] = p_{x_1} \cdots p_{x_n} (p_0 + \cdots + p_{x_{n+1}-1}).$$

In the above inequality, the sum between the parentheses should be set equal to zero if  $x_{n+1} = 0$ . We now obtain

$$F(x; \mathbf{p}) = \sum_{n=0}^{\infty} p_{x_1} \cdots p_{x_n} (p_0 + \cdots + p_{x_{n+1}-1}).$$

This is an explicit formula for the function  $F(x; \mathbf{p})$ . Incidentally, the above explicit representation works when  $x$  is expressed in its terminating expansion as well. The special case  $N = 3, \mathbf{p} = [\frac{1}{2}, 0, \frac{1}{2}]$  gives Cantor's function.

We will prove two properties of these functions.

- (I)  *$F(x; \mathbf{p})$  is strictly increasing unless one of the  $p_i$ 's is equal to zero, and is Hölder continuous with the exponent*

$$\alpha = \frac{\log \frac{1}{r}}{\log N},$$

where  $r = \max \{p_0, \dots, p_{N-1}\}$ .

- (II)  *$F(x; \mathbf{p})$  is singular continuous except when  $\mathbf{p}$  is the uniform distribution:*

$$\mathbf{p} = \left[ \frac{1}{N}, \dots, \frac{1}{N} \right],$$

in which case  $F(x; \mathbf{p}) = x$  on  $[0, 1]$ .

**Proof of (I).** If  $x, y$  in  $[0, 1]$  have the same first  $n$  digits, then from the explicit formula for  $F(x)$  we have

$$|F(x) - F(y)| \leq \sum_{l=n}^{\infty} r^l \leq \frac{r^n}{1-r}.$$

Let  $x, y$  be two distinct numbers in  $[0, 1]$ . Pick a nonnegative  $n$  such that  $N^{-(n+2)} \leq |x - y| < N^{-(n+1)}$ . Then in the nonterminating expansions of  $x$  and  $y$  either they have the same first  $n$  digits or  $x^n = y^n + N^{-n}$ . In the latter case  $y^n = 0.x_1x_2 \cdots x_n \overline{N-1}$  and  $x^n$  have the same first  $n$  digits. Therefore we have

$$\begin{aligned} |F(x) - F(y)| &\leq |F(x) - F(x^n)| + |F(x^n) - F(y^n)| + |F(y^n) - F(y)| \\ &\leq \frac{3r^n}{1-r}. \end{aligned}$$

It follows that for any  $x, y$  in  $[0, 1]$  with  $|x - y| \leq N^{-1}$  we have

$$|F(x) - F(y)| \leq \frac{3}{r^2(1-r)} |x - y|^{\alpha}.$$

This shows that  $F(x)$  is Hölder continuous with exponent  $\alpha$ .

Let  $x = 0.x_1x_2\dots$  (in base  $N$ ) be a point in  $[0, 1]$ . Let  $x^n = 0.x_1x_2\dots x_n$  as before. Then

$$x^n < x \leq x^n + \frac{1}{N^n}.$$

Writing  $x^n + N^{-n}$  as  $0.x_1\dots x_n\overline{N-1}$ , from the explicit formula for  $F(x)$  we have

$$F(x^n + N^{-n}) - F(x^n) = p_{x_1}\dots p_{x_n}.$$

This shows that  $F$  is strictly increasing on  $(0, 1)$  if none of the  $p_i$ 's is equal to zero.

**Proof of (II).** From the above equality we see that

$$D_n F(x) \stackrel{\text{def}}{=} N^n [F(x^n + N^{-n}) - F(x^n)] = N p_{x_1}\dots N p_{x_n}.$$

Since  $F(x)$ , being a monotone function, is a.e. differentiable, we have a.e.,

$$\lim_{n \rightarrow \infty} D_n F(x) = F'(x) \text{ and } F'(x) \text{ is finite.}$$

It follows that

$$F'(x) = \prod_{n=1}^{\infty} N p_{x_n}, \quad \text{a.e.}$$

If all of  $p_i$ 's are equal to  $N^{-1}$ , then  $F'(x) = 1$  a.e. on  $[0, 1]$ . We have  $F(x) = x$  on  $[0, 1]$ .

If at least one of the  $p_i$ 's is not equal to  $N^{-1}$ , say,  $p_1 \neq N^{-1}$ , then for those  $x$  whose expansion has infinitely many 1's the general term  $N p_{x_n}$  of the above infinite product does not converge to 1. Therefore, by an elementary theorem on infinite products, the value of the infinite product, if it exists and is finite, must be 0. On the other hand, the reader can verify easily that the numbers whose expansions have only finitely many 1's form a set of Lebesgue measure zero. Therefore we have proved  $F'(x) = 0$  a.e. and  $F(x)$  is a singular continuous function.

For readers familiar with the strong law of large numbers in probability theory the following argument may be more appealing. By the strong law of large numbers, almost all numbers in  $[0, 1]$  are normal in base  $N$ , namely, if  $s_n(x; i)$  is the number of the digit  $i$  among the first  $n$  digits of the expansion of  $x$ , then for almost all  $x \in [0, 1]$ ,

$$\text{for all } i = 0, \dots, N-1 : \quad \frac{s_n(x; i)}{n} \rightarrow \frac{1}{N} \quad \text{as } n \rightarrow \infty.$$

It follows that, if at least one of the  $p_i$ 's is not equal to  $N^{-1}$ , then for almost all  $x \in [0, 1]$

$$\sqrt[n]{D_n F(x)} = \prod_{i=0}^{N-1} (N p_i)^{s_n(x; i)/n} \rightarrow N \sqrt[N]{p_0 \cdots p_{N-1}} < 1.$$

This implies immediately  $F'(x) = 0$ , a.e.

Elton Pei Hsu  
 Department of Mathematics  
 Northwestern University  
 Evanston, IL 60208, USA