

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 49 (1994)

**Artikel:** Why means in two arguments are special  
**Autor:** Lorenzen, Gunter  
**DOI:** <https://doi.org/10.5169/seals-45418>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 17.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

---



---

## Why Means in Two Arguments are Special

---



---

Gunter Lorenzen

Gunter Lorenzen wurde 1942 geboren. Er studierte Volkswirtschaftslehre an der Universität Kiel und anschliessend Mathematik an der Universität Hamburg, 1974 habilitierte er sich dort für Volkswirtschaftslehre. Lehrstuhlvertretungen in Kiel und Frankfurt und zwei einjährige DAAD-Gastprofessuren an der University of East Asia in Macao folgten. Gegenwärtig ist Lorenzen Professor für Statistik und Ökonometrie an der Universität Hamburg. Sein wissenschaftliches Interesse gilt der quantitativen Wirtschaftsforschung und Statistik.

### 1 Introduction

Starting from the Cauchy-version of the mean value theorem: there exists  $z$  between  $x$  and  $y$  such that

$$\frac{f(x) - f(y)}{g(x) - g(y)} = \frac{f'(z)}{g'(z)}, \quad (1)$$

choosing  $f(x) = x^r, g(x) = x^s$ , and solving for  $z$ , Stolarsky (1975) established the following mean value function:

$$N_{r,s}(x, y) = \left[ \frac{s(x^r - y^r)}{r(x^s - y^s)} \right]^{\frac{1}{r-s}} \quad (r, s \neq 0, r \neq s). \quad (2)$$

$N_{r,s}(x, y)$  has been extensively discussed in Stolarsky (1975, 1980), as well as Leach and Scholander (1978, 1983). The special case of  $s = 1$  was introduced in Galvani (1927), already, and Cisbani (1938) investigated the case  $s = 0$  (which has to be defined by continuity extension).  $N_{r,s}(x, y)$  includes the traditional power mean  $M_s(x, y) = \left( \frac{x^s + y^s}{2} \right)^{1/s}$  at  $r = 2s$ .

We commence by pointing out that generalizations of the power mean  $M_s$  to more than two arguments preserve a specific mean value property, whereas generalizations of the more extensive Stolarsky mean  $N_{r,s}$  do not.

Die Bildung des harmonischen, geometrischen und arithmetischen Mittelwertes von zwei Zahlen kann auf eine *kontinuierliche* Familie von Mittelwertbildungen verallgemeinert werden. Diese Sachlage hängt offenbar in entscheidender Weise damit zusammen, dass genau zwei Argumente betrachtet werden. Gunter Lorenzen geht in seinem Beitrag zuerst der Frage nach, weshalb hier die Anzahl zwei eine Sonderrolle spielt, und gibt anschliessend einige Anwendungen der verallgemeinerten Mittelwertbildungen. *usf*

## 2 Generalized Stolarsky means

In order to generalize  $N_{r,s}$  to be able to handle more than two arguments, we should start from a generalized mean value theorem. Exploitation of the concept of generalized divided differences (Nörlund (1924), Milne-Thomson (1933)) leads to the mean value theorem: there exists  $z$  such that

$$\sum_{i=1}^n \frac{f(x_i)}{\prod_{j \neq i} (g(x_i) - g(x_j))} = \frac{\Delta^{n-1}(z)}{(n-1)!} \tag{3}$$

with  $\min(x_1, \dots, x_n) \leq z \leq \max(x_1, \dots, x_n)$  and where

$$\Delta^{(1)} = \frac{f'}{g'}, \dots, \Delta^{(n-1)} = \frac{(\Delta^{(n-2)})'}{g'}$$

Choosing  $f(x) = x^r$  and  $g(x) = x^s$  and solving for  $z$  again, results in the generalized Stolarsky mean:

$$N_{r,s}(x_1, \dots, x_n) = \left[ \frac{(n-1)! \ s^{n-1}}{r(r-s)(r-2s) \dots (r-(n-2)s)} \sum_{i=1}^n \frac{x_i^r}{\prod_{j \neq i} (x_i^s - x_j^s)} \right]^{\frac{1}{r-(n-1)s}} \tag{4}$$

$(r, s \neq 0; r/s \neq 1, \dots, (n-1)).$

Clearly, at  $n = 2$ , (4) and (2) are identical. Furthermore,  $N_{r,s}(x_1, \dots, x_n)$  includes the power mean  $M_s = (\frac{1}{n} \sum x_i^s)^{1/s}$  at  $r = ns$ . This result follows from

$$\sum \frac{x_i^{ns}}{\prod (x_i^s - x_j^s)} = \sum x_i^s.$$

So the question is, why has the power mean  $M_s$  occupied so much of the attention of the profession, whereas the more general  $N_{r,s}$  and similar expressions (see Leach and Scholander (1984)) seem to have gone more or less unnoticed.

## 3 On a specific mean value property postulated by Kolmogoroff and Nagumo

The reason for putting emphasis on power means seems to be hidden in two papers that were published more than sixty years ago. Kolmogoroff (1930) and Nagumo (1930) simultaneously and independently worked out the axioms required to single out the power mean  $M_s$ . Apart from ‘natural’ properties such as symmetry, homogeneity, boundedness ( $\min(x_i) \leq M \leq \max(x_i)$ ) and monotonicity, they demanded that a mean  $M$  should comply with the following characteristic:

$$M_n(x_1, \dots, x_n) = M_n(M_r(x_1, \dots, x_r) \circ r, x_{r+1}, \dots, x_n) \tag{5}$$

where  $M_r(x_1, \dots, x_r) \circ r$  stands for “ $r$  times the argument  $M_r(x_1, \dots, x_r)$ ”. Kolmogoroff and Nagumo thus demonstrated that it is property (5) by which the power mean  $M_s$  is

distinguished from other means. We have to conclude, therefore, that  $N_{r,s}(x_1, \dots, x_n)$  from (4) is a deficient mean value function in the sense that it lacks property (5) (except at  $r = ns$ ).

The most serious consequence is this: in (3) and consequently in (4) we have (implicitly) assumed the  $x_i$ 's to be different. With some  $x_i$ 's being equal instead, the appropriate number of divided differences in (3) has to be replaced by derivatives. But the result of (4), then, depends on the sequence of differentiation. Specifically,  $N_{r,s}(x, x, y, y)$  say, is different from  $N_{r,s}(x, y)$  and is not even uniquely determined, because  $N_{r,s}(x_1, \dots, x_n)$  is not continuous at  $x_i = x_j$ , except at  $r = ns$ .

To put this consequence in slightly different terms: it is not possible to have a meaningful definition of 'weighted' means if property (5) is being neglected.

But in the case of two arguments only, property (5) is meaningless. In fact, the technique of proof in Kolmogoroff (1930) and Nagumo (1930) breaks down at  $n = 2$ . Thus it is indeed unnecessarily restrictive in the case of two arguments to confine attention to power means. Specifically,  $N_{r,s}(x, y)$  from (2) is deficient in no respect, as it fulfills all the Kolmogoroff-Nagumo axioms. However, generalizing these Stolarsky means (or other means) to be able to handle more than two arguments can be done only, if one is willing to forgo property (5) and suffer the consequences.

#### 4 A mean value function in two arguments that ought to be more popular

At  $r = s + 1$  the mean  $N_{r,s}(x, y)$  reduces to

$$P_s(x, y) = \frac{s(x^{s+1} - y^{s+1})}{(s+1)(x^s - y^s)} \quad s \neq -1, 0$$

$$P_{-1}(x, y) = xy \frac{\ln x - \ln y}{x - y} \quad (s = -1) \quad (6)$$

$$P_0(x, y) = \frac{x - y}{\ln x - \ln y} \quad (s = 0)$$

$P_0(x, y)$  has won some popularity in recent years under the name 'logarithmic mean' (Carlson 1972). We concentrate on this logarithmic mean in section 5 and shall write  $P_0(x, y) = L(x, y)$  here-after.

The following scheme concentrates on the 5 most relevant elements of  $P_s(x, y)$  where  $H$ ,  $G$ ,  $L$  and  $A$  denote the harmonic, geometric, logarithmic, and arithmetic mean, respectively.

$s$	$P_s(x, y)$	$P_s(x, x^{-1})$
-2	$H = G^2/A$	$1/A$
-1	$G^2/L$	$1/L$
$-\frac{1}{2}$	$G$	$1$
0	$L$	$L$
1	$A$	$A$

We conclude that  $P_s(x, y)$  resembles the traditional power mean  $M_s(x, y)$  in bridging the gap from  $H$  to  $G$  to  $A$  and that  $P_s(x, y)$  has the additional advantage of including as well the logarithmic mean  $L$  (and the still unbaptized  $G^2/L$ ).

Stolarsky (1975) proves that  $N_{r,s}(x, y)$  increases monotonically in both the parameters  $r$  and  $s$ . It follows that  $P_s(x, y)$  increases monotonically in  $s$  which leads to the inequality

$$H \leq G^2/L \leq G \leq L \leq A$$

through which we are informed that the logarithmic mean is bounded numerically by the geometric mean and the arithmetic mean.

To prove our point that  $P_s(x, y)$  is a more interesting mean value function than the traditional power mean  $M_s(x, y)$  (as long as we concentrate on two arguments only), we have to demonstrate that the logarithmic mean is useful in applications.

### 5 Applications of the logarithmic mean

Burkhardt (1933) seems to have been the first to give a direct application of the logarithmic mean. Assume the population size  $B(t)$  of birth-cohort to change exponentially:

$$B(t) = ae^{\gamma t}$$

and suppose that the total amount of person-years within a given period  $[t, t + 1]$  (one of the basic concepts of life-table construction) is the problem under investigation.

Then we have

$$\int_t^{t+1} B(x)dx = L(B(t + 1), B(t))$$

because of  $\gamma = \ln B(t + 1) - \ln B(t)$ . To calculate the amount of person-years it is sufficient, therefore, to have census results at the beginning and at the end of the period with the logarithmic mean doing the trick.

Another application of  $L(x, y)$  is to be found in Leach and Scholander (1978): It is well known that the following inequality holds:

$$\left(1 + \frac{1}{n}\right)^n \leq e \leq \left(1 + \frac{1}{n}\right)^{n+1}.$$

So the question arises, if there is a mean  $M(n, n + 1)$  for which

$$\left(1 + \frac{1}{n}\right)^{M(n, n+1)} = e$$

is true. The answer is in the affirmative: choose  $M(n, n + 1) = L(n, n + 1)$ . The resulting identity can easily be extended to

$$\left(1 + \frac{x}{n}\right)^{L(n, n+x)} = e^x.$$

Another identity involving the logarithmic mean, which has some relevance to claim (see Lorenzen 1989), is

$$\tanh^{-1}(x) = \frac{x}{L(1+x, 1-x)}$$

which contrasts with  $x = x/A(1+x, 1-x)$  so that

$$f_s(x) = \frac{x}{P_s(1+x, 1-x)}$$

includes  $f_1(x) = x$  as well as  $f_0(x) = \tanh^{-1}(x)$ .

More applications of the logarithmic mean (and other non-standard means) are to be found in Vartia (1976) and Lorenzen (1990, 1992).

## 6 Conclusion

Whereas it is usually well justified to concentrate on the traditional family of power means, as only these means comply with the Kolmogoroff-Nagumo-property (5), it is unnecessarily restrictive to do so in the case of means in two arguments. Non-standard means in two arguments may have all the required properties and have some relevance to claim within applications.

## References

- [1] Burkhardt, F. (1933): Die Bewegung der Sterblichkeit im Spiegel der Statistik, *Assekuranz Jahrbuch*, 52, 192–208.
- [2] Cisbani, R. (1938): Contributi alla teoria delle medie, *Metron*, 13, 23–60.
- [3] Galvani, L. (1927): Dei limiti a cui tendono alcune medie, *Bollettino della Unione Matematica Italiana*, 6, 173–179.
- [4] Carlson, B.C. (1972): The logarithmic mean, *American Mathematical Monthly*, 79, 615–618.
- [5] Kolmogoroff, A. (1930): Sur la notion de la moyenne, *Atti della Reale Accademia Nazionale dei Lincei (Roma)*, 12, 388–391.
- [6] Leach, E.B. and M.C. Scholander (1978): Extended Mean Values, *American Mathematical Monthly*, 85, 84–90 and 656.
- [7] Leach, E.B. and M.C. Scholander (1983): Extended Mean Values II, *Journal of Mathematical Analysis and Applications*, 92, 207–223.
- [8] Leach, E.B. and M.C. Scholander (1984): Multivariable Extended Mean Values, *Journal of Mathematical Analysis and Applications*, 104, 390–407.
- [9] Lorenzen, G. (1989): Log-ratios and the Logarithmic mean, *Statistical Papers (Statistische Hefte)*, 30, 61–75.
- [10] Lorenzen, G. (1990): Konsistent addierbare relative Änderungen, *Allgemeines Statistisches Archiv*, 74, 336–344.
- [11] Lorenzen, G. (1992): A Reformulation of Pearson's Chi-square Statistic and some Extensions, *Statistics and Probability Letters*, 14, 327–331.

- 
- [12] Milne-Thomson, L.M. (1933): *The Calculus of Finite Differences*, London.
- [13] Nagumo, M. (1930): Über eine Klasse von Mittelwerten, *Japanese Journal of Mathematics*, 71–79.
- [14] Nörlund, N.E. (1924): *Vorlesungen über Differenzenrechnung*, Berlin.
- [15] Stolarsky, K.B. (1975): Generalizations of the Logarithmic Mean, *Mathematics Magazine*, 48, 87–92.
- [16] Stolarsky, K.B. (1980): The Power and Generalized Logarithmic Means, *American Mathematical Monthly*, 87, 545–548.
- [17] Vartia, Y.O. (1976): Ideal Log-Change Index Numbers, *Scandinavian Journal of Statistics*, 3, 121–126.

Gunter Lorenzen  
Institut für Statistik und Ökonometrie  
Universität Hamburg  
Von-Melle-Park 5  
20146 Hamburg  
Deutschland