**Zeitschrift:** Elemente der Mathematik

Herausgeber: Schweizerische Mathematische Gesellschaft

**Band:** 49 (1994)

**Artikel:** The magic world of geometry. I, The isoperimetric problem

Autor: Lunsgaard Hansen, Vagn

**DOI:** https://doi.org/10.5169/seals-45420

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 01.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch


# The Magic World of Geometry — I. The Isoperimetric Problem

## Vagn Lundsgaard Hansen

Vagn Lundsgaard Hansen received his M.Sc. in mathematics and physics from the University of Aarhus, Denmark in 1966 and his Ph.D. in mathematics from the University of Warwick, England in 1972. Since 1980 he is professor of mathematics at The Technical University of Denmark. He has published research papers in topology, geometry and global analysis, and the books *Braids and Coverings* (1989) and *Geometry in Nature* (1993). Also, he was the editor of the *Collected Mathematical Papers of Jakob Nielsen* (1986). He enjoys philosophical discussions, music and family life.

Nature is uncompromisingly effective, and from a geometrical point of view its shapes are full of mathematics. The study of optimal properties of geometric objects is therefore both interesting and important. In this article, I shall present the isoperimetric problem, which is an archetypical problem in this context. The article is the first in a series of three articles with the common subtitle "The Magic World of Geometry" intended to show how geometry appeals to the imagination and how it combines the concrete with the abstract.

Unter dem Sammeltitel The Magic World of Geometry hat V.L. Hansen drei Beiträge verfasst, deren erster hier vorliegt. Der Autor behandelt darin das isoperimetrische Problem in der Ebene: Finde diejenige geschlossene Kurve in der Ebene, welche bei gegebener Länge den grössten Flächeninhalt einschliesst! Er beschreibt den geometrischen Lösungsansatz des grossen Schweizer Mathematikers J. Steiner (1796–1863). Dieser Beweis hat den Vorteil grosser Anschaulichkeit, und er ist damit auch beispielhaft für Steiners Mathematik. Einige Jahre nach Steiners Veröffentlichung machte Weierstrass allerdings darauf aufmerksam, dass Steiner stillschweigend die Existenz einer Lösung vorausgesetzt hatte. Anschaulich scheint die Existenz einer Lösung zwar offensichtlich, aber Weierstrass konnte anhand ähnlicher Fragestellungen zeigen, dass sie nicht ohne weiteres angenommen werden darf. Die Beweislücke wurde später durch Weierstrass und Blaschke geschlossen. ust



The isoperimetric problem is one of the classical geometric variational problems. In its simplest form the problem can be stated as follows: Find the closed plane curve (without self-intersections) of a fixed prescribed length that encloses the maximal plane area.

The ancient Greeks took it for granted that the solution to the problem is what they regarded as the most perfect of all closed curves, namely the circle. However, not until 19th century was a complete proof given. Jacob Steiner (1796–1863) suggested several ingenious proofs, and we shall present one below.

Before we embark on Steiner's proof, it will be appropriate to make some general remarks concerning variational problems. Before Weierstrass (1815–1897) it was accepted without proof that an extremal problem, of a physical or geometrical nature, always has a solution. It is one of the many merits of Weierstrass to have pointed out that this is by no means the case. In a primitive, numerical way we can see this already from the real numbers. For example, the set of fractions

$$\left\{\frac{1}{2},\frac{1}{3},\frac{1}{4},\ldots,\frac{1}{n},\ldots\right\}$$

does not contain a smallest positive number.

Weierstrass gave several simple examples of geometrical problems without a "minimal" solution. In Figure 1 and 2 we give two such examples. In Figure 1 we consider two points A and B on a line, and seek the shortest polygonal path in the plane, which starts in A orthogonally to the line and ends in B. Obviously, there is no such shortest path. In Figure 2 we consider two points on a line through the origin in the plane. If we remove the origin from the plane, there is no longer a shortest path connecting the two points.

It is instructive to learn that during the history of mathematics, even great mathematicians like Dirichlet (1805–1859) and Riemann (1826–1866) have overlooked that the fundamental problem can be hidden where it is least expected. In the calculus of variations the main problem often is the bare question of the existence of an optimal object.

In famous lectures at the University of Berlin in the 1870's, Weierstrass developed general methods to ensure the existence of global maxima and minima of continuous functionals. Using these methods it is not difficult to prove that the isoperimetric problem has a solution if we restrict our attention to n-gons, i.e. closed plane polygonal curves without self-intersections and with n edges. First note that in a coordinate system in the plane, an n-gon can be described by the 2n coordinates of its corners. Observe next that all possible shapes of an n-gon with a fixed prescribed length can be described by the coordinates in a closed and bounded subset of 2n-dimensional real number space. Since the area of an n-gon depends continuously on the 2n coordinates of its corners, the area attains a maximum value on this closed and bounded set according to a fundamental theorem




Fig. 3

of Weierstrass. This proves the existence of a solution to the restricted isoperimetric problem. The existence of a solution to the general isoperimetric problem can now be proved by approximating an arbitrary closed (rectifiable) curve with polygonal curves. More details on the existence of a solution to the isoperimetric problem can be found in references ([1], §5 and §8), ([2], Chp. IV, §2.1, p. 149) and ([3], Chp. VII, §8, p. 375).


It is easy to illustrate by the following experiment that the circle is the solution to the isoperimetric problem. Suspend a closed curve in a small frame. The curve and the threads with which it is attached to the frame can e.g. be sewing thread. Now dip the frame in a suitable soap solution. Thereby the closed curve will be placed on a soap film. Prick a hole in the soap film inside the closed curve. Immediately the closed curve turns into a circle; cf. Figure 3. It follows that the circle must be the solution to the isoperimetric problem, since the potential energy in the soap film outside the curve is proportional to the area, so that the soap film will try to minimize the area, which on the other hand corresponds to the area inside the closed curve being maximized.

The physical experiment is pretty convincing but not sufficient evidence from a mathematical point of view. We could be misled by our senses. Maybe it is only very close to being a circle. Therefore we consider now a closed plane curve  $\mathscr C$  without self-intersections and of a fixed prescribed length L with the property that, among all such curves, it encloses the largest plane area. Following Steiner, we shall then argue mathematically that  $\mathscr C$  must be a circle. It should be mentioned that Steiner also took it for granted that the isoperimetric problem has a solution.

By considering Figure 4 it is clear that  $\mathscr{C}$  must be convex, since if this was not the case we could create a closed curve of the same length L as  $\mathscr{C}$ , but enclosing a larger area, by reflecting an inbuckling to an outbuckling.

Now choose two points A and B on  $\mathcal{C}$  that divide the curve into two arcs of equal length, i.e. length L/2. The chord AB divides the enclosed figure into two pieces, which each must have the same area since, otherwise, we could construct a figure of larger area by replacing the smaller piece with the mirror image in AB of the larger piece; cf. Figure 5.

Consider now the "half" figure bounded by one of the arcs from A to B of length L/2 and the line segment AB. Let P be an arbitrary point on the arc, and consider the triangle APB. Imagine that the arc is made of steel with a hinge at P. Without changing the length of the arc we can then bend the triangle at the corner P. Thereby it is easily seen, that the "half" figure has maximal area precisely when the triangle has a right angle at



P; cf. Figure 6. Since P was arbitrarily chosen, we conclude that the arc from A to B must be a semicircle, and the original closed curve  $\mathscr C$  therefore a circle.

## Literature:

- [1] W. Blaschke: Kreis und Kugel. Verlag von Veit & Comp. 1916.
- [2] R. Courant und D. Hilbert: Methoden der Mathematischen Physik I. 3. Auflage, Springer-Verlag, 1968.
- [3] R. Courant and H. Robbins: What is Mathematics? Oxford University Press, 1941.
- [4] H.G. Eggleston: The Isoperimetric Problem. Chapter 7, Exploring University Mathematics 1, N.J. Hardiman, Pergamon Press, 1967.
- [5] V.L. Hansen: Geometry in Nature. A K Peters, Ltd., Wellesley, Massachusetts, US, 1993.
- [6] S. Hildebrandt and A. Tromba: Mathematics and Optimal Form. Scientific Amer. Library, W.H. Freeman and Co, 1985.
- [7] M. Kline: Mathematical Thought from Ancient to Modern Times. Oxford University Press, 1972.

Vagn Lundsgaard Hansen Mathematical Institute Building 303 Technical University of Denmark DK-2800 Lyngby Denmark