

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 51 (1996)

Artikel: Three-Dimensional Non-Commutative Algebras
Autor: Heimbeck, Günter / Kamupingene, Alfred H.
DOI: <https://doi.org/10.5169/seals-46956>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Three-Dimensional Non-Commutative Algebras

Günter Heimbeck and Alfred H. Kamupingene

Born in 1946, Günter Heimbeck studied at Würzburg University where he attained his Dr.rer.nat.habil. in 1981. After having lectured for two and a half years at Würzburg University he joined the University of the Witwatersrand, Johannesburg in 1984. Since 1987 he is a member of the Department of Mathematics of the University of Namibia in Windhoek. His main interest pertains to geometric algebra and translation planes.

Alfred H. Kamupingene was born in 1956 in Namibia. He studied at Sofia University specializing in mathematical physics. After completing his Ph.D. in 1988 on representations of Lie superalgebras and their applications to Wigner quantum systems he spent two years at the Bulgarian Academy of Sciences as a post-doctoral fellow. Currently he is an Associate Professor at the University of Namibia. His main interest is in algebra, topology and mathematical physics.

As it is well-known, for any prime p , there exists, up to isomorphism, exactly one non-commutative ring of order p^3 ([1], p. 195, Further exercises (6)). The purpose of this short note is to generalize this result to algebras.

Theorem. *For any field F , there exists, up to isomorphism, exactly one 3-dimensional non-commutative F -algebra, namely*

$$\left\{ \left(\begin{array}{cc} \alpha & \beta \\ 0 & \gamma \end{array} \right) \mid \alpha, \beta, \gamma \in F \right\}.$$

In jeder axiomatischen Theorie bemüht sich die Mathematik, über die verschiedenen (nichtisomorphen) "kleinen" Modelle einen Überblick zu gewinnen. So versucht man etwa in der Gruppentheorie, alle Gruppen von kleiner Ordnung aufzuzählen, oder man versucht, für eine Primzahl p , die p -Gruppen der Ordnung p, p^2, p^3 , etc. der Reihe nach zu beschreiben. Ein derartiges Vorgehen liefert in jeder axiomatischen Theorie eine Reihe von interessanten Beispielen. Auch wenn ein tieferes Verständnis gewöhnlich erst auf Grund weiterführender theoretischer Methoden möglich ist, so handelt es sich bei der Behandlung "kleiner" Modelle immer um einen wichtigen Teilschritt. — Der vorliegende Text liefert einen Beitrag zur Aufzählung kleiner Modelle für die Theorie der nichtkommutativen assoziativen Algebren: Es wird gezeigt, dass es zu einem vorgegebenen Körper F bis auf Isomorphie nur *eine* nichtkommutative Algebra der Dimension 3 gibt. *ust*

Proof. Let A be a 3-dimensional non-commutative F -algebra. F will be considered as a subalgebra of A . For any $a \in A$, let $L_a: A \rightarrow A$ be defined by $L_a(x) := ax$. The map $a \in A \mapsto L_a \in \text{End } A$ is obviously an injective algebra homomorphism of A into the F -algebra of the endomorphisms of the F -vector space A . We are going to break up the following proof into several small steps.

a) For any $a \in A \setminus F$,

$$F[a] = \{\lambda + \mu a \mid \lambda, \mu \in F\} \cong \begin{cases} F \times F, \\ F[x]/(x^2). \end{cases}$$

Moreover, $F[a]$ contains some $u \in F[a]$ such that $\text{rk } L_u = 1$.

Proof. Since $F[a]$ is commutative, $F[a] \neq A$. On the other hand, $F[a] \neq F$. Therefore, $F[a]$ is 2-dimensional and $(1, a)$ is a basis of $F[a]$. The minimum polynomial $m \in F[x]$ of a over F is of degree 2, and since $F[a] \cong F[x]/(m)$ is not a field, m is reducible. If m has two distinct roots, by the Chinese Remainder Theorem, $F[x]/(m) \cong F \times F$. If m has a double root, $F[x]/(m) \cong F[x]/(x^2)$. Since m is also the minimum polynomial of L_a , L_a has an eigenvalue λ of geometric multiplicity 2. Now $u := a - \lambda \in F[a]$ and $L_u = L_{a-\lambda} = L_a - \lambda \text{id}_A$ is of rank 1.

b) The commutators of A form a 1-dimensional nilpotent ideal C .

Proof. We choose $a, b \in A \setminus F$ such that $F[a] \neq F[b]$. By a), there exist $u \in F[a]$ and $v \in F[b]$ such that L_u and L_v are both of rank 1. Then $(1, u, v)$ is a basis of A . Since $[\alpha + \beta u + \gamma v, \alpha' + \beta' u + \gamma' v] = \beta\gamma'[u, v] + \gamma\beta'[v, u] = (\beta\gamma' - \gamma\beta')[u, v]$ for all $\alpha, \beta, \gamma, \alpha', \beta', \gamma' \in F$, any commutator of A is a multiple of $c := [u, v] \neq 0$. Since $\text{rk } L_c = \text{rk } L_{[u, v]} = \text{rk}(L_u \circ L_v - L_v \circ L_u) \leq 2$, $I := \text{im } L_c = \{cx \mid x \in A\}$ and $K := \ker L_c = \{x \in A \mid cx = 0\}$ are proper non-trivial right ideals of A . For any $a \in A$, $[a, c]$ is a multiple of c , i.e. $[a, c] = \lambda c$ for some $\lambda \in F$. Hence $ac = c(a + \lambda)$ and therefore, I is also a left ideal of A . Similarly, $ca = (a - \lambda)c$ shows that K is a left ideal. Therefore, I and K are ideals and none of them is equal to the zero ideal. Thus the algebras A/I and A/K are of dimension at most 2 and hence commutative. This implies $\text{span}\{c\} \subset I, K$. Since $\dim I + \dim K = 3$, one of the ideals I and K is 1-dimensional and hence equal to $\text{span}\{c\}$. Therefore, $C := \text{span}\{c\}$ is an ideal. Since $c \in K$, $c^2 = 0$ and hence C is nilpotent.

c) A contains an idempotent element unequal to 0 or 1.

Proof. The proof will be carried out by assuming the converse, i.e. that 0 and 1 are the only idempotent elements of A . By a), we obtain $F[a] \cong F[x]/(x^2)$ for each $a \in A \setminus F$. In particular, if $a \in A$ is a non-unit, then $a^2 = 0$ and $\text{rk } L_a \leq 1$. If $a, b \in A$ are any two non-units, $\text{rk } L_{a+b} = \text{rk}(L_a + L_b) \leq \text{rk } L_a + \text{rk } L_b \leq 2$ and hence, $a + b$ is a non-unit. Therefore, $0 = (a + b)^2 = a^2 + ab + ba + b^2 = ab + ba$, i.e. $ab = -ba$. Now we choose non-zero non-units $a, b \in A$ and $c \in C \setminus \{0\}$ such that $F[a], F[b], F[c]$ are distinct in pairs. Then a and bc commute and hence $bc \in F[a] \cap C = \{0\}$. Therefore, b and c commute as $bc = -cb$. Since $(1, b, c)$ is a basis of A , A is commutative and this constitutes a contradiction.

d)

$$A \cong \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & \gamma \end{pmatrix} \mid \alpha, \beta, \gamma \in F \right\}.$$

Proof. Let $c \in C \setminus \{0\}$. By c), there exists some idempotent element $p \in A \setminus F$. Since $pc \in C$, $pc = \xi c$ for some $\xi \in F$. Since $\xi c = pc = p^2c = p(pc) = p(\xi c) = \xi(pc) = \xi(\xi c) = \xi^2c$ and $c \neq 0$, $\xi^2 = \xi$ and hence, $\xi = 0$ or $\xi = 1$. Therefore, $pc \in \{0, c\}$. Similarly, we obtain $cp \in \{0, c\}$. We may assume that $pc = 0$ because otherwise we are going to replace p by $1 - p$. Then $cp = c$ as $cp \neq pc$. The 2-dimensional subspace $U := \text{span}\{c, p\}$ is obviously an ideal and hence $a \in A \mapsto L_{a|U} \in \text{End } U$ is an algebra homomorphism. The matrices representing $L_{x|U}$ with respect to the basis (c, p) of U for $x \in \{1 - p, p, c\}$ are $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. This suffices to ensure the existence of the asserted isomorphism.

The result about rings of order p^3 mentioned at the beginning of this note becomes, of course, a corollary to our Theorem.

Corollary. *For any prime p , there exists, up to isomorphism, exactly one non-commutative ring of order p^3 .*

Proof. If R is a non-commutative ring of order p^3 , then the additive group of R is an elementary-abelian p -group and hence, R is a 3-dimensional non-commutative $\text{GF}(p)$ -algebra.

References

- [1] P. M. Cohn, Algebra, Volume 2, Second Edition, John Wiley & Sons, Chichester-New York-Brisbane-Toronto-Singapore 1989.
- [2] N. Jacobson, Basic Algebra I, W. H. Freeman and Company, San Francisco 1974.

G. Heimbeck

A. H. Kamupingene

University of Namibia

Department of Mathematics

Private Bag 13301

Windhoek, Namibia