

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 51 (1996)

Artikel: An elementary proof of the convergence of iterated exponentials
Autor: Länger, Helmut
DOI: <https://doi.org/10.5169/seals-46960>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

An elementary proof of the convergence of iterated exponentials

Helmut Länger

Helmut Länger, born in 1951, obtained his doctoral degree in mathematics at the University of Technology in Vienna as a student of Professor Nöbauer. After spending a year in Darmstadt he returned to Vienna in 1977 where he is now working as an associate professor at the Institute of Algebra and Discrete Mathematics of the University of Technology. In the beginning of his career his main research interest was algebra, but in recent years he has also done work on discrete problems arising in biology and in axiomatic quantum mechanics.

What are the solutions of the equations

$$x^{x^x} = \frac{1}{2} \text{ or } x^{x^x} = 2,$$

respectively (cf. [2])? Questions of such a type lead in a natural way to the problem of determining the convergence behaviour of the sequence

$$a, a^a, a^{a^a}, \dots \quad (1)$$

for a positive real number a . Here we are concerned with an iteration process with starting point $a > 0$ and iteration function $f(x) := a^x$.

Für welche Werte von a ist der iterative Prozess gegeben durch $x \rightarrow a^x$ stabil? Für welche Werte von a ist die Folge

$$a, a^a, a^{a^a}, a^{a^{a^a}}, \dots$$

konvergent? Und was ist der Limes? Einfache, reizvolle Fragen mit einfachen, reizvollen Antworten! *ust*

Local stability analysis of the iteration process $x \mapsto a^x$ ($a > 0$). Since f is continuous, the possible (finite) limits of (1) must be fixed points of f . Put $g(x) := x^{\frac{1}{x}}$ for all $x > 0$. Then $\lim_{x \rightarrow 0} g(x) = 0$, g is strictly increasing on $(0, e]$, g is strictly decreasing on $[e, \infty)$ and $\lim_{x \rightarrow \infty} g(x) = 1$. Put $h_1 := (g|_{(0, e]})^{-1}$ and $h_2 := (g|_{[e, \infty)})^{-1}$. It is easy to see that the fixed points of f are exactly the positive real numbers b with $g(b) = a$. Moreover, $f'(b) = \ln b$ for every fixed point b of f . Hence we have the following six cases:

- (i) If $a \in (0, \frac{1}{e^e})$ then $h_1(a) \in (0, \frac{1}{e})$ is the unique fixed point of f and $h_1(a)$ is unstable.
- (ii) If $a = \frac{1}{e^e}$ then $h_1(a) = \frac{1}{e}$ is the unique fixed point of f .
- (iii) If $a \in (\frac{1}{e^e}, 1]$ then $h_1(a) \in (\frac{1}{e}, 1]$ is the unique fixed point of f and $h_1(a)$ is asymptotically stable.
- (iv) If $a \in (1, e^{\frac{1}{e}})$ then there exist exactly two fixed points of f , namely $h_1(a) \in (1, e)$ and $h_2(a) \in (e, \infty)$, $h_1(a)$ is asymptotically stable and $h_2(a)$ is unstable.
- (v) If $a = e^{\frac{1}{e}}$ then $h_1(a) = h_2(a) = e$ is the unique fixed point of f .
- (vi) If $a \in (e^{\frac{1}{e}}, \infty)$ then f has no fixed point.

Now we prove the following

Theorem. *Let a be a positive real number. Then the sequence (1) converges iff $a \in [\frac{1}{e^e}, e^{\frac{1}{e}}]$. In this case the corresponding limit is $h_1(a)$.*

Proof. In the following let x, y, z denote (arbitrary) positive real numbers. Put $a_1 := a$, $a_2 := a^a$, $a_3 := a^{a^a}$, \dots . Define $f^2(x) := f(f(x))$. Then we have

$$(f^2)'(x) = a^{a^x} a^x \ln^2 a \quad \text{and} \quad (f^2)''(x) = a^{a^x} a^x (\ln^3 a)(a^x \ln a + 1).$$

Observe that in case $x < y$ we have

$$\begin{aligned} x^z &< y^z \\ z^x &> z^y \quad \text{if } z < 1 \\ z^x &< z^y \quad \text{if } z > 1. \end{aligned}$$

If $a \in (1, e^{\frac{1}{e}}]$ then f is strictly increasing and hence

$$1 < a_1 < a_2 < a_3 < \dots$$

Moreover, $a_n \leq e$ for all n (which can be proved by induction on n), and therefore (1) converges to $h_1(a)$.

Now consider the only non-trivial remaining case $a \in (0, 1)$. Since $a^a < 1^a = 1$, $h_1(a) < 1$ and f is strictly decreasing, we have

$$a_1 < a_3 < a_5 < \dots < h_1(a) < \dots < a_6 < a_4 < a_2. \quad (2)$$

If $a \in (0, \frac{1}{e^e})$ then $f'(h_1(a)) < -1$ and hence (1) is divergent. If $a \in [\frac{1}{e^e}, 1)$ then we have

$$-1 = \ln \frac{1}{e} = \ln h_1\left(\frac{1}{e^e}\right) \leq \ln h_1(a) = a^{h_1(a)} \ln a < a^u \ln a < 0$$

for all $u > h_1(a)$ and hence $(f^2)'' < 0$ on $(h_1(a), \infty)$. Therefore

$$0 < (f^2)'(u) < (f^2)'(h_1(a)) = \ln^2 h_1(a) \leq 1$$

for all $u > h_1(a)$ which shows that the sequence a_2, a_4, a_6, \dots and therefore also (1) converges to $h_1(a)$.

Remarks.

- (i) The sequence $\frac{1}{4}, \frac{1}{4}^{\frac{1}{4}}, \frac{1}{4}^{\frac{1}{4}^{\frac{1}{4}}}, \dots$ converges to $\frac{1}{2}$, i. e. $x^{x^x} = \frac{1}{2}$ has the unique solution $x = \frac{1}{4}$.
- (ii) The sequence $\sqrt{2}, \sqrt{2}^{\sqrt{2}}, \sqrt{2}^{\sqrt{2}^{\sqrt{2}}}, \dots$ converges to 2 (cf. [2]), i. e. $x^{x^x} = 2$ has the unique solution $x = \sqrt{2}$ which is somewhat surprising.
- (iii) If (1) contains an element $\geq e$ then (1) is divergent.
- (iv) If $a \in (0, \frac{1}{e^e})$ then (1) converges to a limit cycle of order 2.

Acknowledgement. The author is indebted to Dipl.-Ing. H. Postl, Vienna, and Dr. J. Wiesenbauer, TU Vienna, for valuable discussions on the topic of the paper and to the referee for some remarks improving the quality of the paper.

References

- [1] J. Elstrodt: Iterierte Potenzen. Math. Sem.-Ber. 41 (1994), 167–178.
- [2] M. C. Mitchelmore: A matter of definition. Amer. Math. Monthly 81 (1974), 643–647.

Helmut Länger

Technische Universität Wien

Institut für Algebra und Diskrete Mathematik

Abteilung für Mathematik in den Naturwissenschaften

und Mathematische Biologie

Wiedner Hauptstraße 8–10

A-1040 Wien