Elemente der Mathematik
Schweizerische Mathematische Gesellschaft
51 (1996)
On a model of plane geometry
Powers, R.C. / Riedel, T. / Sahoo, P.K.
https://doi.org/10.5169/seals-46969

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. <u>Siehe Rechtliche Hinweise.</u>

# **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. <u>Voir Informations légales.</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. <u>See Legal notice.</u>

**Download PDF: 21.05.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# **On a Model of Plane Geometry**

# R. C. Powers, T. Riedel, and P. K. Sahoo

Robert C. Powers: In 1988, I obtained my doctorate in mathematics at the University of Massachusetts under the supervision of M. F. Janowitz. My interests in mathematics include ordered sets, discrete mathematics, functional equations and geometry. Outside of mathematics, I enjoy spending time with my wife and baby daughter.

Thomas Riedel: After studying physics at Eberhard-Karls Universität Tübingen (Germany), I received my PhD (mathematics) from the University of Massachusetts, Amherst, in 1990 under the direction of B. Schweizer. Most of my work is on functional equations and their applications to probabilistic metric spaces, information theory, geometry and numerical analysis. I am also interested in computers (including their use in education) and physics.

Prasanna K. Sahoo: I obtained my doctorate in mathematics from the University of Waterloo in 1986. My primary research area is functional equations and their applications to areas such as geometry, numerical analysis and economics. Occasionally I work on image thresholding and image compression, and I am mainly interested in applying information theoretic techniques to threshold an image.

In [3], Grünbaum and Mycielski proposed the following model of plane geometry. The points of this model are the points of the Euclidean plane  $\mathbb{R}^2$ . There are four types of lines for this model: a vertical Euclidean line; a horizontal Euclidean line; a translate of the hyperbola  $L = \{(x, y) : x > 0, y = 1/x\}$ ; and a translate of the hyperbola

Modelle der ebenen Geometrie, die mit Ausnahme des Parallelenaxioms alle anderen erfüllen, haben für die Entwicklung der Mathematik eine ausserordentlich wichtige Rolle gespielt. Ihre Entdeckung durch Beltrami, Klein und Poincaré zeigte ja nicht nur die logische Unabhängigkeit des Parallelenaxioms, sondern sie machte auch klar, dass die Mathematik Axiome weitgehend unabhängig von irgendwelchen Bezügen zur Wirklichkeit setzen kann. Ausser den berühmten Modellen von Klein und Poincaré werden jedem Studierenden der Mathematik auch andere vorgeführt, die nur einen Teil der Axiome der ebenen Geometrie erfüllen; die endlichen affinen und projektiven Ebenen sind für diese Zwecke besonders beliebt. Unendliche Modelle sind weniger bekannt. Der vorliegende Beitrag von Powers, Riedel und Sahoo beschäftigt sich mit einer ganzen Klasse von einfach zu beschreibenden unendlichen Modellen dieser Art. Die Autoren gehen dabei auch der Frage nach, unter welchen Bedingungen zwei Modelle ihrer Klasse zueinander isomorph sind. ust  $L^* = \{(x, y) : x < 0, y = -1/x\}$ . We will follow [3] and label this model G2. Notice that G2 does not satisfy the Euclidean, hyperbolic, or elliptic parallel postulates. Thus G2 makes a nice example for students who take a geometry-for-teachers course.



Fig. 1 The lines of G2

There is no reason why one needs to use the hyperbolas 1/x and -1/x (plus translates) as lines of G2. Indeed, if we replace 1/x with  $1/x^2$  and -1/x with  $-1/x^2$ , then we generate yet another model of plane geometry that does not satisfy any of the standard parallel postulates. In [3], the authors proposed the curves  $e^{-x}$  and  $e^x$  as yet another version of model G2. One might think that all these versions of G2 are essentially the same. As our theorem below demonstrates, these versions are not isomorphic as models of incidence geometry. This result parallels the one given in [3] and further developed in [2] and [4].

Let  $f : I \to \mathbb{R}$  and  $g : J \to \mathbb{R}$  be two functions where I and J are subintervals of  $\mathbb{R}$ . We propose the following generalization of G2 using the functions f and g. As before, the points of this generalization of G2 are the points of  $\mathbb{R}^2$ . There are four types of lines: a vertical Euclidean line; a horizontal Euclidean line; a translate of the graph  $\{(x, y) : x \in I, y = f(x)\}$ ; and a translate of the graph  $\{(x, y) : x \in J, y = g(x)\}$ . We denote this interpretation of incidence geometry by  $M_{(f,g)}$ .

In order for  $M_{(f,g)}$  to be a model of incidence geometry we need f and g to be one-to-one. In particular, if these functions are continuous then one needs to be strictly decreasing and the other needs to be strictly increasing. Thus, we will require f(x) to be strictly decreasing on I and g(x) to be strictly increasing on J. If y = f(x) is bounded above by y = m and below by the line y = n, then there is no line in  $M_{(f,g)}$  that passes through the points (0, m + 1) and (1, n - 1). Therefore, we will require f(x) and g(x) to be unbounded from either above or below. It is possible that f and g are unbounded in both directions (e.g.,  $f(x) = \log(-x)$  for x < 0 and  $g(x) = \log(x)$  for x > 0) or in different directions (e.g., f(x) = 1/x for x > 0 and g(x) = -1/x for x > 0). Our goal, however, is to determine those functions f(x) and g(x) such that  $M_{(f,g)}$  and G2 are isomorphic as models of incidence geometry. Toward this end, we will require f(x) and g(x) to be continuous, unbounded from above and bounded from below, and bounded on one side by vertical asymptotes.

**Lemma 1** Let  $F : \mathbb{R}^2 \to \mathbb{R}^2$  be a bijection that induces a map from  $M_{(f,g)}$  onto  $M_{(f,g)}$ . Then

$$F(x,y) = (f_1(x), f_2(y))$$
(1)

for some bijections  $f_1$  and  $f_2$  of  $\mathbb{R}$ .

*Proof* The following geometric fact is clear in  $M_{(f,g)}$ . If L is a line and p is a point not on L such that there is exactly one line through p parallel to L, then L is a horizontal line. It follows from this fact that the family of horizontal lines is mapped to itself. Since a line in  $M_{(f,g)}$  is vertical if and only if it intersects every horizontal line it follows that F maps the family of vertical lines to the family of vertical lines. It now follows that F(x, y) has the form described by equation (1).

We point out that  $G_2 = M_{(1/x, -1/x)}$  satisfies Lemma 1. In fact, we can prove more.

**Lemma 2**  $F : \mathbb{R}^2 \to \mathbb{R}^2$  is a bijection that induces a map from G2 onto G2 if and only if

$$F(x,y) = \begin{pmatrix} \lambda & 0\\ 0 & 1/|\lambda| \end{pmatrix} \begin{pmatrix} x\\ y \end{pmatrix} + \begin{pmatrix} \alpha\\ \beta \end{pmatrix}$$
(2)

for some nonzero  $\lambda \in \mathbb{R}$  and  $\alpha, \beta \in \mathbb{R}$ .

*Proof* By Lemma 1, we know that  $F(x, y) = (f_1(x), f_2(y))$  for some bijections  $f_1$  and  $f_2$  of  $\mathbb{R}$ . Since F maps G2 onto G2 it follows that  $\{F(x, y)|x > a, y = 1/(x - a) + b\}$  is a line of G2. Since  $\{F(x, y)|x > a, y = 1/(x - a) + b\}$  is neither a horizontal nor vertical line it follows that it is a translate of either  $\{(x, y)|x > 0, y = 1/x\}$  or  $\{(x, y)|x < 0, y = -1/x\}$ . Thus, there exist  $a', b' \in \mathbb{R}$  such that

$$(f_1(x) - a')(f_2(y) - b') = 1 \text{ or } -1$$
 (3)

for x > a and y = 1/(x - a) + b. If the above product is 1, then

$$(f_1(x) - a')(f_2(y) - b') = (x - a)(y - b),$$

and so

$$\frac{f_1(x) - a'}{x - a} = \frac{y - b}{f_2(y) - b'}$$

Therefore, there exists a nonzero constant  $\lambda$  such that

$$\frac{f_1(x)-a'}{x-a}=\lambda=\frac{y-b}{f_2(y)-b'}.$$

Thus,  $f_1(x) = \lambda x + \alpha$  and  $f_2(y) = \frac{1}{\lambda}y + \beta$  for some constants  $\alpha$  and  $\beta$  and for all x > a and y = 1/(x-a) + b. Since the line  $\{(f_1(x), f_2(y)) | x > a, y = 1/(x-a) + b\}$  is

bounded below it follows that  $\lambda > 0$ . If the product in equation (3) is -1, then a similar argument will show that there exists a constant  $\lambda < 0$  such that  $f_1(x) = \lambda x + \alpha$  and  $f_2(y) = \frac{1}{(-\lambda)}y + \beta$  for some constants  $\alpha$  and  $\beta$ , for all x > a, and y = 1/(x-a) + b. Since a and b can be chosen arbitrarily it follows that  $f_1(x) = \lambda x + \alpha$  and  $f_2(y) = \frac{1}{(-\lambda)}y + \beta$  for all x and y. Hence F(x, y) satisfies (2).

It is straightforward to verify that if  $F : \mathbb{R}^2 \to \mathbb{R}^2$  is a bijection of the form described by equation (2), then F induces a map from G2 onto G2.

We can now prove our main result.

**Theorem 3** If there exists a continuous bijection  $\phi$  of  $\mathbb{R}^2$  onto itself which induces a map from G2 onto  $M_{(f,g)}$ , then, up to translations, f(x) = a/x for x > 0 and g(x) = -a/x for x < 0 for some positive constant a.

*Proof* The argument used to establish equation (1) in Lemma 1 can be used to show that, for all  $(x, y) \in \mathbb{R}^2$ ,

$$\phi(x, y) = (\phi_1(x), \phi_2(y))$$
(4)

for some bijections  $\phi_1$  and  $\phi_2$  of  $\mathbb{R}$ . Now consider the family of automorphisms of  $M_{(f,g)}$  given by  $\tau_t(x,y) = (x+t,y)$  where  $t \in \mathbb{R}$ . Then, for each t,  $\tau'_t = \phi^{-1}\tau_t\phi$  is an automorphism of G2. It follows from Lemma 2 that

$$\phi_1^{-1}(\phi_1(x) + t) = \lambda x + a$$
(5)

for some nonzero constant  $\lambda$ . If  $\lambda \neq 1$ , then the equation  $x = \lambda x + a$  has  $x = \frac{a}{1-\lambda}$  as a solution. If  $x = \frac{a}{1-\lambda}$  then equation (5) becomes

$$\phi_1(x) + t = \phi_1(x),$$

and so, t = 0. So for  $t \neq 0$ , it follows that  $\lambda = 1$ . Hence

$$\tau'_t = (x + h(t), a(t)y + b(t))$$

where h, a, b are functions and where h satisfies the Cauchy Functional Equation h(r+s) = h(r) + h(s) for all  $r, s \in \mathbb{R}$ . Since  $\phi \circ \tau'_t = \tau_t \circ \phi$  it follows that  $\phi_1(x+h(t)) = \phi_1(x) + t$ . The last equation holds if we replace  $\phi_1(x)$  by  $\phi_1(x) - \phi_1(0)$ , thus we can assume that  $\phi_1(0) = 0$ . If we set x = 0, then we get  $\phi_1(h(t)) = t$  for all t. Hence  $\phi_1 = h^{-1}$  where h is continuous. It follows from Cauchy's Functional Equation (see [1]) that h(t) = kt for some nonzero constant k. Hence  $\phi_1(x) = \frac{1}{k}x$  for all x. A similar argument will show that, for all  $y, \phi_2(y) = k'y$  for some nonzero constant k'. It now follows that equation (4) has, up to composition of a translation, the form

$$\phi(x,y) = (k_1 x, k_2 y)$$

for some nonzero constants  $k_1$  and  $k_2$ .

The line  $L = \{(x,y)|x > 0, y = 1/x\}$  in G2 is mapped by  $\phi$  to the line  $\{(k_1x, k_2y)|x > 0, y = 1/x\}$  in  $M_{(f,g)}$ . The latter must be a translate of either the graph of f or the graph of g. If it is a translate of the graph of f, then there exist  $\alpha$  and  $\beta$  such that

$$k_2 y = f(k_1 x + \alpha) + \beta$$

for all x > 0 and y = 1/x. Since f(x) is assumed to be bounded below and y ranges over the interval  $(0, \infty)$  it follows that  $k_2 > 0$ . This in turn forces  $k_1 > 0$ , since y = 1/x (for x > 0) and f are strictly decreasing functions. Replacing x by  $\frac{x-\alpha}{k_1}$  in the last equation leads to

$$f(x) = \frac{k_1 k_2}{x - \alpha} - \beta \tag{6}$$

for all  $x > \alpha$ .

Note that  $\phi$  maps a translate of  $L = \{(x, y) | x > 0, y = 1/x\}$  to some translate of the graph of f. Since  $\phi$  maps G2 onto  $M_{(f,g)}$ , it follows that  $\phi$  maps

$$L^* = \{(x, y) | x < 0, y = -1/x\}$$

to a translate of the graph of g. Thus

$$g(x) = \frac{-k_1 k_2}{x - \alpha'} - \beta' \tag{7}$$

for all  $x < \alpha'$ . In fact, it is easy to see that  $\alpha = \alpha'$  and  $\beta = \beta'$ .

If the image of L under  $\phi$  is a translate of the graph of g then, using the same type of argument as above, we get  $k_1 < 0$ ,  $k_2 > 0$ ,  $f(x) = \frac{-k_1k_2}{x-\alpha} - \beta$  for all  $x > \alpha$ , and  $g(x) = \frac{k_1k_2}{x-\alpha'} - \beta'$  for all  $x < \alpha'$ .

So f(x) and g(x) are just scaled translations of 1/x. Given that  $M_{(f,g)}$  is closed under translations, we can take f(x) = a/x for x > 0 and g(x) = -a/x for x < 0 where a is some positive constant.

In conclusion, we note that the converse of Theorem 3 is true since, for a > 0,  $\phi(x, y) = (\sqrt{ax}, \sqrt{ay})$  is a bijection of  $\mathbb{R}^2$  which induces a map from G2 onto  $M_{(a/x, -a/x)}$ .

# References

- [1] J. Aczel and J. Dhombres, *Functional equations in several variables*, Cambridge University Press, Cambridge, 1989.
- [2] V. Faber, M. Kuczma and J. Mycielski, Some models of plane geometries and a functional equation, Colloq. Math. 62 (1991), 279–281.
- [3] B. Grünbaum and J. Mycielski, Some models of plane geometry, Amer. Math. Monthly 97 (1990), 839-846.
- [4] R. Powers, T. Riedel and P. K. Sahoo, Some models of geometries and a functional equation, Colloq. Math. 66 (1993), 165–173.

R. C. Powers, T. Riedel, and P. K. Sahoo Department of Mathematics University of Louisville Louisville, KY 40292, USA