

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 52 (1997)

Artikel: Best Lower and Upper Approximates to Irrational Numbers
Autor: Kimberling, Clark
DOI: <https://doi.org/10.5169/seals-2667>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Best Lower and Upper Approximates to Irrational Numbers

Clark Kimberling

Clark Kimberling received his Ph.D. in 1970 from the Illinois Institute of Technology. Aside from number theory, his interests include triangle geometry, playing musical instruments, and composing.

Wherever fractional notation of the form p/q occurs in this paper, p and q denote relatively prime positive integers. The rational number p/q is a best approximate to α (e.g., Lang [L]) if for every b/c having denominator $c < q$,

$$|q\alpha - p| < |c\alpha - b|. \quad (1)$$

These conditions imply $q|\alpha - p/q| < c|\alpha - b/c| < q|\alpha - b/c|$, so that

$$|\alpha - p/q| < |\alpha - b/c|, \quad (2)$$

which is to say that p/q is nearer α than any b/c having $c < q$. However, (1) is stronger than (2), as exemplified by $(\alpha, p/q, b/c) = (1, 3/5, 1/2)$; in other words “best approximate” is “better” than “nearest approximate”.

Given a positive irrational number α , the principal convergents p_i/q_i to α are well known to be the best approximates to α ; this theorem lends itself to a lemma proved in [L]:

Lemma 1 *Suppose $p_0/q_0, p_1/q_1, \dots$ are the principal convergents to a positive irrational number α . If b/c satisfies $|c\alpha - b| < |q_i\alpha - p_i|$, then $c \geq q_{i+1}$.*

Die Approximation irrationaler Zahlen durch rationale Zahlen spielt in der Zahlentheorie eine grosse Rolle. Erinnert sei zum Beispiel an den berühmten Satz von Liouville; er liefert die Möglichkeit, die Transzendenz einer reellen Zahl mit Hilfe ihrer rationalen Approximationen zu beweisen. Clark Kimberling konstruiert im vorliegenden Beitrag auf einfache Weise beste untere und obere rationale Approximationen und setzt sie in Beziehung zu den bereits von Perron eingeführten “besten” und “nächsten Näherungen”. usf

We modify inequality (1), calling p/q a best lower approximate to α if $p/q < \alpha$ and for every $b/c < \alpha$ having $c < q$,

$$q\alpha - p < c\alpha - b; \quad (1L)$$

and calling p/q a best upper approximate to α if $p/q > \alpha$ and for every $b/c < \alpha$ having $c < q$,

$$p - q\alpha < b - c\alpha. \quad (1U)$$

Before solving for p/q in these cases, we note that the analogous problem for “nearest” lower and upper approximates is solved in Perron [P, pp. 55–63], where, ironically, they are called “beste Näherungen”. Perron’s solutions are the same as those obtained below — a surprise in view of the aforementioned nonequivalence of “nearest” and “best”.

For irrational x , let $\|x\|$ denote the distance from x to the integer nearest to x . Let $[x]$ denote the greatest integer $\leq x$, and define $((x))$, the fractional part of x , by $((x)) = x - [x]$. Then

$$\|x\| = \begin{cases} ((x)) & \text{if } ((x)) < 1/2 \\ 1 - ((x)) & \text{otherwise,} \end{cases} \quad (3)$$

and (1) can be written as $\|q\alpha\| < \|c\alpha\|$.

Next we recall some basics about continued fractions, principal convergents, and intermediate convergents. Suppose α has continued fraction $[\![a_0, a_1, a_2, \dots]\!]$, and let

$$p_{-2} = 0, \quad p_{-1} = 1, \quad p_i = a_i p_{i-1} + p_{i-2}$$

$$\text{and} \quad q_{-2} = 1, \quad q_{-1} = 0, \quad q_i = a_i q_{i-1} + q_{i-2}$$

for $i \geq 0$. The principal convergents of α are the rational numbers p_i/q_i for $i \geq 0$.

Now for all nonnegative integers i and j , define

$$p_{i,j} = jp_{i+1} + p_i \text{ and } q_{i,j} = jq_{i+1} + q_i.$$

The fractions

$$\frac{p_{i,j}}{q_{i,j}} = \frac{jp_{i+1} + p_i}{jq_{i+1} + q_i}, \quad 1 \leq j \leq a_{i+2} - 1, \quad (4)$$

are the i -th intermediate convergents of α . As proved in [L, p. 16],

$$\dots < \frac{p_i}{q_i} < \dots < \frac{p_{i,j}}{q_{i,j}} < \frac{p_{i,j+1}}{q_{i,j+1}} < \dots < \frac{p_{i+2}}{q_{i+2}} < \dots \quad \text{if } i \text{ is even,} \quad (5)$$

$$\dots > \frac{p_i}{q_i} > \dots > \frac{p_{i,j}}{q_{i,j}} > \frac{p_{i,j+1}}{q_{i,j+1}} > \dots > \frac{p_{i+2}}{q_{i+2}} > \dots \quad \text{if } i \text{ is odd,} \quad (6)$$

and $p_{i,j-1}q_{ij} - p_{ij}q_{i,j-1} = (-1)^j$ for $i = 0, 1, 2, \dots$ and $j = 1, 2, \dots, a_{i+2} - 1$. If the range of j in (4) is extended to $0 \leq j \leq a_{i+2} - 1$, then the principal convergents are included among the intermediate convergents. We shall refer to both kinds as simply convergents, those in (5) as even-indexed convergents, and those in (6) as odd-indexed convergents.

Suppose now that $q \geq 1$. Taking $x = q\alpha$ in (3) gives $\|q\alpha\| = |q\alpha - p|$, where

$$p = \begin{cases} \lfloor q\alpha \rfloor & \text{if } ((q\alpha)) < 1/2 \\ \lfloor q\alpha \rfloor + 1 & \text{otherwise,} \end{cases} \quad \text{so that} \quad \|q\alpha\| = \begin{cases} ((q\alpha)) & \text{if } ((q\alpha)) < 1/2 \\ 1 - ((q\alpha)) & \text{otherwise.} \end{cases}$$

Lemma 2 If p_i/q_i are the principal convergents to a positive irrational number α and i is even, then $((q_{i+2}\alpha)) < 1/2 < ((q_{i+1}\alpha))$,

$$((jq_{i+1}\alpha)) = j((q_{i+1}\alpha)) - j + 1, \quad (7)$$

and

$$((jq_{i+1}\alpha)) + ((q_i\alpha)) > 1 \quad (8)$$

for $j = 1, 2, \dots, a_{i+2} - 1$.

Proof. The first assertion merely expresses the fact that the integer nearest $q_i\alpha$ is p_i for even $i > 0$, and that the integer nearest $q_{i+1}\alpha$ is $p_i + 1$. Continuing, it is well known ([L, p. 8]) that

$$p_{i+1} - q_{i+1}\alpha < 1/q_{i+2},$$

so that $((q_{i+1}\alpha)) > 1 - 1/q_{i+2}$. Now $a_{i+2} \leq a_{i+2}q_{i+1} + q_i = q_{i+2}$, whence

$$0 < j((q_{i+1}\alpha)) - j + 1 < 1 \text{ for } 1 \leq j \leq a_{i+2} - 1.$$

Since $((jq_{i+1}\alpha))$ is an irrational number having the same fractional part as $j((q_{i+1}\alpha)) - j + 1$, identity (7) is proved. Continuing,

$$p_{i+2}q_{i+1} - q_{i+2}p_{i+1} = -1 < 0 = p_{i+2}q_{i+2} - q_{i+2}p_{i+2},$$

so that

$$\alpha > \frac{p_{i+2}}{q_{i+2}} > \frac{p_{i+2} - p_{i+1}}{q_{i+2} - q_{i+1}}.$$

Then $\alpha(a_{i+2}q_{i+1} + q_i - q_{i+1}) > a_{i+2}p_{i+1} + p_i - p_{i+1}$, which implies

$$\frac{((q_i\alpha))}{1 - ((q_{i+1}\alpha))} = \frac{q_i\alpha - p_i}{p_{i+1} - q_{i+1}\alpha} > a_{i+2} - 1.$$

Thus for $0 \leq j \leq a_{i+2} - 1$, we have

$$j(1 - ((q_{i+1}\alpha))) < ((q_i\alpha)),$$

so that

$$j((q_{i+1}\alpha)) - j + 1 + ((q_i\alpha)) > 1,$$

and (8) follows from (7). \square

Lemma 3 If p_{ij}/q_{ij} are the convergents to a positive irrational number α and i is even, then

$$((q_{i0}\alpha)) > ((q_{i1}\alpha)) > ((q_{i2}\alpha)) > \dots > ((q_{i,a_{i+2}-1}\alpha)) > 1 - ((q_{i+1,0}\alpha)) > ((q_{i+2,0}\alpha)).$$

Proof. When $j = 0$, we are dealing with principal convergents, hence best approximates to α , so that

$$\|q_{i0}\alpha\| > \|q_{i+1,0}\alpha\| > \|q_{i+2,0}\alpha\|,$$

or equivalently,

$$((q_{i0}\alpha)) > 1 - ((q_{i+1,0}\alpha)) > ((q_{i+2,0}\alpha)).$$

Next, using Lemma 2, we find for $j = 0, 1, \dots, a_{i+2} - 2$ that

$$((jq_{i+1}\alpha)) = j((q_{i+1}\alpha)) - j + 1 > (j+1)((q_{i+1}\alpha)) - (j+1) + 1 = (((j+1)q_{i+1}\alpha)),$$

so that $((q_{ij}\alpha)) > ((q_{i,j+1}\alpha))$. Finally,

$$1 - ((q_{i+1}\alpha)) = p_{i+1} - q_{i+1}\alpha < 1/q_{i+2} < 1/a_{i+2},$$

whence $a_{i+2}((q_{i+1}\alpha)) > a_{i+2} - 1$, and

$$((q_{i,a_{i+2}-1}\alpha)) = (a_{i+2} - 1)((q_{i+1}\alpha)) - (a_{i+2} - 1) + 1 > 1 - ((q_{i+1}\alpha)). \quad \square$$

Theorem 1 *The best lower approximates to a positive irrational number α are the even-indexed convergents to α .*

Proof. Suppose q_{ij} is an even-indexed convergent, and c is a positive integer such that $((c\alpha)) < ((q_{ij}\alpha))$. We wish to show that $c > q_{ij}$. By Lemma 3, $((c\alpha)) < ((q_i\alpha))$, so that $\|c\alpha\| < \|q_i\alpha\|$, which by Lemma 1 implies $c \geq q_{i+1}$. If $c = q_{i+1}$, then $((c\alpha)) = ((q_{i+1}\alpha))$, which by Lemma 3 implies

$$((c\alpha)) > 1 - ((q_{ij}\alpha)) > ((q_{ij}\alpha)),$$

a contradiction. Also, clearly, $c \neq q_{ij}$, so it remains to consider the possibility that $q_{i+1} < c < q_{ij}$; write $c = mq_{i+1} + h$, where $1 \leq h < q_{i+1}$ and $1 \leq m \leq j$. Then

$$((c\alpha)) = ((mq_{i+1}\alpha + h\alpha)),$$

and, using Lemma 2,

$$((mq_{i+1}\alpha)) + ((h\alpha)) - 1 = ((c\alpha)) < ((q_{ij}\alpha)) = ((jq_{i+1}\alpha + q_i\alpha)) = ((jq_{i+1}\alpha)) + ((q_i\alpha)) - 1,$$

so that

$$((h\alpha)) < ((jq_{i+1}\alpha)) - ((mq_{i+1}\alpha)) + ((q_i\alpha)).$$

Identity (7) easily gives $((jq_{i+1}\alpha)) \leq ((mq_{i+1}\alpha))$, so that $((h\alpha)) < ((q_i\alpha))$, and by Lemma 1, $h > q_{i+1}$, a contradiction. \square

Lemma 4 *Let p_i/q_i denote the principal convergents to a positive irrational number $\alpha = [a_0, a_1, a_2, \dots]$, and let p'_i/q'_i denote the principal convergents to the number $\alpha' = a_0 + 1 - \alpha$. If $\alpha - a_0 < 1/2$, then $p'_0/q'_0 = 0/1$, $p'_1/q'_1 = 1/1$, and*

$$\frac{p'_i}{q'_i} = \frac{(a_0 + 1)q_{i-1} - p_{i-1}}{q_{i-1}}$$

for $i = 2, 3, \dots$. If $\alpha - a_0 > 1/2$, then $a_1 = 1$, $p'_0/q'_0 = 0/1$, $p'_1/q'_1 = 1/(a_2 + 1)$, and

$$\frac{p'_i}{q'_i} = \frac{(a_0 + 1)q_{i+1} - p_{i+1}}{q_{i+1}}$$

for $i = 2, 3, \dots$

Proof. It is easy to verify that

$$\alpha' = \begin{cases} [0, 1, a_1 - 1, a_2, a_3, a_4, \dots] & \text{if } \alpha - a_0 < 1/2 \\ [0, a_2 + 1, a_3, a_4, \dots] & \text{if } \alpha - a_0 > 1/2. \end{cases}$$

The rest of the proof is routine and omitted. \square

Theorem 2 *The best upper approximates to a positive irrational number α are the odd-indexed convergents to α .*

Proof. Suppose p_{ij}/q_{ij} is an odd-indexed convergent to α . Suppose also, contrary to (1U), that there exists b/c , with denominator $c < q_{ij}$, such that

$$p_{ij} - q_{ij}\alpha \geq b - c\alpha.$$

Substituting $a_0 + 1 - \alpha'$ for α and using principal convergents, we then have

$$\begin{aligned} (jp_{i+1} + p_i) - (jq_{i+1} + q_i)(a_0 + 1 - \alpha') &\geq b - c(a_0 + 1 - \alpha'), \\ (jq_{i+1} + q_i)\alpha' - \left(j((a_0 + 1)q_{i+1} - p_{i+1}) + (a_0 + 1)q_i - p_i\right) &\geq c\alpha' - (ca_0 + c - b). \end{aligned} \tag{9}$$

If $\alpha - a_0 < 1/2$, then by Lemma 4, inequality (9) can be written as

$$(jq'_{i+2} + q'_{i+1})\alpha' - (jp'_{i+2} + p'_{i+1}) \geq c\alpha' - (ca_0 + c - b),$$

so that

$$q'_{i+1,j}\alpha' - p'_{i+1,j} \geq c\alpha' - (ca_0 + c - b),$$

contrary to Theorem 1, since $p'_{i+1,j}/q'_{i+1,j}$ is an even-indexed convergent to α' and

$$c < q_{ij} = jq_{i+1,j} + q_i = jq'_{i+2,j} + q'_{i+1} = q'_{i+1,j}.$$

On the other hand, if $\alpha - a_0 > 1/2$, then (9) can be written as

$$(jq'_i + q'_{i-1})\alpha' - (jp'_i + p'_{i-1}) \geq c\alpha' - (ca_0 + c - b),$$

so that

$$q'_{i-1,j}\alpha' - p'_{i-1,j} \geq c\alpha' - (ca_0 + c - b),$$

contrary to Theorem 1. \square

References

- [L] Serge Lang, *Introduction to Diophantine Approximations*, Addison-Wesley, Reading, Mass., 1966.
- [P] Oskar Perron, *Die Lehre von den Kettenbrüchen*, Chelsea, New York, 1950.

Clark Kimberling
 Department of Mathematics
 University of Evansville
 1800 Lincoln Avenue
 Evansville, Indiana 47722