

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 55 (2000)

Artikel: A Property of Euler's Elastic Curve
Autor: Moll, Victor H. / Neill, Pamela A. / Nowalsky, Judith L.
DOI: <https://doi.org/10.5169/seals-5638>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 03.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A Property of Euler's Elastic Curve

Victor H. Moll, Pamela A. Neill,
Judith L. Nowalsky, Leonardo Solanilla

Victor H. Moll was born in Santiago, Chile. He completed his undergraduate education at the Universidad Santa Maria in Valparaiso. He studied under Henry McKean at the Courant Institute and joined the Department of Mathematics at Tulane University in the wonderful city of New Orleans. His current mathematical interests lie in Symbolic Computation and the evaluations of definite integrals. His work can be found in <http://www.math.tulane.edu:80/vhm>.

Pamela Andrea F. Neill is a first generation American. She graduated from the University of New Orleans, May 1979, with a BS in Civil Engineering. Then she worked in foundations for the US Corps of Engineers for 7 years and is now an assistant professor at Delgado Community College in New Orleans. She received an MST in Mathematics from Loyola University December, 1992.

Judith L. Nowalsky was born in New Orleans, Louisiana. She graduated from Newcomb College of Tulane University in 1981 with a degree in Economics. Later she obtained a Master of Science in Teaching at Loyola in 1992. She obtained a MS in Mathematics at Tulane in 1998.

Leonardo Solanilla was born in Ibaguè, Tolima, Colombian Andes. He grew up wild and spoiled by the love of his family. From 1981 to 1985 he attended La Universidad de los Andes in Bogotà, after which he received his BS in Electrical Engineering. He enrolled in the doctoral program at Tulane University in 1993, getting his PhD in 1999. As this note is being written, he holds a postdoctoral position in the Instituto de Física y Matemáticas at the Universidad Michoacana de San Nicolás in the sunny Morelia, Mexico.

Die Fourierreentwicklung glatter, periodischer Funktionen dürfte den meisten Leserinnen und Lesern bekannt sein. Das Studium komplexer, doppeltperiodischer Funktionen führt auf die elegante Theorie der Weierstrass'schen \wp -Funktion. Deren Umkehrfunktionen geben Anlass zu den elliptischen Integralen, welche – historisch gesehen – am Anfang der Entwicklung standen. Fagnano, Euler, Legendre und Gauss haben wesentliche Beiträge dazu geleistet. Erst Abel und Jacobi führten – unabhängig voneinander, wie die Korrespondenz zwischen A.-M. Legendre und C.G.J. Jacobi belegt – die elliptischen Funktionen ein. Der vorliegende Beitrag gibt zunächst einen Überblick über die Untersuchungen von Euler und Legendre über elliptische bzw. lemniskatische Integrale und schliesst mit einer Verallgemeinerung der klassischen Formel von Legendre. *jk*

1 Introduction

During the first two decades of the 19th century, Legendre developed the theory of elliptic integrals. His work [5] appeared in 1811 and his monumental treatise [6] in 1825. Shortly after that, Abel published his work [1] on the inversion of elliptic integrals and on the properties of the elliptic functions defined by this procedure. One of Legendre's most elegant formulae appears on [5] page 61. This is his famous relation:

$$\begin{aligned} & \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} \times \int_0^1 \sqrt{\frac{1-(k')^2x^2}{1-x^2}} dx + \\ & \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-(k')^2x^2)}} \times \int_0^1 \sqrt{\frac{1-k^2x^2}{1-x^2}} dx - \\ & \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} \times \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-(k')^2x^2)}} = \frac{\pi}{2}. \end{aligned} \quad (1.1)$$

The terms in (1.1) are the classical elliptic integrals that made their debut in the calculation of the length of the ellipse and the lemniscate. The reader is referred to [7] for details on this topic and to [2] for the history of Legendre's relation (1.1).

The lemniscatic integral ((1.3), below) appears in the calculation of the arclength of the lemniscate of equation $(x^2 + y^2)^2 = a^2(x^2 - y^2)$. Siegel [8] makes this example his starting point in his book on abelian functions. The parametrization of the lemniscate

$$x = \sqrt{\frac{r^2 + r^4}{2}} \quad \text{and} \quad y = \sqrt{\frac{r^2 - r^4}{2}}, \quad (1.2)$$

with $r = \sqrt{x^2 + y^2}$, yields the expression

$$L = \int_0^1 \frac{dx}{\sqrt{1-x^4}} \quad (1.3)$$

for the total arclength. This lemniscatic integral was studied by Euler in [4] and is the special case $k = \sqrt{-1}$ of the *elliptic integral of the first kind*

$$K(k) := \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}$$

later studied by Legendre in [6]. In this case (1.1) becomes

$$\int_0^1 \frac{dx}{\sqrt{1-x^4}} \times \int_0^1 \frac{x^2}{\sqrt{1-x^4}} dx = \frac{\pi}{4}. \quad (1.4)$$

In this paper we describe Euler's method to prove (1.4) and establish a generalization that deals with the elastic curve

$$f_n(x) := \int_0^x \frac{t^n}{\sqrt{1-t^{2n}}} dt$$

for which we prove that

$$R_n \times L_n = \frac{\pi}{2n},$$

where $R_n = f_n(1)$ is the so-called *main radius*, and L_n is the length of the curve from $x = 0$ to $x = 1$. The special case $n = 2$ yields Euler's result.

Section 2 recalls a standard proof of (1.1) based on the fact that the Legendre integrals satisfy a differential equation. Section 3 describes Euler's original proof, its generalization and discusses the issue of convergence, a fact that Euler was happy to ignore. Although Euler did not explicitly address the issue of convergence in [3], his familiarity with Stirling's formula dates from at least 1736.

2 Legendre's proof

The first proof of Legendre's relation (1.1) is based on a differential equation satisfied by the elliptic integrals

$$K(k) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}} \quad \text{and} \quad E(k) = \int_0^1 \sqrt{\frac{1-k^2x^2}{1-x^2}} dx.$$

Among the many identities satisfied by these functions we employ an expression for their derivatives.

Proposition 2.1 *The functions $K(k)$ and $E(k)$ satisfy*

$$\begin{aligned} k(k')^2 \frac{dK}{dk} &= E - (k')^2 K \\ k \frac{dE}{dk} &= E - K, \end{aligned} \tag{2.1}$$

where $k' = \sqrt{1-k^2}$ is the conjugate modulus.

Proof. This follows directly from the definitions. \square

Proposition 2.2 *Let $K'(k) = K(k')$ and $E'(k) = E(k')$. Then the function $KE' + EK' - KK'$ is constant.*

Proof. Employ Proposition 2.1 to check that the derivative is identically 0. \square

Legendre then evaluates the constant at the modulus $k = \frac{1}{2}\sqrt{2-\sqrt{3}}$ and its complement $k' = \frac{1}{2}\sqrt{2+\sqrt{3}}$. In this paper we complete Legendre's proof by using the modulus $k = \sqrt{-1}$. This is explained in the next section.

3 Euler's direct proof

In [3] Euler developed his theory of infinite products and used it in [4] to prove the relation

$$\int_0^1 \frac{dx}{\sqrt{1-x^4}} \times \int_0^1 \frac{x^2}{\sqrt{1-x^4}} dx = \frac{\pi}{4}. \quad (3.1)$$

In this paper we generalize Euler's method and prove the following result.

Theorem 3.1 *The generalized elastic curve*

$$f_n(x) := \int_0^x \frac{t^n}{\sqrt{1-t^{2n}}} dt \quad (3.2)$$

satisfies

$$R_n \times L_n = \frac{\pi}{2n},$$

R_n is the main radius, the value $f_n(1)$, and L_n is the length of the curve from $x = 0$ to $x = 1$.

Proof. We have

$$R_n = \int_0^1 \frac{t^n}{\sqrt{1-t^{2n}}} dt \quad \text{and} \quad L_n = \int_0^1 \frac{dt}{\sqrt{1-t^{2n}}}.$$

Integrate the relation

$$d \left(t^k \sqrt{1-t^{2n}} \right) = \frac{kt^{k-1} dt - (k+n)t^{2n+k-1} dt}{\sqrt{1-t^{2n}}}$$

from 0 to 1 to produce the recursive formula

$$\int_0^1 \frac{t^{k-1}}{\sqrt{1-t^{2n}}} dt = \frac{k+n}{k} \int_0^1 \frac{t^{2n+k-1}}{\sqrt{1-t^{2n}}} dt. \quad (3.3)$$

The value $k = n + 1$ in (3.3) yields

$$R_n = \frac{2n+1}{n+1} \int_0^1 \frac{t^{3n}}{\sqrt{1-t^{2n}}} dt. \quad (3.4)$$

Then the value $k = 3n + 1$ produces

$$\int_0^1 \frac{t^{3n}}{\sqrt{1-t^{2n}}} dt = \frac{4n+1}{3n+1} \int_0^1 \frac{t^{5n}}{\sqrt{1-t^{2n}}} dt,$$

so (3.4) produces

$$R_n = \frac{2n+1}{n+1} \times \frac{4n+1}{3n+1} \int_0^1 \frac{t^{5n}}{\sqrt{1-t^{2n}}} dt.$$

Iterating (3.3) we obtain, after m steps,

$$R_n = \prod_{j=1}^m \frac{2jn+1}{(2j-1)n+1} \times \int_0^1 \frac{t^{(2m+1)n}}{\sqrt{1-t^{2n}}} dt. \quad (3.5)$$

The next step is to justify the passage to the limit in (3.5) as $m \rightarrow \infty$, with n fixed. Observe that the left hand side is *independent* of m , so it remains R_n after $m \rightarrow \infty$. The difficulty in passing to the limit is that the product in (3.5) diverges. The general term p_j satisfies

$$1 - p_j = \frac{-n}{(2j-1)n+1}$$

and the divergence of the product follows from that of the harmonic series. The divergence is cured by introducing scaling factors both in the integral and the product. The proof is omitted in Eulerian fashion.

Proposition 3.2 *The functions*

$$\frac{1}{2m+1} \int_0^1 \frac{t^{(2m+1)n}}{\sqrt{1-t^{2n}}} dt \quad \text{and} \quad (2m+1) \times \prod_{j=1}^m \frac{2jn+1}{(2j-1)n+1}$$

have non-zero limits as $m \rightarrow \infty$.

Therefore from (3.5) we obtain

$$R_n = \lim_{m \rightarrow \infty} \prod_{j=1}^{2m} (jn+1)^{(-1)^j} \times \int_0^1 \frac{t^{(2m+1)n}}{\sqrt{1-t^{2n}}} dt$$

where we have employed

$$\prod_{j=1}^m \frac{2jn+1}{(2j-1)n+1} = \prod_{j=1}^{2m} (jn+1)^{(-1)^j}$$

in order to simplify the notation. A similar argument shows that

$$\begin{aligned} L_n &= \prod_{j=1}^m \frac{(2j-1)n+1}{2(j-1)n+1} \int_0^1 \frac{t^{2mn}}{\sqrt{1-t^{2n}}} dt \\ &= \lim_{m \rightarrow \infty} \prod_{j=1}^{2m} (jn+1)^{(-1)^{j+1}} \int_0^1 \frac{t^{2mn}}{\sqrt{1-t^{2n}}} dt. \end{aligned} \quad (3.6)$$

The final step is to introduce the auxiliary quantities

$$A_n := \int_0^1 \frac{t^{n-1}}{\sqrt{1-t^{2n}}} dt \quad \text{and} \quad B_n := \int_0^1 \frac{t^{2n-1}}{\sqrt{1-t^{2n}}} dt.$$

We now show that the quotient L_n/A_n can be evaluated explicitly and that the value of A_n is elementary. This produces an expression for L_n . A similar statement holds for R_n/B_n and B_n .

Observe first that

$$A_n = \int_0^1 \frac{t^{n-1}}{\sqrt{1-t^{2n}}} dt = \frac{1}{n} \int_0^1 \frac{dx}{\sqrt{1-x^2}} = \frac{\pi}{2n} \quad (3.7)$$

and similarly $B_n = 1/n$. Now consider the recursion (3.3) for odd multiples of n to produce

$$A_n = \lim_{m \rightarrow \infty} \prod_{j=1}^{2m} (jn)^{(-1)^j} \times \int_0^1 \frac{t^{(2m+1)n-1}}{\sqrt{1-t^{2n}}} dt \quad (3.8)$$

and similarly the even multiples of n yield

$$B_n = \frac{1}{n} \lim_{m \rightarrow \infty} \prod_{j=1}^{2m+1} (jn)^{(-1)^{j+1}} \times \int_0^1 \frac{t^{2(m+1)n-1}}{\sqrt{1-t^{2n}}} dt,$$

in the exact manner as the derivation of (3.5). Therefore using (3.6) and (3.8), and passing to the limit as $m \rightarrow \infty$ so that the integrals disappear, we obtain

$$\frac{L_n}{A_n} = \prod_{j=1}^{\infty} [(jn+1)^{(-1)^{j+1}} \times (jn)^{(-1)^{j+1}}],$$

so (3.7) yields

$$L_n = \frac{\pi}{2n} \times \prod_{j=1}^{\infty} [(jn+1)^{(-1)^{j+1}} \times (jn)^{(-1)^{j+1}}].$$

Similarly, using $B_n = 1/n$,

$$R_n = \prod_{j=1}^{\infty} [(jn+1)^{(-1)^j} \times (jn)^{(-1)^j}].$$

The formula $R_n \times L_n = \pi/2n$ follows directly from here. \square

4 Conclusions

In this paper we have established that the main radius R_n of the generalized elastic curve (3.2) and the length L_n of this curve satisfy $R_n \times L_n = \pi/2n$. The case $n = 2$ corresponds to the classical Legendre's formula for elliptic integrals.

References

- [1] Abel, N.: *Recherches sur les fonctions elliptiques*. Crelle Journal **1**, 1827.
- [2] Duren, P.: *The Legendre relation for elliptic integrals* in Paul Halmos, *Celebrating 50 years of Mathematics*. Editors John H. Ewing and F. W. Gehring, 305–315. Springer-Verlag, 1991.
- [3] Euler, L.: *Animadversiones in Rectificationem Ellipsis*. Comment 154 Enestroemianus Index, opuscula varii argumenti. **2**, 1750, 121–166.
- [4] Euler, L.: *De miris proprietatibus curvae elasticae sub aequatione $y = \int \frac{xx}{\sqrt{1-x^4}} dx$ contentae*. Comment 605 Enestroemianus Index. Acta academiae scientiarum Petrop. 1782: II (1786) 34–61. Reprinted in *Opera Omnia*, ser. 1, **21**, 91–118.
- [5] Legendre, A.M.: *Exercices de calcul integral sur diverses ordres de transcendentes*. Paris. 1811.
- [6] Legendre, A.M.: *Traité des fonctions elliptiques et des integrales Euleriennes*. Paris. 1825.
- [7] McKean, H.; Moll, V.: *Elliptic Curves: Function Theory, Geometry, Arithmetic*. Cambridge University Press, 1997.
- [8] Siegel, C. L.: *Topics in Complex Function Theory I, Elliptic Functions and Uniformization*. Wiley-Interscience, 1969.
- [9] Struik, D.J. (Ed.): *A Source Book in Mathematics, 1200-1800*. Harvard University Press, Cambridge, Massachusetts, 1969.

Victor H. Moll
Department of Mathematics
Tulane University
New Orleans, LA 70118, USA
e-mail: vhm@math.tulane.edu

Pamela A. Neill
Department of Mathematics
Delgado Community College
New Orleans, LA 70119, USA
e-mail: pneill@pop3.dcc.edu

Judith L. Nowalsky
Department of Mathematics
Tulane University
New Orleans, LA 70118, USA
e-mail: judithn@math.tulane.edu

Leonardo Solanilla
Instituto de Fisica y Matematicas
Universidad Michoacana
Edificio C3, Ciudad Universitaria
Morelia CP 58040, Michoacan, Mexico