

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 56 (2001)

**Artikel:** On the Representation of Permutation as Products of Transpositions  
**Autor:** Neuenschwander, Daniel  
**DOI:** <https://doi.org/10.5169/seals-6666>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 23.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

---

## On the Representation of Permutations as Products of Transpositions

---

Daniel Neuenschwander

Daniel Neuenschwander was born in 1963 near Neuchâtel (CH). After his doctorate (under the direction of H. Carnal) at the University of Bern in 1991, he was appointed professor of mathematics at the Biel School of Engineering and Architecture of the Bern University of Applied Sciences (the former HTL). After having got a part-time professorship for actuarial sciences at the University of Lausanne, he completed his habilitation in 1996. He teaches mathematics of insurance and finance also at the University of Bern and holds a position at the Federal section of cryptology at the Department of Defense. He has taught at the University of Dortmund and has been invited professor at the University of Nancy I. His main mathematical working-area is probability theory (especially on algebraic structures) and its applications in insurance, finance, and cryptology. Non-mathematical interests include music (organ, harpsichord, piano) and travelling.

The subject of this note is the representation of permutations of finite sets as products of a minimal number of simple transpositions.

Consider the following theorem:

**Theorem 1**

- a) *If a permutation  $\varphi$  which is a product of  $n$  transpositions cannot be written as a product of fewer than  $n$  transpositions, then for any transposition occurring in this product both elements belong to the same cycle of  $\varphi$ .*

Einer Menge bestehend aus  $n$  Elementen ist in natürlicher Weise die Menge  $S_n$  der Permutationen der Elemente dieser Menge zugeordnet. Es ist bekannt, dass  $S_n$  eine Gruppe mit  $n!$  Elementen ist. Bereits  $S_3$  liefert ein Beispiel einer nicht-kommutativen Gruppe, und die Untergruppe  $A_n \subset S_n$  vom Index 2, die sogenannte alternierende Gruppe, ist für  $n > 4$  einfach, was zur Nicht-Lösbarkeit algebraischer Gleichungen vom Grad grösser als vier durch Radikale führt. Andererseits ist es einfach zu beweisen, dass jede Permutation  $\pi \in S_n$  als Produkt von Zykeln darstellbar ist. Spezielle Zykeln sind die Transpositionen, d.h. die Vertauschungen zweier Elemente; es zeigt sich, dass  $\pi$  insbesondere als Produkt von Transpositionen darstellbar ist. Im vorliegenden Beitrag wird nun auf elementare Weise gezeigt, welches die minimale Anzahl von Transpositionen ist, die man zur Darstellung einer Permutation benötigt, und auf wieviele Arten eine solche Darstellung erfolgen kann. *jk*

**b)** A permutation  $\varphi$  which is a cycle of length  $n + 1$  cannot be written as a product of fewer than  $n$  transpositions.

Part b) of Theorem 1 is well-known, see e.g. Schwenk (1984), Lossers (1986), and the literature cited there. We show how assertions a) and b) can be proved together by induction on  $n \in \mathbb{N}_0$ . We will use the word "cycle" in the sense that a cycle can also be one-elemented. However, transpositions will always be genuine, i.e. two-elemented.

*Proof of Theorem 1.* For a cycle  $\varphi$ , denote by  $|\varphi|$  its length. For  $n = 0$ , a) and b) are trivial. Assume them to be true for all  $0 \leq k \leq n$  and consider  $\varphi = \varphi'(a, b)$ , where  $\varphi'$  is a product of  $n$  transpositions and  $n + 1$  is the minimum number of transpositions which is necessary to represent  $\varphi$ . Let  $c_i$  ( $1 \leq i \leq m$ ) denote the disjoint cycles of  $\varphi'$ . If both  $a, b$  belonged to the same  $c_{i_0}$ , then the permutation  $c_{i_0}(a, b)$  as a product of not fewer than  $|c_{i_0}|$  transpositions (induction hypothesis b)) would consist of two disjoint cycles of  $|c_{i_0}|$  elements together and could therefore be represented by  $|c_{i_0}| - 2$  transpositions, which is a contradiction. So  $a, b$  belong to two different  $c_i$ 's, thus any of the cycles  $(a, b), c_i$  ( $1 \leq i \leq m$ ) is part of one of the disjoint cycles  $d_i$  of  $\varphi$ , and therefore for any transposition of  $\varphi$  in a decomposition of  $\varphi$  into a minimal number of transpositions both interchanged elements belong to the same  $d_i$ , which proves a). Now assume in addition that  $\varphi$  is a cycle itself, i.e. w.l.o.g.  $\varphi = (1, 2, \dots, L)$ ,  $a, b \in \mathbb{N}$ ,  $a < b$ . We show that  $L = n + 2$ . We have  $\varphi' = \varphi''\varphi'''$ , where

$$\varphi'' = (1, 2, \dots, a, \varphi(b), \varphi^2(b), \dots, L)$$

and

$$\varphi''' = (\varphi(a), \varphi^2(a), \dots, \varphi^{-1}(b), b).$$

We have  $|\varphi''| = L - (b - a)$  and  $|\varphi'''| = b - a$ , hence (by counting transpositions in a minimal transposition decomposition and using induction hypotheses a) and b))

$$n + 1 = |\varphi''| + |\varphi'''| - 1 = L - (b - a) + b - a - 1,$$

hence  $L = n + 2$ .  $\square$

In this context, a natural question is also in how many ways a cycle of length  $n$  can be represented as a product of  $n - 1$  transpositions. Let  $N_n$  be this number.

**Theorem 2**  $N_n = n^{n-2}$  ( $n \in \mathbb{N}$ ).

This property has been proved by Lossers (1986), using the formula for the number of point-labeled trees of  $n$  points. See also the literature cited in Lossers (1986) for other references to this theorem. Here, let us present a self-contained proof based on Theorem 1a) and Abel's identity:

$$\sum_{k=0}^m \binom{m}{k} (x+k)^{k-1} (y+m-k)^{m-k} = x^{-1} (x+y+m)^m. \quad (1)$$

*Proof of Theorem 2.* For  $n = 1$  the assertion is trivial. We assume it to be true for  $k \leq n$  and prove it for  $n + 1$ . The cycle

$$\psi = (1, 2, \dots, n+1) = (a_1, b_1)(a_2, b_2) \cdots (a_n, b_n)$$

$(a_k < b_k \ (1 \leq k \leq n))$  can be represented as follows:

$$\psi = \psi'(a_n, b_n)$$

with

$$\begin{aligned}\psi' &= \psi''\psi''', \\ \psi'' &= (1, 2, \dots, a_n, \psi(b_n), \psi^2(b_n), \dots, n+1), \\ \psi''' &= (\psi(a_n), \psi^2(a_n), \dots, \psi^{-1}(b_n), b_n).\end{aligned}$$

Writing  $k := b_n - a_n$ , the triple  $(\psi'', \psi''', (a_n, b_n))$  is uniquely determined by the pair  $(k, a_n)$ . For a given  $k$ , there exist  $n+1-k$  possibilities to choose  $a_n$ . As (in consideration of Theorem 1a)) for all  $(a_k, b_k)$  ( $1 \leq k \leq n-1$ ), the elements  $a_k, b_k$  belong to the same cycle of  $\psi'$ , there are (for fixed  $k, a_n$ )  $\binom{n-1}{k-1} N_k N_{n+1-k}$  possibilities for  $\psi'$ , thus by the induction hypothesis and (1) (with  $x := y := 1, m := n-1$ ) we calculate

$$\begin{aligned}N_{n+1} &= \sum_{k=1}^n (n+1-k) \binom{n-1}{k-1} N_k N_{n+1-k} \\ &= \sum_{k=1}^n (n+1-k) \binom{n-1}{k-1} k^{k-2} (n+1-k)^{n+1-k-2} \\ &= \sum_{k=0}^{n-1} \binom{n-1}{k} (k+1)^{k-1} (n-k)^{n-1-k} \\ &= (n+1)^{n-1}.\end{aligned}\quad \square$$

## References

- [1] Schwenk, A.J., Problem E 3058. *Amer. Math. Monthly* (1984), 516.
- [2] Lossers, O.P., Solution to Problem E 3058. *Amer. Math. Monthly* (1986), 820–821.

Daniel Neuenschwander  
 Université de Lausanne  
 Institut des Sciences Actuarielles  
 CH-1015 Lausanne  
 Universität Bern  
 Institut für mathematische Statistik und Versicherungslehre  
 Sidlerstrasse 5  
 CH-3012 Bern  
 Sektion Kryptologie  
 Departement VBS  
 CH-3003 Bern  
 e-mail: neuens@bluewindow.ch