

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 57 (2002)

Artikel: Some constrained optimization problems in elementary statistics
Autor: Stadje, Wolfgang
DOI: <https://doi.org/10.5169/seals-7602>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Some constrained optimization problems in elementary statistics

Wolfgang Stadje

Wolfgang Stadje promovierte und habilitierte in Mathematik an der Universität Göttingen. Er ist Professor an der Universität Osnabrück und arbeitet hauptsächlich über stochastische Prozesse und ihre Anwendungen, z.B. auf Warteschlangen- und Lagerhaltungsmodelle, sequentielle Analysis sowie über analytische und kombinatorische Probleme in der Wahrscheinlichkeitstheorie.

1 Introduction

Let $x_1, \dots, x_n \in \mathbb{R}$ be n measurements of some quantity. The following elementary functions of x_1, \dots, x_n are of particular statistical importance:

- the arithmetic mean $\bar{x} = n^{-1} \sum_{i=1}^n x_i$;
- the variance $s^2 = n^{-1} \sum_{i=1}^n (x_i - \bar{x})^2$;
- the extreme values $M = \max_{1 \leq i \leq n} x_i$ and $m = \min_{1 \leq i \leq n} x_i$ and the range $R = M - m$;
- in the case of positive x_i 's, the geometric mean $\bar{x}_g = (\prod_{i=1}^n x_i)^{1/n}$ and the harmonic mean $\bar{x}_h = n / \sum_{i=1}^n 1/x_i$.

The arithmetic mean is certainly the most widely used (and oldest) method to combine discordant measurements in order to summarize the data in a single value. The geometric and the harmonic mean are measures of location used in special circumstances, for example to determine an average of n successive price increases, or an average price if the same amount of some goods is purchased n times at different prices (see for

Zur Bündelung der in n Meßwerten einer Größe enthaltenen Information sind das arithmetische Mittel, die Varianz, Maximum und Minimum sowie die Spannweite die gebräuchlichsten statistischen Merkmale. Die vorliegende Arbeit geht der Frage nach, welche Werte diese statistischen Funktionen jeweils maximal annehmen können, wenn Bedingungen an die anderen gestellt werden. Mehrere solche Optimierungsaufgaben mit Nebenbedingungen haben ästhetisch ansprechende Lösungen, die mit Hilfe verschiedener elementarer Techniken hergeleitet werden. Neben dem arithmetischen werden außerdem das geometrische und das harmonische Mittel betrachtet.

example [1]). The three means satisfy the inequalities $\bar{x} \geq \bar{x}_g \geq \bar{x}_h$. The variance is the classical measure of dispersion around the center of location, while the range gives the full amount of variability without indicating how concentrated ‘most of’ the measurements are. \bar{x} and s^2 are also important quantities in mechanics: if a total mass 1 is distributed on the real line in portions of size $1/n$ at the points x_1, \dots, x_n , then \bar{x} and s^2 are the center of gravity and the moment of inertia around this center, respectively.

It seems interesting to find the extremal values of these functions under constraints on one or several of the others, especially if the solutions turn out to have an appealing analytic form. This is indeed the case, and in this note we will solve the following six of these constrained optimization problems:

- (P1)** Maximize s^2 subject to $\bar{x} = \mu$ and $0 \leq x_i \leq c$, $i = 1, \dots, n$ (with given $\mu \in \mathbb{R}$ and $c > 0$).
- (P2)** Maximize s^2 subject to $\bar{x} = \mu$, $R = r$ ($\mu \in \mathbb{R}$, $r > 0$ given).
- (P3)** Maximize $M = \max_{1 \leq i \leq n} x_i$ subject to $\bar{x} = \mu$ and $s^2 = \sigma^2$ (with given $\mu \in \mathbb{R}$ and $\sigma^2 > 0$).
- (P4)** Maximize the range R subject to $\bar{x} = \mu$, $s^2 = \sigma^2$.
- (P5)** Maximize \bar{x}_g/\bar{x} subject to $x_1, \dots, x_n > 0$, $1 - \delta_0 \leq x_i/\bar{x} \leq 1 + \delta_1$ for given $\delta_0 \in [0, 1)$ and $\delta_1 \in [0, 1)$, not both equal to zero.
- (P6)** Minimize \bar{x}_h subject to $\bar{x} = \mu$, $s^2 = \sigma^2$ and $x_i \geq c$, $i = 1, \dots, n$ (with given $\sigma^2 > 0$, $c > 0$ and $\mu \in (c, \infty)$).

Solutions. The extremal values in the above problems are given by the following formulas:

- (P1)** $s_{\max}^2 = n^{-1}(j(c - \mu)^2 + (n - j - 1)\mu^2 + ((n - 1)\mu - jc)^2)$, where the nonnegative integer j is defined by $j < n\mu/c \leq j + 1$.

$$\text{(P2)} \quad s_{\max}^2 = \begin{cases} \frac{n^2 - 1}{4n^2} r^2 & \text{if } n \text{ is odd,} \\ \frac{1}{4n} r^2 & \text{if } n \text{ is even.} \end{cases}$$

$$\text{(P3)} \quad M_{\max} = \mu + (n - 1)^{\frac{1}{2}} \sigma^2.$$

$$\text{(P4)} \quad R_{\max} = (2n)^{\frac{1}{2}} \sigma^2.$$

$$\text{(P5)} \quad (\bar{x}_g/\bar{x})_{\max} = ((1 - \delta_0)^k (1 + \delta_1)^{n-k-1} (1 + k(\delta_0 + \delta_1)) - (n - 1)\delta_1)^{-1/n}, \text{ where } k = \lceil n\delta_1/(\delta_0 + \delta_1) \rceil.$$

- (P6)** For this problem we can only prove that

$$(\bar{x}_h)_{\min} \geq \frac{c(\sigma^2 + \mu^2 - c\mu)}{\sigma^2 - \mu^2 + c\mu}.$$

The exact value of the minimum is unknown.

Of course, $[a]$ is defined to be the largest integer m satisfying $m \leq a$.

We will prove these results and also give the values of x_1, \dots, x_n for which the extrema are attained. Many similar problems whose solutions seem to be unknown can be easily formulated. An example is (P6); the corresponding maximization problem is also open.

A basic reference text on inequalities of the type considered here is [3]. An interesting paper in the spirit of this note is [2], where the following is proved: For any $m \geq 3$ the maximal value α_m of $n^{-1} \sum_{i=1}^n x_i^m$ subject to $\bar{x} = 0$ and $s^2 = 1$ is attained at

$$(x_1, \dots, x_n) = ((n-1)^{1/2}, -(n-1)^{-1/2}, \dots, -(n-1)^{-1/2}),$$

bounded by $((n-1)^m + (-1)^m)/n(n-1)^{(m/2)-1}$ and satisfies $\alpha_{2m} \geq \alpha_{m+1}^2 + \alpha_m^2$.

2 Derivations

Theorem 1 *In problem (P1), choose j such that $j < n\mu/c \leq j+1$. Then the maximum is attained at $x^0 = (x_1^0, \dots, x_n^0)$, where*

$$x_1^0 = \dots = x_j^0 = c, \quad x_{j+1}^0 = n\mu - jc, \quad x_{j+2}^0 = \dots = x_n^0 = 0, \quad (2.1)$$

and is equal to

$$s_{\max}^2 = n^{-1} \left[j(c - \mu)^2 + (n - j - 1)\mu^2 + ((n - 1)\mu - jc)^2 \right]. \quad (2.2)$$

Proof. Since $ns^2 = \sum_{i=1}^n x_i^2 - n\mu^2$, we have to maximize $\sum_{i=1}^n x_i^2$ subject to $\bar{x} = \mu$ and, without restriction of generality, $c \geq x_1 \geq \dots \geq x_n \geq 0$. Suppose that the maximum under these constraints is attained at some point (u_1, \dots, u_n) satisfying $u_k = c$, $k \leq i_0$, for some $i_0 \in \{0, 1, \dots, j-1\}$ and $u_{i_0+1} < c$. Then

$$u_1 + \dots + u_{i_0} + u_{i_0+1} < i_0 c + c \leq jc < n\mu,$$

so that $u_{i_0+2} > 0$. But

$$\begin{aligned} (u_{i_0+1} + \varepsilon)^2 + (u_{i_0+2} - \varepsilon)^2 &= u_{i_0+1}^2 + u_{i_0+2}^2 + 2\varepsilon^2 + 2\varepsilon(u_{i_0+1} - u_{i_0+2}) \\ &> u_{i_0+1}^2 + u_{i_0+2}^2 \end{aligned}$$

for all $\varepsilon > 0$. In particular, taking $\varepsilon \in (0, \min[c - u_{i_0+1}, u_{i_0+2}])$ yields the contradiction that the maximum is not attained at (u_1, \dots, u_n) .

Thus, if $x_1^0 \geq \dots \geq x_n^0$ is a maximizing point, we must have $x_1^0 = \dots = x_j^0 = c$.

If $j+2 \leq n$ and $x_{j+2}^0 > 0$, reasoning as above shows that x^0 could not be a maximum. Hence, $x_i^0 = 0$ for $i > j+1$ and $x_{j+1}^0 = n\mu - x_1^0 - \dots - x_j^0 = n\mu - jc$. This proves (2.1), and (2.2) is now immediate. \square

Theorem 2 In (P2) the maximal value is attained at

$$x^0 = \begin{cases} \left(\mu - \frac{r}{2}, \dots, \mu - \frac{r}{2}, \mu + \frac{r}{2}, \dots, \mu + \frac{r}{2}\right) & \text{if } n \text{ is even,} \\ \left(\mu - \frac{n+1}{2n}r, \dots, \mu - \frac{n+1}{2n}r, \mu + \frac{n-1}{2n}r, \dots, \mu + \frac{n-1}{2n}r\right) & \text{if } n \text{ is odd,} \end{cases} \tag{2.3}$$

and is equal to

$$s_{\max}^2 = \begin{cases} \frac{1}{4n}r^2 & \text{if } n \text{ is even,} \\ \frac{n^2-1}{4n^2}r^2 & \text{if } n \text{ is odd.} \end{cases} \tag{2.4}$$

Proof. We may assume that $\mu = 0$. Let $x = (x_1, \dots, x_n)$ be a point at which the maximum is attained. We show that up to a permutation x must be of the form $(\alpha, \dots, \alpha, \beta, \dots, \beta)$ with $\alpha < 0 < \beta$. If this does not hold, then there is a nonnegative component smaller than $\max_{1 \leq i \leq n} x_i$ or a nonpositive component greater than $\min_{1 \leq i \leq n} x_i$. Thus let us suppose that one of the nonnegative x_i , say x_{i_0} , is smaller than $\max_{1 \leq i \leq n} x_i$.

Then all negative x_i must be equal to $\min_{1 \leq i \leq n} x_i$, because if $0 > x_{j_0} > \min_{1 \leq i \leq n} x_i$ for some x_{j_0} , then the point x' defined by $x'_{i_0} = x_{i_0} + \varepsilon$, $x'_{j_0} = x_{j_0} - \varepsilon$, $x'_i = x_i$ for $i \neq i_0, j_0$ has the arithmetic mean $\bar{x}' = 0$ and has, for small $\varepsilon > 0$, the same range as x , but

$$\sum_{i=1}^n (x'_i)^2 = 2\varepsilon^2 + 2(x_{i_0} - x_{j_0})\varepsilon + \sum_{i=1}^n x_i^2 > \sum_{i=1}^n x_i^2 \text{ for all } \varepsilon > 0.$$

Furthermore, all nonnegative x_i except x_{i_0} are equal to $\max_{1 \leq i \leq n} x_i$, because if $0 \leq x_{i_1} \leq x_{i_1} < \max_{1 \leq i \leq n} x_i$ for some $i_1 \neq i_0$, the point x'' defined by $x''_{i_0} = x_{i_0} - \varepsilon$, $x''_{i_1} = x_{i_1} + \varepsilon$ and $x''_i = x_i$ for $i \neq i_0, i_1$ has a larger sum of squares than x but the same range for small $\varepsilon > 0$, and mean $\bar{x}'' = 0$. It follows that up to a permutation x is of the form

$$(\alpha, \dots, \alpha, \gamma, \beta, \dots, \beta), \quad k \text{ } \alpha\text{'s and } n - k - 1 \text{ } \beta\text{'s,}$$

where $\alpha < 0 \leq \gamma \leq \beta$ and $k \in \{1, \dots, n - 2\}$. We may assume that $\gamma = \beta$, since if $\gamma < \beta$ the point

$$(\alpha - (n - 1)^{-1}\varepsilon, \dots, \alpha - (n - 1)^{-1}\varepsilon, \gamma + \varepsilon, \beta - (n - 1)^{-1}\varepsilon, \dots, \beta - (n - 1)^{-1}\varepsilon), \quad \varepsilon > 0$$

gives the same mean value and the same range as x , as long as $0 < \gamma + \varepsilon < \beta$, while its sum of squares is

$$\begin{aligned} &k\alpha^2 + \gamma^2 + (n - k - 1)\beta^2 + \varepsilon^2 \left(\frac{k}{(n - 1)^2} + 1 + \frac{n - k - 1}{(n - 1)^2} \right) \\ &+ 2\varepsilon \left(\gamma - \frac{\alpha k}{n - 1} - \frac{\beta(n - k - 1)}{n - 1} \right) > k\alpha^2 + \gamma^2 + (n - k - 1)\beta^2; \end{aligned}$$

the inequality follows from

$$\gamma - \frac{\alpha k}{n-1} - \frac{\beta(n-k-1)}{n-1} = \frac{n}{n-1} \gamma \geq 0$$

(note that $k\alpha + \gamma + (n-k-1)\beta = 0$).

Thus the maximum is attained at some point $(\alpha, \dots, \alpha, \alpha+r, \dots, \alpha+r)$ and we have to find an $\alpha < 0$ and a $k \in \{1, \dots, n-1\}$ such that

$$k\alpha^2 + (n-k)(\alpha+r)^2$$

is maximized subject to $k\alpha + (n-k)(\alpha+r) = 0$. From this constraint we obtain

$$\alpha = -\frac{n-k}{n}r, \quad \alpha+r = \frac{k}{n}r,$$

so that

$$\begin{aligned} k\alpha^2 + (n-k)(\alpha+r)^2 &= k \left(\frac{n-k}{n} \right)^2 r^2 + (n-k) \left(\frac{k}{n} \right)^2 r^2 \\ &= k(n-k) \frac{r^2}{n}. \end{aligned}$$

If n is even (odd), the maximum is attained for $k = n/2$ ($k = (n-1)/2$). A short calculation now yields the expression for s_{\max}^2 . The theorem is proved. \square

Theorem 3 *In problem (P3) the maximal value of M is equal to*

$$M_{\max} = (n-1)^{1/2} \sigma,$$

and is attained at x^0 , where

$$x_1^0 = \mu + (n-1)^{1/2} \sigma, \quad x_2^0 = \dots = x_n^0 = \mu - (n-1)^{-1/2} \sigma.$$

Proof. By symmetry, we can equivalently maximize x_1 subject to $\bar{x} = \mu$, $s^2 = \sigma^2$. We form the Lagrange function

$$L(x, \lambda_1, \lambda_2) = x_1 + \lambda_1(\bar{x} - \mu) + \lambda_2(s^2 - \sigma^2).$$

Setting its partial derivatives equal to zero, we obtain

$$\begin{aligned} 0 &= \frac{\partial L}{\partial x_1} = 1 + n^{-1} \lambda_1 + 2n^{-1} \lambda_2 (x_1 - \mu), \\ 0 &= \frac{\partial L}{\partial x_2} = n^{-1} \lambda_1 + 2n^{-1} \lambda_2 (x_2 - \mu), \\ &\vdots \\ 0 &= \frac{\partial L}{\partial x_n} = n^{-1} \lambda_1 + 2n^{-1} \lambda_2 (x_n - \mu). \end{aligned}$$

Summing these equations yields

$$0 = 1 + \lambda_1 + 2n^{-1}\lambda_2 \left(\sum_{i=1}^n x_i - n\mu \right) = 1 + \lambda_1,$$

i.e., $\lambda_1 = -1$, so that

$$x_1 = \mu + \frac{1-n}{2\lambda_2}, \quad x_2 = \dots = x_n = \mu + \frac{1}{2\lambda_2}.$$

By the constraint on the variance,

$$n\sigma^2 = \sum_{i=1}^n (x_i - \mu)^2 = \frac{(1-n)^2}{4\lambda_2^2} + \frac{n-1}{4\lambda_2^2}.$$

Therefore,

$$\lambda_2 = \pm(2\sigma)^{-1}(n-1)^{1/2}.$$

We obtain the two points x^0 and x^1 , where x^1 is defined by

$$x_1^1 = \mu - (n-1)^{1/2}\sigma, \quad x_2^1 = \dots = x_n^1 = \mu + (n-1)^{-1/2}\sigma.$$

The Jacobian matrix of the mapping $(\bar{x} - \mu, s^2 - \sigma^2)$ from \mathbb{R}^n to \mathbb{R}^2 has rank 2 at x^0 and at x^1 ; hence $M_{\max} = x_1^0 = \mu + (n-1)^{1/2}\sigma$. Note that the minimum possible value of x_1 is $\mu - (n-1)^{1/2}\sigma$. \square

Theorem 4 *In problem (P4) the maximum range is equal to*

$$R_{\max} = (2n)^{1/2}\sigma,$$

and is attained at the point x^0 given by

$$x_1^0 = \mu + (n/2)^{1/2}\sigma, \quad x_2^0 = \mu - (n/2)^{1/2}\sigma, \quad x_3^0 = \dots = x_n^0 = \mu.$$

Proof. Change the variables to $u_i = x_i - \mu$, $i = 1, \dots, n$. Then we have to maximize $u_1 - u_2$ subject to

$$\bar{u} = 0, \quad \sum_{i=1}^n u_i^2 = n\sigma^2.$$

As in the proof of Theorem 2, we consider the corresponding Lagrange function

$$L(u, \lambda_1, \lambda_2) = u_1 - u_2 + \lambda_1 \sum_{i=1}^n u_i + \lambda_2 \left(\sum_{i=1}^n u_i^2 - n\sigma^2 \right).$$

Set its derivatives equal to zero:

$$\begin{aligned} 0 &= \frac{\partial L}{\partial u_1} = 1 + \lambda_1 + 2\lambda_2 u_1, \\ 0 &= \frac{\partial L}{\partial u_2} = -1 + \lambda_1 + 2\lambda_2 u_2, \\ 0 &= \frac{\partial L}{\partial u_3} = \lambda_1 + 2\lambda_2 u_3, \\ &\vdots \\ 0 &= \frac{\partial L}{\partial u_n} = \lambda_1 + 2\lambda_2 u_n. \end{aligned}$$

Adding these equations, we find that $\lambda_1 = 0$,

$$u_1 = -(2\lambda_2)^{-1}, \quad u_2 = (2\lambda_2)^{-1}, \quad u_3 = \dots = u_n = 0$$

and

$$n\sigma^2 = (2\lambda_2)^{-1} + (2\lambda_2)^{-2}, \quad \text{i.e., } \lambda_2 = \pm(2n)^{-1/2}\sigma^{-1}.$$

It follows that the maximum of $u_1 - u_2$ is attained at

$$u^0 = ((n/2)^{1/2}\sigma, -(n/2)^{1/2}\sigma, 0, \dots, 0)$$

(and the minimum at $-u^0$). □

For our next result we need the following

Lemma *Let $-\delta_0 \leq a \leq n\delta_1$. The minimum of the function*

$$F(y_1, \dots, y_n) = \prod_{i=1}^n (1 + y_i),$$

subject to the restrictions $y_1, \dots, y_n \in [-\delta_0, \delta_1]$, $\sum_{i=1}^n y_i = a$, is only attained at points that have at least $n - 1$ components in $\{-\delta_0, \delta_1\}$.

Proof. If $n = 2$, we have to minimize $F(y_1, y_2) = ay_1 - y_1^2$ over the interval $\max[-\delta_0, a - \delta_1] \leq y_1 \leq \min[\delta_1, a + \delta_0]$. The minimum is attained at $y_1 = \delta_1, y_2 = a - \delta_1$ if $\delta_1 \leq a + \delta_0$ and at $y_1 = a + \delta_0, y_2 = -\delta_0$ if $\delta_1 > a + \delta_0$. The only other way to attain the minimum is to permute the components.

Now we proceed by induction on n . Suppose the assertion is true for $n - 1$ for some $n \geq 3$. The function $F(y_1, \dots, y_n)$ has no minimum on the set $A = \{y \in \mathbb{R}^n \mid \sum_i y_i = a, \min_i y_i > -1\}$. Indeed, if $y \in A$, the point $y' \in \mathbb{R}^n$ with components $y'_1 = y_1 - \eta$, $y'_2 = y_2 + \eta$ and $y'_i = y_i$ for $i > 2$ is in A for small $|\eta|$, and $F(y') = F(y) + [\eta(y_1 - y_2) - \eta^2] \prod_{i=3}^n (1 + y_i)$. By suitable choice of η (positive or negative and close enough to zero) we obtain $F(y') < F(y)$. It follows that any absolute minimum (u_1, \dots, u_n) of $F(y_1, \dots, y_n)$ subject to $y_1, \dots, y_n \in [-\delta_0, \delta_1]$, $\sum_i y_i = a$ must lie on the boundary, i.e., must have one component in $\{-\delta_0, \delta_1\}$. Assume that $u_n = \delta_1$. Then (u_1, \dots, u_{n-1}) is an absolute minimum of $\prod_{i=1}^{n-1} (1 + y_i)$, subject to $y_i \in [-\delta_0, \delta_1]$, $\sum_{i=1}^{n-1} y_i = a - \delta_1$. Since there are $u_1, \dots, u_{n-1} \in [-\delta_0, \delta_1]$ satisfying $\sum_{i=1}^{n-1} y_i = a - \delta_1$, it is clear that $-(n-1)\delta_0 \leq a - \delta_1 \leq (n-1)\delta_0$. Hence we can apply the induction hypothesis and conclude that among u_1, \dots, u_{n-1} at least $n - 2$ are in $\{-\delta_0, \delta_1\}$. □

Theorem 5 In problem (P5) the maximal value of \bar{x}_g/\bar{x} under the given restrictions is equal to

$$\left((1 - \delta_0)^k (1 + \delta_1)^{n-k-1} (1 + k(\delta_0 + \delta_1) - (n - 1)\delta_1) \right)^{-1/n},$$

where $k = \lceil n\delta_1/(\delta_0 + \delta_1) \rceil$, and is attained at

$$x_i - \bar{x} = \begin{cases} -\bar{x}\delta_0 & i = 1, \dots, k, \\ \bar{x}\delta_1 & i = k + 1, \dots, n - 1, \\ \bar{x}((\delta_0 + \delta_1)k - (n - 1)\delta_1) & i = n. \end{cases}$$

Proof. Let $y_i = x_i - \bar{x}$, $i = 1, \dots, n$. We may assume that $\bar{x} = 1$ and then have to minimize

$$F(y_1, \dots, y_n) = \prod_{i=1}^n (1 + y_i),$$

subject to

$$y_1, \dots, y_n \in [-\delta_0, \delta_1], \quad \sum_{i=1}^n y_i = 0.$$

Let the minimum be attained at (u_1, \dots, u_n) . By the lemma, at least $n - 1$ of the u_i 's are in $\{-\delta_0, \delta_1\}$. Suppose (without loss in generality) that $-\delta_0$ occurs k times, δ_1 occurs ℓ times and that the $-\delta_0$'s are left of the δ_1 's: $u_1 = u_2 = \dots = u_k = -\delta_0$ and $u_{k+1} = \dots = u_{k+\ell} = \delta_1$, where $k + \ell \in \{n - 1, n\}$.

Case 1. Let $k + \ell = n - 1$. Then it follows that

$$k\delta_0 - \ell\delta_1 = -\sum_{i=1}^{n-1} y_i = y_n \in [-\delta_0, \delta_1]. \tag{2.5}$$

But (2.5) implies that

$$k \leq n\delta_1/(\delta_0 + \delta_1) \leq k + 1. \tag{2.6}$$

If $n\delta_1/(\delta_0 + \delta_1)$ is not an integer, then, by (2.6), $k = \lceil n\delta_1/(\delta_0 + \delta_1) \rceil$. If $n\delta_1/(\delta_0 + \delta_1)$ is an integer, then $n\delta_1/(\delta_0 + \delta_1) \in \{k, k + 1\}$. Assume that $n\delta_1/(\delta_0 + \delta_1) = k + 1$. Then $\ell = n - 1 - k = n\delta_0/(\delta_0 + \delta_1)$ and, by (2.5),

$$u_n = k\delta_0 - \ell\delta_1 = \left(\frac{n\delta_1}{\delta_0 + \delta_1} - 1 \right) \delta_0 - \frac{n\delta_0}{\delta_0 + \delta_1} \delta_1 = -\delta_0,$$

which contradicts the definition of k . Thus $n\delta_1/(\delta_0 + \delta_1) = k$.

Case 2. Let $k + \ell = n$. Then

$$0 = k\delta_0 - \ell\delta_1 = k\delta_0 - (n - k)\delta_1,$$

so that $k = n\delta_1/(\delta_0 + \delta_1)$.

Hence $k = \lceil n\delta_1/(\delta_0 + \delta_1) \rceil$ in both cases, and there are at least $n - k - 1$ components equal to δ_1 . The n -th component is given by $\sum_i u_i = 0$, i.e.,

$$u_n = \lceil n\delta_1/(\delta_0 + \delta_1) \rceil (\delta_0 + \delta_1) - (n - 1)\delta_1.$$

The rest of the proof is now straightforward. □

Theorem 6 *The minimum of \bar{x}_h in problem (P6) satisfies*

$$(\bar{x}_h)_{\min} \geq \frac{c(\sigma^2 + \mu^2 - c\mu)}{\sigma^2 - \mu^2 + c\mu}. \quad (2.7)$$

Proof. The problem is equivalent to finding an upper bound for $n^{-1} \sum_{i=1}^n 1/x_i$ under the given constraints. Let $H(x) = 1/x$, $x > 0$. Suppose that some parabola $P(x) = a_0x^2 + a_1x + a_2$ satisfies

$$H(x) \leq P(x) \quad \text{for all } x \in [c, \infty). \quad (2.8)$$

Then

$$n^{-1} \sum_{i=1}^n \frac{1}{x_i} = n^{-1} \sum_{i=1}^n H(x_i) \leq n^{-1} \sum_{i=1}^n P(x_i) \quad (2.9)$$

yielding an upper bound containing μ , σ^2 and the coefficients of P . For any $x_0 \in [c, \infty)$ and any $a > 0$, the special parabola

$$P_{x_0,a}(x) = a(x - x_0)^2 - x_0^{-2}(x - x_0) + x_0^{-1}$$

is tangential to the hyperbola H at x_0 , i.e., has a double point of intersection with H at x_0 . It is easy to check that H and $P_{x_0,a}$ only intersect at x_0 and at $b = (ax_0^2)^{-1}$. Therefore, if $b \leq c$, inequality (2.8) holds. Let $a = (cx_0^2)^{-1}$. Then $b = c$, so that we can use $P_{x_0,a}$ in (2.9) and obtain

$$\begin{aligned} n^{-1} \sum_{i=1}^n \frac{1}{x_i} &\leq n^{-1} \sum_{i=1}^n \left[(cx_0^2)^{-1}(x_i - x_0)^2 - x_0^{-2}(x_i - x_0) + x_0^{-1} \right] \\ &= (cx_0^2)^{-1} \left(n^{-1} \sum_{i=1}^n x_i^2 + x_0^2 - 2x_0\mu \right) - x_0^{-2}\mu + 2x_0^{-1} \\ &= (cx_0^2)^{-1} (\sigma^2 + \mu^2 + x_0^2 - 2x_0\mu) - x_0^{-2}\mu + 2x_0^{-1} \\ &= \frac{\sigma^2 + \mu^2 - c\mu}{cx_0^2} - \frac{2}{x_0} \left(\frac{\mu}{c} - 1 \right) + \frac{1}{c} \end{aligned} \quad (2.10)$$

for all $x_0 \geq c$. As a function of $x_0 \in [c, \infty)$, the right-hand side of (2.9) is minimal for

$$x_0 = (\sigma^2 + \mu^2 - c\mu)/(\mu - c); \quad (2.11)$$

note that the value of x_0 given in (2.11) is well-defined and in (c, ∞) , because $\mu > c$ and $\sigma^2 + \mu^2 - c\mu > (\mu - c)^2 + c\mu - c^2 \geq c(\mu - c)$. Inserting this x_0 in (2.10), the right-hand side becomes

$$\frac{1}{c} \left(1 - \frac{(\mu - c)^2}{\sigma^2 + \mu^2 - c\mu} \right),$$

which yields the lower bound in (2.7). The theorem is proved. \square

References

- [1] Bosch, K.: *Statistik-Taschenbuch*. Oldenbourg, München und Wien 1992.
- [2] Lakshmanamurti, M.: On the upper bound of $\sum_{i=1}^n x_i^m$ subject to the conditions $\sum_{i=1}^n x_i = 0$ and $\sum_{i=1}^n x_i^2 = n$. *Math. Student* 18 (1950), 111–116.
- [3] Mitrinović, D.S.: *Analytic Inequalities*. Springer, Berlin etc. 1970.

Wolfgang Stadje
Universität Osnabrück
Fachbereich Mathematik/Informatik
D-49069 Osnabrück, Deutschland
e-mail: wolfgang@mathematik.uni-osnabrueck.de