

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 59 (2004)

**Artikel:** Analogies are interesting!  
**Autor:** Sastry, K.R.S.  
**DOI:** <https://doi.org/10.5169/seals-9308>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 14.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

---

---

## Analogy are interesting!

---

---



K.R.S. Sastry

K.R.S. Sastry obtained his Bachelor's and Master's degrees in mathematics from the University of Mysore in India. He taught mathematics, initially in the State of Karnataka in India, later on in Ethiopia. After his retirement in 1994 he returned to the city of Bangalore in India, where he continues to work on mathematical problems.

Let an equilateral triangle  $ABC$  be inscribed in a circle. Let  $P$  be any point on the minor  $\widehat{BC}$  (arc  $BC$ ) as in Fig. 1. Then it is well-known that  $AP = BP + PC$ . (This can be easily proved using Ptolemy's theorem, trigonometry or by other means.) Is not this property pretty? Obviously, we do not expect a non equilateral triangle to exhibit such a property. However, in a given triangle  $ABC$  it seems reasonable to expect the existence of at least one point  $P$  with the above property. Surprisingly, a non equilateral isosceles triangle does *not* possess such a point. Our discussion examines the reason and considers interesting special cases.

Next, we look at the right angled triangle  $ABC$  in which  $\angle BAC = \pi/2$ . If  $AD$  is drawn perpendicular to  $BC$  (shown in Fig. 2), then it is well-known that  $AD^2 = BD \cdot DC$ . In fact, if  $D'$  is the mid point of  $BC$  of the same  $\triangle ABC$ , then also it trivially follows that  $AD'^2 = BD' \cdot D'C$  because  $D'$  is the center of the circle  $ABC$ . The second part of our discussion focuses on the characterization of  $\triangle ABC$  in which at least one point  $D$  exists on  $BC$  with the property just mentioned. Before we begin the discussion, let us recall the necessary background euclidean geometry results.

Analogien spielen bekanntlich eine wichtige Rolle beim Entdecken neuer Erkenntnisse in der Mathematik. An diese Tatsache knüpft der Autor in der nachfolgenden Arbeit an. Ausgehend von dem bekannten Ergebnis, dass für einen Punkt  $P$  auf dem Umkreis eines gleichseitigen Dreiecks  $ABC$  zwischen  $B$  und  $C$  die Beziehung  $AP = BP + PC$  besteht, wird in Umkehrung dazu nach notwendigen und hinreichenden Bedingungen an die Seiten eines beliebigen Dreiecks  $ABC$  gesucht, so dass ein Punkt  $P$  auf dem Umkreis des gegebenen Dreiecks zwischen  $B$  und  $C$  existiert, der der obigen Gleichung genügt. Ausgangspunkt für das zweite Beispiel ist der Höhensatz im rechtwinkligen Dreieck. Verallgemeinernd dazu wird nach notwendigen und hinreichenden Bedingungen an die Seiten eines beliebigen Dreiecks  $ABC$  gefragt, so dass ein Punkt  $D$  auf der Seite  $BC$  existiert, der die Beziehung  $AD^2 = BD \cdot DC$  erfüllt.


Fig. 1  $AP = BP + PC$ Fig. 2  $AD^2 = BD \cdot DC$  and  $AD'^2 = BD' \cdot D'C$ 

### Background material

We use the standard notation throughout our discussion. In  $\triangle ABC$ ,  $a, b, c$  denote either the sides or the lengths  $BC, CA, AB$  and  $A, B, C$  denote either the vertices or the measures of  $\angle BAC, \angle CBA, \angle ACB$ , respectively. We assume the reader's familiarity with the angle bisector theorem, properties of similar triangles, intersecting chords of a circle and so on. In addition, we need the result that is sometimes called Stewart's theorem. Therefore, we derive it below as an application of the cosine rule. This result is central to our discussion.

**Theorem 0** *Let  $D$  be a point on the side  $BC$  of  $\triangle ABC$ . If  $BD/DC = m/n$ , then*

$$AD^2 = \frac{mb^2 + nc^2}{m+n} - \frac{mn a^2}{(m+n)^2}. \quad (*)$$

Fig. 3  $\frac{BD}{DC} = \frac{m}{n}$ ,  $AD^2 = ?$ 

*Proof.* We refer to Fig. 3. If  $BD/DC = m/n$ , then by the property of equal ratios  $BD/(BD+DC) = m/(m+n)$ , so  $BD = ma/(m+n)$  and  $DC = na/(m+n)$ . We now apply the cosine rule for  $\triangle ABD$ . This gives

$$\begin{aligned} AD^2 &= AB^2 + BD^2 - 2 \cdot AB \cdot BD \cdot \cos(\angle ABD) \\ &= c^2 + \frac{m^2 a^2}{(m+n)^2} - 2 \cdot c \cdot \frac{ma}{m+n} \cdot \frac{c^2 + a^2 - b^2}{2ca}. \end{aligned}$$

The simplification (left to the reader) of the above expression leads to the result given in (\*).  $\square$

**Remark 1** The result (\*) is true for any position of the point  $D$  on  $BC$  that is within  $BC$  as in Fig. 3, on  $BC$  extended or  $CB$  extended. In such an extended case we should observe the usual sign convention, i.e.,  $BD > 0$ ,  $DC > 0$  if  $D$  is as in Fig. 3,  $BD > 0$ ,  $DC < 0$  if  $D$  is to the right of  $BC$  and  $BD < 0$ ,  $DC > 0$  if  $D$  is to the left of  $BC$ .

### The analogies

We first discuss the equilateral triangle analogy mentioned in the introduction. Theorem 1 shows that the sides of a  $\triangle ABC$  must be expressible in a certain form for the existence of a point  $P$  on the minor  $\widehat{BC}$  with the property  $AP = BP + PC$ . We will then deduce a number of corollaries. Except for Corollary 6, Theorem 1 and the Corollaries 1 to 5 are true for positive real numbers  $a, b, c, m$  and  $n$ .


**Theorem 1** *Let  $\triangle ABC$  be inscribed in a circle. Then a point  $P$  exists on the minor  $\widehat{BC}$  such that  $AP = BP + PC$ , if and only if the sides have the form  $a = mu^2 + nv^2$ ,  $b = u(mu + nv)$ ,  $c = v(mu + nv)$ . Here  $u$  and  $v$  are positive real numbers and  $m/n$  is the ratio in which  $AP$  divides the side  $BC$ .*

*Proof.* We refer to Fig. 4. From (\*) we have

$$AD^2 = \frac{mb^2 + nc^2}{m+n} - \frac{mna^2}{(m+n)^2}. \quad (1)$$

Since the triangles  $BDP$  and  $ADC$  are similar, we find that

$$\frac{BD}{AD} = \frac{BP}{AC}, \text{ i.e., } BP = \frac{mab}{(m+n)AD}.$$

Fig. 4 When  $AP = BP + PC$ ?

Likewise from the similar triangles  $CDP$  and  $ADB$  we find that

$$PC = \frac{nac}{(m+n)AD}.$$

Hence  $AP = BP + PC$ , if and only if

$$\begin{aligned} AD \cdot AP &= \frac{mab}{m+n} + \frac{nac}{m+n}, \quad \text{or} \\ AD^2 + AD \cdot DP &= \frac{a(mb+nc)}{(m+n)}. \end{aligned} \quad (2)$$

In Fig. 4,  $AP$  and  $BC$  are intersecting chords of the circle. Therefore,  $AD \cdot DP = BD \cdot DC = mna^2/(m+n)^2$ . Using the preceding result and substituting (1) into (2) gives

$$a(mb+nc) = mb^2 + nc^2.$$

Hence, the sides  $a, b, c$  of  $\triangle ABC$  must be expressible as  $a = (mb^2 + nc^2)/(mb + nc)$ ,  $b, c$ . We now take  $b = u, c = v$  as arbitrary positive real numbers and multiply the sides by  $(mu + nv)$ . This similarity transformation enlarges  $AP, BP, PC$  also by the same factor and hence the property  $AP = BP + PC$  still holds. This gives the characterization mentioned in Theorem 1.  $\square$

We now deduce a number of corollaries from Theorem 1. Corollary 1 gives another but simpler necessary and sufficient condition for  $\triangle ABC$  to have the desired point  $P$ .

**Corollary 1** *Let  $\triangle ABC$  be inscribed in a circle. A point  $P$  exists on the minor  $\widehat{BC}$  such that  $AP = BP + PC$ , if and only if the sides  $a, b, c$  satisfy the inequalities  $b > a > c$  or  $c > a > b$ .*

*Proof.* Theorem 1 tells us that the desired point  $P$  exists if and only if  $a = (mb^2 + nc^2)/(mb + nc)$ . This is equivalent to  $m/n = c(c-a)/b(a-b)$  or  $a = b = c$ , the latter being excluded here. Since  $AP$  intersects  $BC$  internally, the ratio  $m/n > 0$ . This is possible if and only if  $a, b, c$  satisfy  $b > a > c$  or  $c > a > b$  and the proof is complete.  $\square$

In fact, Corollary 1 provides greater insight into the subject under discussion as we see below.

**Corollary 2** A non equilateral isosceles triangle  $\Delta ABC$  does not contain a point  $P$  on the minor  $\widehat{BC}$  such that  $AP = BP + PC$ .

*Proof.* The hypothesis of Corollary 2 implies exactly one of  $a = b$ ,  $a = c$ ,  $b = c$ . This is neither  $b > a > c$  nor  $c > a > b$ . Hence,  $P$  does not exist. Alternatively, Theorem 1 shows that if one of the three above equalities holds then the other two *must* follow. This forces  $\Delta ABC$  to be equilateral contradicting the hypothesis of Corollary 2.  $\square$

Corollary 3 shows the uniqueness of the point  $P$ .

**Corollary 3** If the sides of  $\Delta ABC$  satisfy  $b > a > c$  or  $c > a > b$ , then there exists a unique point  $P$  on the minor  $\widehat{BC}$  such that  $AP = BP + PC$ .

*Proof.* In the proof of Corollary 1 we saw that  $m/n = c(c-a)/b(a-b)$ . Hence the ratio  $m/n$  is uniquely determined by  $a, b, c$ . Therefore, the point  $D$  on  $BC$  and hence  $P$  are unique, too.  $\square$

**Corollary 4** Suppose that a set of points  $P_i$  exists on the minor  $\widehat{BC}$  of  $\Delta ABC$  such that  $AP_i = BP_i + PC_i$ ,  $i = 1, 2, \dots$ . Then,  $\Delta ABC$  must be equilateral.

*Proof.* Since  $\Delta ABC$  is fixed,  $a, b, c$  remain constant but the ratio  $m/n$  varies as  $P$  moves over  $P_i$ . In other words, the equation  $mb(a-b) = nc(c-a)$  can hold for many pairs of values of  $m, n$ , if and only if  $a = b$  and  $c = a$  implying that  $\Delta ABC$  must be equilateral.  $\square$

**Corollary 5 (Hoppe triangles)** Let  $AP$  (Fig. 4) bisect  $\angle BAC$  of  $\Delta ABC$ . Then,  $AP = BP + PC$ , if and only if the sides  $b, a, c$  are in arithmetic progression.

*Proof.* The angle bisector theorem states that  $AP$ , i.e.,  $AD$  bisects  $\angle BAC$ , if and only if  $m/n = c/b$ . Hence,  $AP = BP + PC$ , if and only if  $a = (mb^2 + nc^2)/(mb + nc) = \frac{1}{2}(b+c)$ . This establishes the corollary. In fact, here  $AP = 2BP$  and  $AD = \frac{1}{2}\sqrt{3bc}$ .  $\square$

**Remark 2** Dickson in his monumental work, *History of the Theory of Numbers* [1] tells that Hoppe gives a complete discussion of Heron triangles (triangles with integer sides and area) in which the sides are in arithmetic progression. To honour Hoppe we call a triangle *Hoppe triangle* if the sides are in arithmetic progression. At this point we mention that there is a vast literature on Heron triangles and the reader is invited to refer to [3, 4, 6, 7] to get a glimpse of the many interesting Heron problems.

**Corollary 6** Suppose the elements  $a, b, c, m, n$  of  $\Delta ABC$  are positive integers. If  $AD$  is a median of  $\Delta ABC$ , then  $AP = BP + PC$ , if and only if  $AB + AC$  is either a square or twice a square.

*Proof.* Since  $AD$  is a median it follows that  $BD = DC$  and  $m = n = 1$ . Therefore,  $a = u^2 + v^2$ ,  $b = u(u+v)$ ,  $c = v(u+v)$ . Now,  $AB + AC = (u+v)^2$  is a (perfect) square. However, if  $u$  and  $v$  are both odd integers, then  $\gcd(a, b, c) = 2$ . The sides of the reduced (but similar) triangle will have  $AB + AC$  twice a square. Conversely, if  $AB + AC = w^2$  or  $2w^2$ , then we can always choose  $v = 1$ ,  $u = w - 1$  to get a desired  $\Delta ABC$ .  $\square$

**An example.** To make the argument clearer, suppose  $AB + AC = 36 = 6^2$ . Then,  $u + v = 6$ . If  $u = 5$ ,  $v = 1$ , then  $a = 26$ ,  $b = 30$ ,  $c = 6$  and  $\gcd(a, b, c) = 2$ . If we divide the sides by 2, then the reduced similar  $\Delta ABC$  has sides  $a = 13$ ,  $b = 15$ ,  $c = 3$  and  $AB + AC = 18 = 2(3)^2$  is twice a square. We remark that under a similarity transformation, the property  $AP = BP + PC$  is invariant.

Next, we consider the right angled triangle analogue mentioned in the introduction. Theorem 2 provides a characterization of  $\Delta ABC$  in which there is a point  $D$  on the side  $BC$  such that  $AD^2 = BD \cdot DC$ . In other words,  $AD$  is the geometric mean between  $BD$  and  $DC$ .

**Theorem 2** *In triangle  $ABC$  a point  $D$  on the side  $BC$  exists with the property  $AD^2 = BD \cdot DC$ , if and only if the sides  $a$ ,  $b$ ,  $c$  satisfy the inequalities  $\sqrt{2}a \geq b + c > a$ .*




Fig. 5 When is  $AD$  the geometric mean of  $BD$  and  $DC$ ?

*Proof.* We refer to Fig. 5. From  $(*)$  the equation  $AD^2 = BD \cdot DC$  is equivalent to

$$\frac{mb^2 + nc^2}{(m+n)} - \frac{mna^2}{(m+n)^2} = \frac{mna^2}{(m+n)^2}.$$

We rewrite the above equation as a quadratic in  $m/n$ . This yields

$$\left(\frac{m}{n}\right)^2 b^2 + (b^2 + c^2 - 2a^2) \left(\frac{m}{n}\right) + c^2 = 0$$

with the solutions

$$\frac{m}{n} = \frac{-b^2 - c^2 + 2a^2 \pm \sqrt{((b+c)^2 - 2a^2)((b-c)^2 - 2a^2)}}{2b^2}. \quad (3)$$

By the triangle inequality it follows that  $b - c < a$ . Hence,  $(b - c)^2 - 2a^2 \leq 0$ . To get a real root for  $m/n$  we must therefore have  $(b + c)^2 - 2a^2 \leq 0$ . That is  $\sqrt{2}a \geq b + c > a$ , the characterization asserted by the theorem.  $\square$

**Remark 3** In general, equation (3) yields two roots for  $m/n > 0$  and hence two positions for  $D$  on  $BC$  such that the cevian  $AD$  is the geometric mean of  $BD$  and  $DC$ .  $D$  exists uniquely when  $m/n$  is a double root. This happens, if and only if  $\sqrt{2}a = b + c$ . Corollary 7 tells more about such triangles.

**Corollary 7** *The point  $D$  on side  $BC$  of  $\Delta ABC$  exists uniquely with the property  $AD^2 = BD \cdot DC$ , if and only if  $AD$  bisects  $\angle BAC$ .*

*Proof.* In Remark 3 we saw that  $D$  exists uniquely, if and only if  $\sqrt{2}a = b + c$ . In this case, equation (3) shows that  $m/n = (-b^2 - c^2 + 2a^2)/2b^2 = c/b$ . The angle bisector theorem says that  $m/n$ , i.e.,  $BD/DC = c/b$ , if and only if  $AD$  bisects  $\angle BAC$ . This establishes Corollary 7. Moreover, in this case we find that  $AD = \sqrt{\frac{1}{2}bc}$ .  $\square$

Corollary 8 gives the length of the tangent at the vertex  $A$  to the circle  $ABC$  intercepted by the side  $BC$ .

**Corollary 8** *Let  $\Delta ABC$  be inscribed in a circle. Suppose the tangent at  $A$  meets the side  $BC$  (extended) at  $D$ . Then,  $AD = abc/|c^2 - b^2|$ .*

*Proof.* The tangent-secant property says that  $AD^2 = BD \cdot DC$ . If  $c > b$ , then  $D$  lies to the right of  $BC$  and if  $c < b$ , then  $D$  lies to the left of  $BC$ . If  $c = b$ , then  $AD$  is parallel to  $BC$  and undefined as a length. In either case,  $BD/DC = |c^2/b^2|$ . Hence,  $BD = c^2a/|c^2 - b^2|$ ,  $DC = b^2a/|c^2 - b^2|$  and the assertion of the corollary follows.  $\square$

## Conclusion

The reader witnessed an interesting discussion from the consideration of analogies of known results. Euclidean geometry has been studied for thousands of years and yet new results appear on the pages of mathematics journals around the world. We invite the reader to make a new contribution to the following lovely but unsolved problem in natural number solutions (as far as I am aware) on the equilateral triangle:

Let  $P$  be a point in the plane of an equilateral triangle  $ABC$ . Assume that  $PA = p$ ,  $PB = q$ ,  $PC = r$  are natural numbers. Determine the length  $s$  in natural numbers of a side of  $\Delta ABC$  in terms of  $p$ ,  $q$ ,  $r$ .

This is a famous problem with rich history. Reference [2] gives the details. The numbers  $p$ ,  $q$ ,  $r$ ,  $s$  are inter-related by the equation

$$3(p^4 + q^4 + r^4 + s^4) = (p^2 + q^2 + r^2 + s^2)^2.$$

The remarkable symmetry exhibited by the above equation shows that any three of the four values  $p$ ,  $q$ ,  $r$ ,  $s$  may be taken as the distances of the point  $P$  and the fourth one gives the side length of the equilateral triangle. The natural number solutions for  $p$ ,  $q$ ,  $r$ ,  $s$  are known in the special case when the point  $P$  lies on a side or on the circum circle of the equilateral  $\Delta ABC$ . These two instances are covered by setting  $p = q + r$  and then solving for  $s$ . Another special but new solution is found in [5]. Who is the lucky person who finds the general solution of the above problem/equation in natural numbers?

**Acknowledgement.** The author thanks Ms Christa Dobers who kindly produced the LaTeX version of my hand written paper.

## References

- [1] Dickson, L.E.: *History of the Theory of Numbers*. Vol. II, Chelsea, New York 1971, 171–201.
- [2] Mitrović, D.S. et al.: *Recent Advances in Geometric Inequalities*. Kluwer Academic Publishers, Dordrecht 1989, 391–392 (Chapter XIII).
- [3] Sastry, K.R.S.: Heron angles. *Mathematics and Comp. Education* 28 (1994), 24–35.
- [4] Sastry, K.R.S.: Heron triangles: a new perspective. *Aust. Math. Soc. Gazette* 26 (1999), 160–168.
- [5] Sastry, K.R.S.: Natural Number Solutions to  $3(p^4 + q^4 + r^4 + s^4) = (p^2 + q^2 + r^2 + s^2)^2$ . *Mathematics and Comp. Education* 34 (2000), 6–11.
- [6] Sastry, K.R.S.: Heron Triangles: A Gergonne-and-Median Perspective. *Forum Geometricorum* 1 (2001), 17–24.
- [7] Sastry, K.R.S.: Heron Ladders. *Math. Spectrum* 34 (2001/2002), 61–64.
- [8] Sastry, K.R.S.: Brahmagupta quadrilaterals: a description. *Crux Mathematicorum with MM* (to appear).

K.R.S. Sastry  
Jeevan Sandhya  
Doddakalsandra Post  
Raghuvana Halli  
Bangalore 560 062, India