

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 62 (2007)

Artikel: Also set-valued functions do not like iterative roots
Autor: Jarczyk, Witold / Zhang, Weinian
DOI: <https://doi.org/10.5169/seals-98912>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Also set-valued functions do not like iterative roots

Witold Jarczyk and Weinian Zhang

Witold Jarczyk obtained his PhD and habilitation from the Silesian University in 1983 and 1993, respectively. Presently, he is professor of mathematics at the University of Zielona Góra in Poland. His interests lie in functional equations and inequalities, iteration theory and discrete dynamical systems.

Weinian Zhang received his PhD from the Peking University in 1990. Presently, he is professor of mathematics at the Sichuan University in China. His interests lie in bifurcation theory of differential equations, iteration theory and dynamical systems.

1 Introduction

It seems that it is Ch. Babbage who first, yet at the beginning of the 19th century, wrote on iterative roots explicitly. Given a mapping $f : X \rightarrow X$ and a positive integer $n \geq 2$, the problem is to find a mapping $g : X \rightarrow X$ such that the n -th iterate of g , the composition g^n of n copies of g , is f , i.e., to solve the functional equation

$$g^n = f. \tag{1.1}$$

In [1] Babbage studied (1.1) for f being the identity mapping. After him a lot of results concerning the general case of (1.1) in various settings have been proved. Many of them can be found in the monographs [12] and [13] by M. Kuczma and M. Kuczma, B. Choczewski, R. Ger, respectively, as well as in the book [19] by Gy. Targonski. Some recent results have been presented in the survey papers [3] and [2].

Die Aufgabe, die n -te iterative Wurzel einer Abbildung $f : X \rightarrow X$ zu finden, besteht darin, eine Funktion $g : X \rightarrow X$ so zu bestimmen, dass $g^n = g \circ g \circ \dots \circ g = f$ (n -fache Hintereinanderausführung) gilt. Für dieses Problem sind sowohl kombinatorische, als auch analytische Resultate bekannt. So besitzt beispielsweise $f : [0, 1] \rightarrow [0, 1]$, gegeben durch $f(x) = 4x(1 - x)$, keine iterative Wurzel. Die Autoren untersuchen in dieser Arbeit das analoge Problem für mengenwertige Abbildungen $f : X \rightarrow 2^X$. Es zeigt sich, dass selbst Monotonie- und Stetigkeitsannahmen, die bei gewöhnlichen Funktionen Existenz von Wurzeln sicher stellen, hierfür in diesem Fall im allgemeinen nicht ausreichen.

The purely combinatorial paper [10] by R. Isaacs gave a description of solutions to (1.1) for an arbitrary bijection f . The case of general f was completely solved by G. Zimmermann, Ph.D. student of Targonski, in her not well-known doctoral thesis [22] (see also [17] by G. Riggert noticing that *Zimmermann* is the maiden name of Riggert).





Fig. 1: Hat function

Fig. 2: Parabola $y = 4x(1 - x)$

It turns out that even very simple and nice functions can have no roots. For instance, this is the case if f is the so-called *hat function*, i.e. $f(x) = \min\{2x, 2 - 2x\}$ for $x \in [0, 1]$ (see Fig. 1) or f is the celebrated *parabola* $y = 4x(1 - x)$ for $x \in [0, 1]$ (see Fig. 2). Of course, lack of roots for these functions can be deduced from Zimmermann's work. However, the reader surely can give a short and quite elementary argument in both cases. The above mentioned functions represent two important classes of mappings: piecewise monotone functions and polynomials. As follows, from [5], even in the class of piecewise monotone functions equation (1.1) leads to non-trivial questions. For f in this class sufficient conditions for nonexistence and existence of roots can be found in [21]. For polynomials the lack of roots is also a rather common phenomenon. A fundamental paper is [16] by R.E. Rice, B. Schweizer and A. Sklar, published in the *Monthly* almost 25 years ago. The answer to its title question "*When is $f(f(z)) = az^2 + bz + c$?*" is *never*. Similar results concerning some other polynomials can be found in [7] and [8]. Nonexistence of roots, both formal and holomorphic, was indicated by S. Bogatyi in his important article [6].

Difficulties appearing when solving equation (1.1), even in the class of continuous monotone self-mappings of an interval, have been enlightened in the crucial paper [9] by P.D. Humke and M. Laczkovich. Roughly speaking, they proved that the set of functions having a root is an analytic but non-Borel subset of the space $C([0, 1], \mathbb{R})$ endowed with the sup-norm. The papers [18] and [4] by K. Simon and A. Blokh, respectively, show that this set is small in $C([0, 1], [0, 1])$ both from the category (see [18, 4]) and measure-theoretical (cf. [18]) points of view. Nonexistence of roots is typical also for some regular functions (see [20]).

Recently some natural ideas of using set-valued functions have been examined (cf., for instance, [14, 15, 11]). One can consider replacing single-valued functions by set-valued functions in (1.1) both for f and g . It seems that up to now there are no notions leading to a satisfactory result in such a case.

In this paper we show that the phenomenon of lack of iterative roots appears also when studying some set-valued functions with exactly one value not being a singleton. Even imposing assumptions like continuity or strict monotonicity on the “single-valued parts” of such a set-valued function does not guarantee the existence of its square roots (see Example 2). This shows that maybe the situation for set-valued functions is even more sophisticated since those assumptions usually allow us to find roots in the single-valued case.

2 Main results

Given a set-valued function $f : X \rightarrow 2^Y$, the image $f(A)$ of a set $A \subset X$ is defined by

$$f(A) = \bigcup_{x \in A} f(x).$$

Then we can introduce the composition $g \circ f$ of set-valued functions $f : X \rightarrow 2^Y$ and $g : Y \rightarrow 2^Z$ by the familiar formula

$$(g \circ f)(x) = g(f(x)).$$

Clearly this operation is associative. So, for every positive integer n , we can define the n -th iterate of $g : X \rightarrow 2^X$ as the composition of n copies of g :

$$g^n = \underbrace{g \circ \dots \circ g}_{n\text{-times}}.$$

Consequently, the problem of looking for solutions $g : X \rightarrow 2^X$ to (1.1) for set-valued functions f is posed in a proper way.

Remark that if $g : X \rightarrow 2^X$ is an iterative root of $f : X \rightarrow 2^X$ then f and g commute, i.e. $f \circ g = g \circ f$. In fact, assume that $g^k = f$ for a positive integer k and fix an $x \in X$. If $z \in f(g(x))$ then $z \in f(y)$ for a $y \in g(x)$, that is, $z \in g^k(g(x)) = g(g^k(x)) = g(f(x))$. Conversely, if $z \in g(f(x))$ then $z \in g(y)$ for a $y \in f(x)$, so $z \in g(f(x)) = g(g^k(x)) = g^k(g(x)) = f(g(x))$.

In what follows, we consider X as an arbitrary set and let $\#A$ denote the cardinality of a subset $A \subset X$.

Proposition. Consider a set-valued function $f : X \rightarrow 2^X$ and let $g : X \rightarrow 2^X$ be its iterative square root. If there is a point $c \in X$ such that

- (i) $\#f(x) = 1$ for every $x \in X \setminus \{c\}$ and
- (ii) $f(x_0) = \{c\}$ for an $x_0 \in X$,

then $\#g(c) \leq 1$.

Proof. Suppose that

$$\#g(c) \geq 2. \tag{2.2}$$

It follows from (i) that g has non-void values only. Fix a $p \in g(x_0)$. Then

$$g(p) \subset g(g(x_0)) = f(x_0) = \{c\},$$

that is in fact $g(p) = \{c\}$, whence

$$f(p) = g(g(p)) = g(c).$$

Therefore, by (2.2) and (i), we get $p = c$. Thus, we have proved that $g(x_0) = \{c\}$, which gives

$$g(c) = g^2(x_0) = f(x_0) = \{c\},$$

a contradiction to (2.2). \square

Our main results are simple consequences of the proposition.

Theorem 1. *Let $f : X \rightarrow 2^X$. If there are a point $c \in X$ and a positive integer n such that (i) and (ii) hold,*

(iii) $\#f(c) > n$, and

(iv) $\#\{x \in X : f(x) = \{y\}\} \leq n$ for every $y \in X$,

then f has no iterative square roots.

Proof. Suppose that f has a square root $g : X \rightarrow 2^X$. By (i) and (iii) all the values of f and consequently of g are non-void.

Firstly, we claim that

$$\#g(x) \leq n \quad \text{for } x \in X \setminus \{c\}. \quad (2.3)$$

In order to see this, fix an $x \in X \setminus \{c\}$. Take any $v \in g(x)$. Since $f(x)$ is a singleton and

$$f(x) = g(g(x)) = \bigcup_{u \in g(x)} g(u),$$

for every $u \in g(x)$ we have $g(u) = g(v)$, whence $f(u) = f(v)$. This gives the inclusion

$$g(x) \subset \{u \in X : f(u) = f(v)\}. \quad (2.4)$$

If $g(x)$ is not a singleton then, according to (2.4) and (i), $f(v)$ is a singleton, whence using (2.4) again and (iv) we complete the proof of (2.3).

Since all the values of g are non-void, it follows from the proposition that $g(c) = \{u\}$ with a $u \in X$. Then $g(u) = g^2(c) = f(c)$, whence, by (iii), we have $\#g(u) > n$. So, $u \neq c$, which contradicts (2.3). \square

Theorem 2. *Let $f : X \rightarrow 2^X$. If there is a point $c \in X$ such that (i) and (ii) hold,*

(v) $\#f(c) > 1$, and

(vi) $c \in f(c)$,

then f has no iterative square roots.

Proof. Suppose that f has a square root $g : X \rightarrow 2^X$. By (i) and (v) all the values of g are non-void. Therefore, it follows from the proposition that $g(c) = \{u\}$ for a $u \in X$.

Then $g(u) = g^2(c) = f(c)$, whence, by (v), we have $\#g(u) > 1$, implying that $u \neq c$. On account of (v) the set $g(u)$ contains a point $v \in X \setminus \{c\}$. Moreover, according to (vi), $c \in f(c) = g(u)$. Therefore, since

$$g(v) \cup g(c) \subset g(g(u)) = f(u),$$

(i) gives $g(v) = g(c)$. Consequently, $f(v) = f(c)$, which contradicts (i) and (v). \square

3 Examples

1. Consider $f : [0, 1] \rightarrow 2^{[0,1]}$ given by

$$f(x) = \begin{cases} \frac{3}{2}x, & \text{if } x \in [0, \frac{1}{2}), \\ [\frac{1}{2}, \frac{3}{4}], & \text{if } x = \frac{1}{2}, \\ x, & \text{if } x \in (\frac{1}{2}, 1]. \end{cases}$$

Then assumptions (i)–(iv) in Theorem 1 are satisfied with $c = 1/2$ and $n = 2$. Consequently, f has no square root.

2. There are some properties, like e.g. strict monotonicity, continuity, lack of fixed points of f in the interior of its interval domain, which guarantee the existence of iterative roots of single-valued functions (cf., e.g., [12, Chap. XV] and [13, Chap. 11]). For set-valued functions the situation is more complicated, which can be seen by considering $f_1 : [0, 1] \rightarrow 2^{[0,1]}$ and $f_2 : [0, 1] \rightarrow 2^{[0,1]}$ defined by

$$f_1(x) = \begin{cases} \frac{5}{3}x, & \text{if } x \in [0, \frac{1}{2}), \\ [\frac{2}{3}, \frac{5}{6}], & \text{if } x = \frac{1}{2}, \\ \frac{2}{3}(x-1) + 1, & \text{if } x \in (\frac{1}{2}, 1], \end{cases}$$

$$f_2(x) = \begin{cases} \frac{4}{3}x, & \text{if } x \in [0, \frac{1}{2}), \\ [\frac{2}{3}, \frac{5}{6}], & \text{if } x = \frac{1}{2}, \\ \frac{1}{3}(x-1) + 1, & \text{if } x \in (\frac{1}{2}, 1], \end{cases}$$

respectively. Both of them are upper semicontinuous and have no fixed points in $(0, 1)$. Moreover, $f_1|_{[0, 1/2)}$ and $f_1|_{(1/2, 1]}$ are both strictly increasing and the (single-valued) function $f_1|_{[0, 1] \setminus \{1/2\}}$ is continuous. For f_2 we have even more: $f_2|_{[0, 1] \setminus \{1/2\}}$ is strictly increasing and continuous. Nevertheless, by Theorem 1, where we take $c = 1/2$ and $n = 3 - j$ for f_j ($j = 1, 2$), both f_1 and f_2 have no square roots. Observe also that $1/2 \notin f_1(1/2)$ and $1/2 \notin f_2(1/2)$, that is, condition (vi) is not satisfied. Consequently, Theorem 1 does not follow from Theorem 2.

3. In the case of f_3 as shown in Fig. 5 we have no roots again, as observed for $n = 4$.

4. The shape of the graph of f_4 (see Fig. 6), given by

$$f_4(x) = \begin{cases} \frac{1}{2}x, & \text{if } x \in [0, \frac{1}{2}), \\ [\frac{1}{4}, \frac{3}{4}], & \text{if } x = \frac{1}{2}, \\ \frac{1}{2}(x-1) + 1, & \text{if } x \in (\frac{1}{2}, 1], \end{cases}$$




Fig. 3: f_1 with $n = 2$




Fig. 4: f_2 with $n = 1$




Fig. 5: f_3 with $n = 4$




Fig. 6: f_4 has roots

is similar to the graph of f_2 , but f_4 has a square root. One can easily verify that $g : [0, 1] \rightarrow 2^{[0,1]}$, defined by

$$g(x) = \begin{cases} \frac{1}{\sqrt{2}}x, & \text{if } x \in [0, \frac{1}{2}), \\ [\frac{1}{2\sqrt{2}}, 1 - \frac{1}{2\sqrt{2}}], & \text{if } x = \frac{1}{2}, \\ \frac{1}{\sqrt{2}}(x - 1) + 1, & \text{if } x \in (\frac{1}{2}, 1], \end{cases}$$



satisfies $g^2 = f$. Observe, however, that Condition (ii) fails, where c has to be $1/2$.

5. Consider the set-valued functions $f_5 : [0, 1] \rightarrow 2^{[0,1]}$ and $f_6 : [0, 1] \rightarrow 2^{[0,1]}$ defined by

$$f_5(x) = \begin{cases} \frac{3}{2}x, & \text{if } x \in [0, \frac{1}{2}), \\ \{\frac{1}{2}, \frac{3}{4}\}, & \text{if } x = \frac{1}{2}, \\ x, & \text{if } x \in (\frac{1}{2}, 1], \end{cases}$$

$$f_6(x) = \begin{cases} \frac{3}{2}x, & \text{if } x \in [0, \frac{1}{2}), \\ [\frac{1}{2}, \frac{3}{4}], & \text{if } x = \frac{1}{2}, \\ \frac{1}{2}, & \text{if } x \in (\frac{1}{2}, \frac{3}{4}], \\ 2(x-1)+1, & \text{if } x \in (\frac{3}{4}, 1], \end{cases}$$

respectively. Condition (iii) is not satisfied by f_5 since $c = 1/2$, $n = 2$ and $\#f_5(c) = 2$. For f_6 condition (iii) is not satisfied because $c = 1/2$, $n = \aleph_0$ and $\#f_6(c) = \aleph_0$. However, they both satisfy (v) and (vi). Theorem 2 shows that none of them has a square root. Consequently, this also implies that Theorem 2 does not follow from Theorem 1.

Fig. 7: f_5 with $n = 2$ Fig. 8: f_6 with $n = \infty$

Acknowledgment

The work was supported by the University of Zielona Góra, NSFC#10471101, NOYF-B#10428104 and SRFDP#20050610003.

References

- [1] Babbage, Ch.: Essay towards the calculus of functions. *Philosoph. Transact.* (1815), 389–423; Essay towards the calculus of functions, II. *Ibid.* (1816), 179–256.
- [2] Baron, K.: *Recent results in the theory of functional equations in a single variable*. Seminar LV 15 (2003), 1–16, <http://www.mathematik.uni-karlsruhe.de/~sem1v>
- [3] Baron, K.; Jarczyk, W.: Recent results on functional equations in a single variable, perspectives and open problems. *Aequationes Math.* 61 (2001), 1–48.
- [4] Blokh, A.M.: The set of all iterates is nowhere dense in $C([0, 1], [0, 1])$. *Trans. Amer. Math. Soc.* 333 (1992), 787–798.
- [5] Blokh, A.M.; Coven, E.; Misiurewicz, M.; Nitecki, Z.: Roots of continuous piecewise monotone maps of an interval. *Acta Math. Univ. Comenian. (N.S.)* 60 (1991), 3–10.
- [6] Bogatyı, S.: On the nonexistence of iterative roots. *Topology Appl.* (1997), 97–123.
- [7] Bronshtein, E.M.: *On an iterative square root of a quadratic trinomial* [Russian]. *Geometric problems in the theory of functions and sets* [Russian]. 24–27, Kalinin. Gos. Univ., Kalinin 1989.
- [8] Choczewski, B.; Kuczma, M.: *On iterative roots of polynomials*. European Conference on Iteration Theory (Lisbon, 1991), 59–67, World Sci. Publishing, Singapore 1992.

- [9] Humke, P.D.; Laczkovich, M.: The Borel structure of iterates of continuous functions. *Proc. Edinburgh Math. Soc.*(2) 32 (1989), 483–493.
- [10] Isaacs, R.: Iterates of fractional order. *Canad. J. Math.* 2 (1950), 409–416.
- [11] Jarczyk, W.; Powierża, T.: On the smallest set-valued iterative roots of bijections. *Dynamical Systems and Functional Equations* (Murcia, 2000). *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* 13 (2003), 1889–1893.
- [12] Kuczma, M.: *Functional equations in a single variable*. Monografie Mat. 46, Polish Scientific Publishers, Warszawa 1968.
- [13] Kuczma, M.; Choczewski, B.; Ger, R.: *Iterative functional equations*. Encyclopedia of Mathematics and Its Applications 32, Cambridge University Press, Cambridge 1990.
- [14] Powierża, T.: Strong set-valued iterative roots. *Grazer Math. Ber.* 344 (2001), 51–56.
- [15] Powierża, T.: Higher order set-valued iterative roots of bijections. *Publ. Math. Debrecen* 61 (2002), 315–324.
- [16] Rice, R.E.; Schweizer, B.; Sklar, A.: When is $f(f(z)) = az^2 + bz + c$? *Amer. Math. Monthly* 87 (1980), 252–263.
- [17] Riggert, G.: n -te iterative Wurzeln von beliebigen Abbildungen. *Aequationes Math.* 14 (1976), 208.
- [18] Simon, K.: Some dual statements concerning Wiener measure and Baire category. *Proc. Amer. Math. Soc.* 106 (1989), 455–463.
- [19] Targonski, Gy.: *Topics in iteration theory*. Vandenhoeck and Ruprecht, Göttingen 1981.
- [20] Zhang, W.: A generic property of globally smooth iterative roots. *Sci. China Ser. A* 38 (1995), 267–272.
- [21] Zhang, W.: PM functions, their characteristic intervals and iterative roots. *Ann. Polon. Math.* 65 (1997), 119–128.
- [22] Zimmermann, G.: *Über die Existenz iterativer Wurzeln von Abbildungen*. Doctoral Dissertation, Marburg/Lahn 1978.

Witold Jarczyk

Faculty of Mathematics, Computer Science and Econometrics

University of Zielona Góra

Szafrana 4a

PL–65-516, Zielona Góra, Poland

e-mail: wjarczyk@uz.zgora.pl

Weinian Zhang (Corresponding author)

Department of Mathematics

Sichuan University

Chengdu

Sichuan 610064, P.R. China

e-mail: matzwn@126.com or matzwn@sohu.com