

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 62 (2007)

Artikel: A new family of almost identities
Autor: Maze, Gérard / Minder, Lorenz
DOI: <https://doi.org/10.5169/seals-98913>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A new family of almost identities

Gérard Maze and Lorenz Minder

Gérard Maze received his Ph.D. from the University of Notre Dame in 2002. He currently holds a postdoctoral position at the University of Zurich. His research interests lie in the field of Applied Algebra.

Lorenz Minder is currently a Ph.D. student at the École Polytechnique Fédérale de Lausanne (EPFL). His research interests cover cryptography and coding theory.

1 Introduction

It is well-known that a class of “almost integers” can be found using the theory of modular functions, and a few spectacular examples are given by Ramanujan [5]. They can be generated using some amazing properties of the j -function. Some of the numbers which are close approximations of integers are $\exp(\pi\sqrt{163})$ (sometimes known as Ramanujan’s constant), $\exp(\pi\sqrt{37})$ and $\exp(\pi\sqrt{58})$. These irrationals come close to an integer as follows:

$$\begin{aligned}\exp(\pi\sqrt{37}) &= 199148648 - 0.219 \dots \cdot 10^{-4}, \\ \exp(\pi\sqrt{58}) &= 24591257752 - 0.177 \dots \cdot 10^{-6}, \\ \exp(\pi\sqrt{163}) &= 262537412640768744 - 0.749 \dots \cdot 10^{-12}.\end{aligned}$$

Another surprising result comes from the average length of a segment in an isosceles right triangle with catheti of unit length. If l is this average length, then

$$l = \frac{1}{30} \left(2 + 4\sqrt{2} + (4 + \sqrt{2}) \sinh^{-1}(1) \right) = 0.4142933026 \dots = (\sqrt{2} - 1) - 0.8 \dots \cdot 10^{-4}.$$

„Almost identities“ sind mathematische Ausdrücke, die gegen „schöne“ Zahlen zu konvergieren scheinen, stattdessen aber – mit oft schier unglaublicher Präzision – danebentreffen. Verblüffende almost identities sind faszinierend. Noch faszinierender ist aber meist die Frage, die almost identities unweigerlich aufwerfen, nämlich: Ist es Zufall oder gibt es einen tieferen Grund? In dieser Arbeit stellen die Autoren eine Familie von almost identities vor und geben anschliessend eine Analyse, die das Phänomen in diesen Fällen erklärt.

Such astonishing non-equalities are usually called almost identities or non-identities. Many examples of such unexpected behaviour are known [6]. The four examples above are however different in essence: the first three come from a deep property of a complex mathematical object (the j -function) and the last has a good chance of being a genuine arithmetical coincidence.

A natural question that comes to mind in presence of such a non-identity is therefore whether or not the phenomenon is purely coincidental, or comes from a more subtle process. For instance, in the equation

$$e^\pi - \pi = 19.999099979 \dots,$$

it is not clear at all whether the almost identity pops up from a deep connection between e and π or just because the expression *happens* to be close to 20.

Recently, J.M. Borwein and P.B. Borwein discovered several families of almost identities [2], leading to a systematic study of such phenomena. These were based on mathematical concepts that lead to clear explanations. Among the non-identities studied by these authors, let us mention the following striking example:

$$\sum_{k=-\infty}^{\infty} \frac{1}{10^{(k/100)^2}} \cong 100 \sqrt{\frac{\pi}{\ln(10)}},$$

correct to at least 18,000 digits. In this situation, the almost identity is *not* a coincidence. From the same viewpoint, let us mention as well the sequence

$$h_n = \frac{n!}{2(\ln(2))^{n+1}},$$

for $1 \leq n \leq 17$, discovered by D. Hickerson. These numbers are close to integers due to the fact that the above quotient is the dominant term in an infinite series whose sum is the number of possible outcomes of a race between n people (where ties are allowed). See [6] for the exact expression of these numbers. Here, once again, no coincidence.

While we were studying the function

$$f(x) = \sum_{k=1}^{\infty} \frac{1}{1 + 2^k x}, \quad x \in]0, 1],$$

that appears in the analysis of the complexity of the binary gcd algorithm, we came to find a new family of almost identities. Let us define the real numbers u_n as follows:

$$u_n := \ln(2) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{(2^{k/2} + 2^{-k/2})^n}, \quad n \in \mathbb{N} \setminus \{0\}.$$

The following equalities show the very strange behaviour of the almost identities generated

by the sequence $\{u_n\}$:

$$\begin{aligned}
 u_1 &= \pi + 0.53 \dots \cdot 10^{-11} \\
 u_2 &= 1 + 0.48 \dots \cdot 10^{-10} \\
 u_3 &= \frac{\pi}{2^3} + 0.22 \dots \cdot 10^{-9} \\
 u_4 &= \frac{1}{6} + 0.67 \dots \cdot 10^{-9} \\
 u_5 &= \frac{3\pi}{2^7} + 0.15 \dots \cdot 10^{-8} \\
 u_6 &= \frac{1}{30} + 0.29 \dots \cdot 10^{-8} \\
 \vdots &\quad \vdots \quad \vdots
 \end{aligned}$$

This article presents an explanation of this phenomenon. We study the cases with $n = 1$ and $n = 2$ by using the theory of Mellin transforms and leave the remaining cases together with the recurrence relation (1.1) below to the extended version of this article [4]. We also present a generalization of the phenomenon, leading to, e.g., the almost identity

$$\ln(4) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{2^{-k} + 2^k} = \pi + 0.82 \dots \cdot 10^{-5}.$$

The proof of the recurrence relation

$$u_n = \left(\frac{1}{4} \cdot \frac{n-2}{n-1} \right) u_{n-2} + r_n \quad (1.1)$$

is not given in this article, but it can be found in the extended version of this work [4], where also the explicit values of r_n satisfying $0 < r_n \leq r_{10} = 0.7227399 \dots \cdot 10^{-8}$, $\forall n \in \mathbb{N}$, are described.

In this article, we will use the notation $f(x) \sim_a g(x)$ for equivalent functions in a neighbourhood of a and $\log_2 x$ for the logarithm in base 2 of x . Also, the set \mathbb{N} is considered to contain the integer 0 in the sequel.

2 Preliminaries

The first two cases on our list are

$$u_1 = \ln(2) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{2^{-k/2} + 2^{k/2}} = \pi + 0.53 \dots \cdot 10^{-11}$$

and

$$u_2 = \ln(2) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{(2^{-k/2} + 2^{k/2})^2} = 1 + 0.48 \dots \cdot 10^{-10}.$$

Let us define the complex functions g_1 and g_2 as

$$g_1(x) = -2 \cdot (\arctan(\sqrt{x}) - \pi/2) \quad \text{and} \quad g_2(x) = \frac{1}{1+x}, \quad \Re x > 0,$$

as well as the functions G_1 and G_2 defined as

$$G_n(x) = \sum_{k=1}^{\infty} g_n(2^k x), \quad \Re x > 0, \quad n = 1, 2.$$

The convergence of G_1 is justified by the fact that in a neighbourhood of $+\infty$ we have

$$\arctan t - \pi/2 = - \int_t^{\infty} \frac{1}{1+v^2} dv = - \int_t^{\infty} \left(\frac{1}{v^2} - \frac{1}{v^4} + \frac{1}{v^6} + \dots \right) dv = O(1/t).$$

The following equalities are justified because G_1 and G_2 converge uniformly on compact subsets of their domains, and therefore, the derivative can be interchanged with the sum. Therefore,

$$\begin{aligned} \lim_{m \rightarrow \infty} \frac{d}{du} [G_n(2^{-u})] \Big|_{u=m} &= \lim_{m \rightarrow \infty} \sum_{k=1}^{\infty} \frac{d}{du} [g_n(2^{k-u})] \Big|_{u=m} \\ &= \lim_{m \rightarrow \infty} \ln(2) \cdot \sum_{k=1}^{\infty} \frac{(2^{(k-m)})^{n/2}}{(1+2^{(k-m)})^n} \\ &= \lim_{m \rightarrow \infty} \ln(2) \cdot \sum_{k=1}^{\infty} \frac{1}{(2^{-(k-m)/2} + 2^{(k-m)/2})^n} \\ &= u_n, \end{aligned} \tag{2.1}$$

where the limit is understood with $m \in \mathbb{N}$. The game plan is then to express the functions G_1 and G_2 in a completely different manner in order to compute these limits. The keystone of this process is the Mellin transform [3]. Recall that the Mellin transform of a locally Lebesgue integrable function $f(x)$ over $]0, \infty[$ is the function

$$f^*(s) = \int_0^{\infty} f(x) x^{s-1} dx.$$

The conditions $f(x) \sim_0 O(x^u)$ and $f(x) \sim_{\infty} O(x^v)$, with $u > v$ guarantee that $f^*(s)$ exists in the strip $-u < \Re s < -v$. Mellin's inversion formula [3, p. 13] states that if f is continuous and $c \in]-u, -v[$, then

$$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} f^*(s) x^{-s} ds,$$

and in a neighbourhood of 0, we have

$$f(x) = \sum_{\Re s_l < c} \text{Res}(f^*(s) x^{-s}, s_l),$$

where the summation is over the poles s_l of the function $f^*(s)x^{-s}$ whose real part is strictly smaller than c .

Let $g(x)$ be a locally Lebesgue integrable function over $]0, \infty[$, $f(x) = \sum_{k=1}^{\infty} g(2^k x)$, and suppose that the convergence is uniform in $]0, \infty[$. Then,

$$\begin{aligned} f^*(s) &= \int_0^{\infty} \sum_{k=1}^{\infty} g(2^k x) x^{s-1} dx \\ &= \sum_{k=1}^{\infty} \int_0^{\infty} g(y) y^{s-1} 2^{-ks} dy \\ &= \frac{g^*(s)}{2^s - 1}. \end{aligned} \tag{2.2}$$

3 The case $n = 1$

Proposition 1. *For $x > 0$, we have*

$$G_1(x) = -\frac{\pi}{2} - \pi \log_2(x) + \sqrt{x} S_1(x) - \sum_{k=1}^{\infty} \frac{\sin(2k\pi \log_2(x))}{k \cdot \cosh(2k\pi^2 / \ln(2))},$$

where $S_1(x)$ is a power series in x , which converges in $[0, 1[$.

Proof. As announced earlier, the idea is to use Mellin transforms in a back and forth process to reveal another expression of G_1 . Using (2.2), we can write

$$G_1^*(s) = \frac{g_1^*(s)}{2^s - 1}. \tag{3.1}$$

In order to compute g_1^* , recall that in a neighbourhood of $+\infty$ we have $\arctan t - \pi/2 = O(1/t)$. So, we can perform an integration by parts, as long as $\Re s \in]0, 1/2[$:

$$\begin{aligned} g_1^*(s) &= -2 \int_0^{\infty} (\arctan(\sqrt{x}) - \pi/2) x^{s-1} dx \\ &= -2 \cdot \left[(\arctan(\sqrt{x}) - \pi/2) \cdot \frac{x^s}{s} \Big|_0^{\infty} - \frac{1}{2s} \int_0^{\infty} \frac{x^{s-1/2}}{1+x} dx \right] \\ &= \frac{1}{s} \int_0^{\infty} \frac{x^{s-1/2}}{1+x} dx \\ &= \frac{\pi}{s \cos \pi s}. \end{aligned}$$

The last equality comes from the relation

$$\int_0^{\infty} \frac{x^{s-1}}{1+x} dx = \frac{\pi}{\sin \pi s}.$$

Using Mellin's inversion formula with $c = 1/4$ and (3.1), we get

$$\begin{aligned} G_1(x) &= \frac{1}{2\pi i} \int_{1/4-i\infty}^{1/4+i\infty} \left(\frac{\pi}{s \cos \pi s} \right) \frac{x^{-s}}{2^s - 1} ds \\ &= \sum_{\Re s_l < 1/4} \text{Res} \left(\left(\frac{\pi}{s \cos \pi s} \right) \frac{x^{-s}}{2^s - 1}, s_l \right). \end{aligned}$$

The poles of the function $\left(\frac{\pi}{s \cos \pi s} \right) \frac{x^{-s}}{2^s - 1}$ can be partitioned as follows:

- i) $s = 0$ is a pole of order two,
- ii) the real simple poles $-1/2 + k$, $k \in \mathbb{Z}$,
- iii) the imaginary simple poles $2k\pi i / \ln(2)$, $k \in \mathbb{Z} \setminus \{0\}$.

The residues are then

$$\begin{aligned} -\pi \log_2(x) - \frac{\pi}{2} &\quad \text{at } s = 0, \\ -\frac{(-2)^{k+2}}{(1+2k)(2^{k+1} - \sqrt{2})} \sqrt{x} x^k &\quad \text{at } s = -1/2 - k, \quad k \in \mathbb{N}, \\ \frac{1}{2i} \cdot \frac{\exp(-2k\pi i \log_2(x))}{k \cdot \cosh(2k\pi^2 / \ln(2))} &\quad \text{at } s = 2k\pi i / \ln(2), \quad k \in \mathbb{Z} \setminus \{0\}, \end{aligned}$$

and the above sum becomes

$$\begin{aligned} G_1(x) &= -\frac{\pi}{2} - \pi \log_2(x) + \sum_{k=0}^{\infty} \frac{(-2)^{k+2}}{(1+2k)(-2^{k+1} + \sqrt{2})} \sqrt{x} x^k \\ &\quad - \sum_{k=1}^{\infty} \frac{\sin(2k\pi \log_2(x))}{k \cdot \cosh(2k\pi^2 / \ln(2))}, \end{aligned}$$

which proves the proposition. \square

Corollary 2. $u_1 = \pi + \sum_{k=1}^{\infty} \frac{2\pi}{\cosh(2k\pi^2 / \ln(2))}$.

Proof. Based on (2.1), we have

$$\begin{aligned} u_1 &= \lim_{m \rightarrow \infty} \frac{d}{du} [G_1(2^{-u})] \Big|_{u=m} \\ &= \pi + \lim_{u \rightarrow \infty} \left[e^{-u/2} S_1(e^{-u}) \right]' + \sum_{k=1}^{\infty} \frac{2\pi}{\cosh(2k\pi^2 / \ln(2))} \end{aligned}$$

and the last limit being equal to zero, the corollary is proven. \square

The case $n = 1$ is then settled since the sum on the right-hand side of the equality of Corollary 2 is in fact small:

$$u_1 - \pi = \sum_{k=1}^{\infty} \frac{2\pi}{\cosh(2k\pi^2/\ln(2))} = 0.538914478 \dots \cdot 10^{-11}.$$

4 The case $n = 2$

We can compute u_2 in a similar way as we computed u_1 . Note that once u_1 and u_2 are computed, the recurrence relation (1.1) gives the complete sequence $\{u_i\}$.

Proposition 3. *For $x > 0$, we have*

$$G_2(x) = -\frac{1}{2} - \log_2(x) + S_2(x) - \frac{2\pi}{\ln(2)} \sum_{k=1}^{\infty} \frac{\sin(2k\pi \log_2(x))}{\sinh(2k\pi^2/\ln(2))},$$

where $S_2(x)$ is a power series in x , converging in $[0, 1[$ such that $S_2(x) = 0$.

Proof. The proof follows the same lines as in the first case. First,

$$g_2^*(s) = \int_0^{\infty} \frac{x^{s-1}}{1+x} dx = \frac{\pi}{\sin \pi s},$$

and thus, once again based on (2.2) and (3.1), we have

$$\begin{aligned} G_2(x) &= \int_{1/2-i\infty}^{1/2+i\infty} G_2^*(s)x^{-s} ds \\ &= \int_{1/2-i\infty}^{1/2+i\infty} \left(\frac{\pi}{\sin \pi s} \right) \frac{x^{-s}}{2^s - 1} ds \\ &= \sum_{\Re s_l < 1/2} \text{Res} \left(\left(\frac{\pi}{\sin \pi s} \right) \frac{x^{-s}}{2^s - 1}, s_l \right). \end{aligned}$$

The poles of the function can be partitioned as follows:

- i) $s = 0$ is a pole of order two,
- ii) the real simple poles $k, k \in \mathbb{Z} \setminus \{0\}$,
- iii) the imaginary simple poles $2k\pi i / \ln(2), k \in \mathbb{Z} \setminus \{0\}$.

The residues are then

$$\begin{aligned} -\log_2(x) - \frac{1}{2} &\quad \text{at } s = 0, \\ -\frac{(-2)^k}{2^k - 1} x^k &\quad \text{at } s = -k, \quad k = 1, 2, 3, \dots, \\ \frac{\pi}{i} \cdot \frac{\exp(-2k\pi i \log_2(x))}{\ln(2) \cdot \sinh(2k\pi^2/\ln(2))} &\quad \text{at } s = 2k\pi i / \ln(2), \quad k \in \mathbb{Z} \setminus \{0\}. \end{aligned}$$

The new expression of G_2 is therefore

$$G_2(x) = -\frac{1}{2} - \log_2(x) - \sum_{k=1}^{\infty} \frac{(-2)^k}{2^k - 1} x^k - \frac{2\pi}{\ln(2)} \sum_{k=1}^{\infty} \frac{\sin(2k\pi \log_2(x))}{\sinh(2k\pi^2/\ln(2))}.$$

This concludes the proof. \square

Corollary 4. $u_2 = 1 + \frac{2\pi}{\ln(2)} \sum_{k=1}^{\infty} \frac{2k\pi}{\sinh(2k\pi^2/\ln(2))}$.

Proof. We use the same trick as in Corollary 2:

$$\begin{aligned} u_2 &= \lim_{m \rightarrow \infty} \frac{d}{du} [G_2(2^{-u})] \Big|_{u=m} \\ &= 1 + \lim_{u \rightarrow \infty} [S_2(e^{-u})'] + \frac{2\pi}{\ln(2)} \sum_{k=1}^{\infty} \frac{2k\pi}{\sinh(2k\pi^2/\ln(2))} \end{aligned}$$

and the limit being equal to zero, the corollary is proven. \square

Once again, this shows why the number u_2 is almost an integer. Indeed, the sum on the right-hand side is fairly small:

$$u_2 - 1 = \frac{2\pi}{\ln(2)} \sum_{k=1}^{\infty} \frac{2k\pi}{\sinh(2k\pi^2/\ln(2))} = 0.4885108992 \dots \cdot 10^{-10}.$$

5 Final remarks

Having found the roots of the mystery related to the non-equalities $u_1 \neq \pi$ and $u_2 \neq 1$, let us mention that the same tools used so far lead to a complete understanding of the non-equalities concerning u_3, u_4, \dots and why they are so close to “good arithmetic numbers”. The interested reader can find the details in the extended version of this article [4].

We end this article with the following remark: Our arguments do not depend on the presence of 2 in the denominator of

$$\frac{1}{(2^{-k/2} + 2^{k/2})^n}.$$

One could argue that any sum of the type

$$\ln(m) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{(m^{-k/2} + m^{k/2})^n}$$

has the potential to lie close to \mathbb{Q} or $\pi\mathbb{Q}$ depending on the parity of n . As a matter of fact, we have, for example,

$$\begin{aligned} \ln(4) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{2^{-k} + 2^k} &= \pi + 0.82 \dots \cdot 10^{-5}, \\ \ln(9) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{3^{-k} + 3^k} &= \pi + 0.15 \dots \cdot 10^{-2}, \\ \ln(4) \cdot \sum_{k=-\infty}^{\infty} \frac{1}{(2^{-k} + 2^k)^2} &= 1 + 0.37 \dots \cdot 10^{-4}. \end{aligned}$$

In fact, it can be proven that the choice $m = 2$ is the best one can do in order to maximize the resemblance with elements in $\mathbb{Q} \cup \pi\mathbb{Q}$, see [4].

References

- [1] Andrews, G.E.; Askey, R.; Roy, R.: *Special functions*. Encyclopedia of Mathematics and its Applications, 71, Cambridge University Press, 1999.
- [2] Borwein, J.M.; Borwein, P.B.: Strange Series and High Precision Fraud. *Amer. Math. Monthly* 99 (1992), 622–640.
- [3] Flajolet, P.; Gourdon, X.; Dumas, P.: Mellin transforms and asymptotics: harmonic sums. *Theoret. Comput. Sci. (Special volume on mathematical analysis of algorithms)* 144 (1995), 3–58.
- [4] Maze, G.; Minder, L.: A New Family of Almost Identities (extended version).
<http://www.arxiv.org/abs/math.GM/0409014>.
- [5] Ramanujan, S.: Modular Equations and Approximations to π . *Quart. J. Pure Appl. Math.* 45 (1914/15), 350–372.
- [6] Weisstein, E.W.: Almost Integer. From MathWorld – A Wolfram Web Resource, at
<http://mathworld.wolfram.com/AlmostInteger.html>.

Gérard Maze
 Mathematics Institute
 University of Zurich
 CH-8057 Zurich, Switzerland
 e-mail: gmaze@math.unizh.ch

Lorenz Minder
 Laboratoire de Mathématiques Algorithmiques
 École Polytechnique Fédérale de Lausanne
 CH-1015 Lausanne, Switzerland
 e-mail: lorenz.minder@epfl.ch